



Endoscopy After the COVID-19 Pandemic—What Will Be Different?

Rashid N. Lui, MBChB, FRCP^{1,2}
Raymond S. Y. Tang, MD^{1,2}
Philip W. Y. Chiu, MD, FRCSEd^{2,3,*}

Address

¹Division of Gastroenterology and Hepatology, Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Hong Kong, SAR, China

²Endoscopy Centre, Prince of Wales Hospital, The Chinese University of Hong Kong, Hong Kong, SAR, China

Email: philipchiu@surgery.cuhk.edu.hk

³Division of Upper GI and Metabolic Surgery, Department of Surgery, The Chinese University of Hong Kong, Hong Kong, SAR, China

Published online: 26 January 2022

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2022

This article is part of the Topical Collection on *Endoscopy*

Abstract

Purpose of Review This paper reviews the latest literature regarding the impact of COVID on endoscopy service provision.

Recent Findings Endoscopy has been shown to be largely safe when appropriate infection prevention and control measures are in place. Endoscopy training and education has been profoundly affected though novel training models to overcome this have been developed. Proper handling of delayed or cancelled procedures is of utmost importance to minimize delays in diagnosis and treatment of diseases such as cancer. Adoption of new technologies such as non-endoscopy alternatives and telehealth may be a viable alternative to minimize infection risks.

Summary This pandemic has led to tangible differences in how we provide endoscopy service in the future. Future research focusing on better risk stratification of patients who need endoscopy, validating novel endoscopy training models, and adopting new technologies are urgently needed to support these changes in the post-pandemic world.

Introduction

The COVID (coronavirus disease) pandemic is still ravaging in many parts of the world and has led to an unprecedented impact on healthcare systems globally. Initially, many parts of the world had to stop elective endoscopies due to the risk of potential infections, lack of personal protective equipment (PPE), and redeployment of healthcare professionals (HCPs) to COVID-related patient services. [1•, 2•, 3•, 4–10] After the initial phase, many centers resumed limited endoscopy services with the implementation of stringent infection prevention and control (IPC) policies such as universal masking for patients, pre-endoscopy symptoms screening and COVID testing, and using high-level personal protective equipment for upper endoscopies as they are potentially aerosol-generating. [11, 12] Thankfully, with widespread implementation of these measures, no large nosocomial outbreak of the virus has been reported so far. As the pandemic waxed and waned, many authorities issued guidance

on how to safely and effectively restart routine clinical services. [13, 14, 15•, 16–18] The rapid development of vaccines and the widespread implementation of vaccination programs worldwide was originally seen as a game changer, [19–23] but this has led to the surges of highly infectious COVID variants. [24•, 25] Hopefully, with efforts to strengthen containment and to further increase the vaccination uptake rates, even in younger individuals and children, [26] the pandemic can be controlled to a certain degree. Regardless of when this pandemic will really end, it is evident that COVID has profoundly impacted healthcare systems worldwide, including the delivery of endoscopy services. It is evident that some of the changes adopted will likely be staying with us in perpetuity. In this review, we provide a summary of the literature and some of our insights on the future of endoscopy in the post-COVID era (Table 1).

Procedural Workflow for Endoscopy

Pre-endoscopy

Infection-Minimized Pathways

Hayee et al. [27] have suggested separating hospitals into “hot” and “cold” aka COVID-minimized sites to stratify the risk of endoscopy to patients, taking into consideration individual patient risks such as those with cancer, on immunosuppression etc. A study of more than 6200 patients by the SCOTS project group from the UK showed that providing such a COVID-minimized environment can offer a clear road map for performing safe endoscopy during the recovery phase of the pandemic. [28•] In the future even after the COVID pandemic has settled, these infection-minimized pathways may still have a role to reduce the risk of nosocomial outbreaks and provide better protection to patients at high risk of acquiring infections.

Universal Screening and Testing

Regardless of the infectious agent, routine screening of patients for FTOCC (fever, travel history, occupational history, clustering history, and contact history) will likely be here to stay as a simple and easily implementable safeguard to protect other patients and HCPs.

Table 1 Changes regarding endoscopy service provision after the COVID pandemic

Procedural workflow for endoscopy

Pre-endoscopy	<ul style="list-style-type: none"> - Infection minimized pathways - Universal screening and testing - Patient PPE
Endoscopy	- Adequate PPE for endoscopists (N95 respirators or equivalent for upper gastrointestinal endoscopies)
Post-endoscopy	<ul style="list-style-type: none"> - Segregation of recovery areas - Phone follow-up of patients
Lasting impact	
Endoscopy training and education	<ul style="list-style-type: none"> - Web-based lectures - Video-based education - National and international webinars - Simulation-based training - Objective skill assessment tools - Leveraging social media platforms for alternative means of education and training
Endoscopist well-being	<ul style="list-style-type: none"> - Enhance mental health support for endoscopists - Identify structural causes of burnout and anxiety in clinical programs
Proper handling of delayed or cancelled procedures	<ul style="list-style-type: none"> - Resume endoscopy services such as CRC screening - Logbook of all subjects who had endoscopy delayed or cancelled due to the pandemic - Alternative risk stratification tools such as incorporating FIT
Adoption of non-endoscopy alternatives	<ul style="list-style-type: none"> - Adopt new technologies such as capsule endoscopy, CT colonography - Telehealth
Aftermath	
Preparedness for future pandemics	<ul style="list-style-type: none"> - Stockpile PPE, medical equipment, and crucial medications - Develop surge capacity within healthcare systems - Establish pandemic preparedness committees to develop clinical frameworks for future outbreaks - Strengthen collaboration and liaison with local public health officials - Develop better predictive models

Abbreviations: *PPE* personal protective equipment; *CRC* colorectal cancer; *FIT* fecal immunochemical test; *CT* computed tomography

In the future, if another pandemic arises, universal testing for the detection of this hypothetical pathogen will likely be implemented. A trend to move from laboratory-based tests to point-of-care tests will also likely be present, thus allowing for the development of decentralized, rapid, sensitive, and low-cost diagnostics which can be widely implemented as screening tests. [29] With the widespread vaccination of HCPs and the general population, stringent screening measures that were required prior to elective endoscopy will likely be relaxed similar to the current pandemic. For example, the American Gastroenterological Association updated their guidance recently with recommendations against the routine pre-procedure testing for SARS-CoV-2 in patients scheduled to undergo endoscopy. [30]

Patient PPE

It is highly likely that standard IPC measures will be fundamentally changed after the pandemic. Asking patients to wear facemasks at all times and adhere to social distancing policies where appropriate, especially for patients that are potentially infectious sources, will likely be continued in the near future.

Endoscopy*Aerosol-Generating Procedures*

Our understanding of which procedures are considered aerosol-generating procedures (AGPs) has enhanced significantly during this pandemic. There is mounting evidence that upper GI (gastrointestinal) endoscopy including esophagogastroduodenoscopy (EGD), endoscopic retrograde cholangiopancreatography (ERCP), and endoscopic ultrasound should be considered as AGPs, and endoscopists should wear enhanced PPE with N95/FFP3 respirators or equivalent. [11, 31] In our center, the routine use of a continuous dental sucker has already been widely implemented for all upper GI endoscopies which has been shown to significantly reduce the amount of aerosols, [11] with this practice likely to be continued in the future in the post-COVID era.

The risk for lower GI endoscopy is less conclusive. Although viral RNA has been detected in stool samples of COVID patients, whether this confers a risk for fecal–oral transmission similar to prior outbreaks related to toilet fumes during the SARS (severe acute respiratory syndrome) pandemic has not been well established, [32] though theoretically the risk of transmission via aerosols or fomites is largely similar between SARS-CoV-1 and SARS-CoV-2. [33] Regardless, many authorities consider lower GI endoscopy as potentially AGP as the procedure usually lasts longer than other endoscopies thus increasing exposure time to the endoscopist, requiring air insufflation, with the risk of the patient passing flatus.

Availability of Negative-Pressure Facilities and Appropriate Building Engineering Controls

Endoscopy for patients with suspected or confirmed COVID should be performed in negative pressure facilities for further protection of endoscopy HCPs and other patients. In the future, if another respiratory pathogen causes large-scale infections, these policies will likely be re-enacted. Hospitals where the aforementioned negative pressure rooms are not available should consider adding these facilities during redevelopment as part of a preparedness plan for future pandemics.

The COVID pandemic has led to renewed interest in the potential role of air change rates as an infection control strategy. [34] Appropriate building engineering controls including sufficient and effective ventilation, particle filtration and air disinfection, avoiding air recirculation, and avoiding over-crowding have all been suggested. [35]

For better or for worse, the days of endoscopists performing procedures without facemasks will likely be over. Despite evidence from Johnston et al. [36] showing that endoscopy poses a clear risk for transmission of potentially infectious biologic samples, it took a global pandemic to truly increase the uptake rate of basic IPC measures. Thankfully, the risk of COVID transmission is considered low if adherence to strict preventive measures is in place. [37]

Post-endoscopy

Segregation of Recovery Areas

Many centers have implemented workflows where patients with suspected or confirmed COVID are cohorted in separate locations throughout their patient journey, including a segregation of recovery areas. [4] In the future, facing a theoretical new pathogen, these specific workflows can and should be re-implemented to minimize the risk of cross-infection and allow for easier contact tracing.

Phone Follow-Up

Some authorities have suggested conducting a routine phone follow-up in 7 to 14 days after endoscopy to enquire if there are any new symptoms or new diagnoses of COVID. [7]. Studies have shown that client satisfaction is improved and failure-to-attend rates are lowered with phone consultations, and that it can effectively minimize in-person consultations [38] Regardless of the infectious agent, this will likely be considered good clinical practice and may become commonplace in the future.

At the end of the day, the safety of endoscopy for both patients and HCPs is of utmost importance. Stringent measures that are carefully adhered to before, during, and after endoscopy can mitigate the risk of infection. The silver lining is that this is an opportunity to introduce new care models and enhance our preparedness for future pandemics. [39]

Lasting Impact and Aftermath

Endoscopy Training and Education

The COVID pandemic has had a profound impact on endoscopy procedure volumes [40] and endoscopy training. [41•, 42–45] Concerns have been raised regarding difficulties encountered by trainees to meet pre-COVID

training targets given the many restrictions to endoscopy at the height of the pandemic. Some authors have suggested that a new multimodal educational model should be adopted, encompassing web-based didactic learning, simulation-based training, and skill-based goal setting for competency assessment. [46]

Distance learning including local web-based lectures, video-based education with interaction, and national and international webinars or lectures, primarily via gastrointestinal societies, has become an important educational resource. [47] These measures will likely continue in the long term and provide the essential basics of endoscopy to trainees in a safe manner, with minimal utilization of scarce PPE resources, and will likely enhance patient safety.

Simulation-based training has improved substantially in recent years. More sophisticated endoscopy simulators may incorporate elements of both mechanical simulation and augmented reality [46]/virtual reality. [48] In more resource-limited regions, low-cost training models have also been developed [49]

Future endoscopy training may focus less on the number of procedures performed, and shift to the use of objective skill assessment tools in the clinical setting to monitor the learning curves of trainees or fellows. [46] Though there may be pros and cons of such a competency-based curriculum, it offers a unique hybrid learning opportunity for continued training during pandemics.

Increasingly, social media platforms such as Twitter offer an alternative means of education and training. Although in the past these channels were considered unstructured, an increasing rigor to such education modalities has been seen [50] with some resources such as #MondayNightIBD offering continuing medical education credits to participants. [51]

The pandemic likely accelerated the uptake of many of these educational measures. Whether these innovations can stand the test of time would depend on whether changes in policy, training requirements, monetary investment, trainer buy-in and time are present. [52] It is also encouraging to note that with adaptive mechanisms, procedure volumes for trainees and/or fellows have returned near to baseline without an ongoing impact on endoscopy training. [53] Time will tell whether these new educational and training models are on par with traditional methods that are mainly based on procedure numbers.

Endoscopist Well-Being

The well-being has been an issue for further study during this pandemic. In one multinational survey by Pawlak et al. [41•], more than 70% of endoscopy trainees reported concerns that the COVID could prolong their training, with more than 50% reporting symptoms of anxiety and approximately

18% reporting symptoms of burnout. In a more recent study of Southeast Asian gastroenterologists, more than 70% were still significantly affected by the pandemic and burnout was also common. Safeguards for mental health were found to be suboptimal with more than 50% of gastroenterologists being unaware of or did not have access to mental health support. [54] The pandemic has laid bare the lack of psychosocial and mental health support in many regions worldwide and renewed efforts to make workplace improvements for these important issues for endoscopists are urgently needed.

Proper Handling of Delayed or Cancelled Procedures

In a large retrospective study from Hong Kong, despite a drastic decrease in hospitalizations for gastrointestinal ailments and reduction in endoscopy volumes, data seems to suggest that this did not affect emergency endoscopy or surgeries, nor lead to excessive mortality. [55] However, emerging data suggests that there are tangible, detrimental effects of the pandemic on diagnosing new cancers such as gastric and colorectal cancers (CRCs), and that delays in endoscopy may potentially lead to upstaging of cancer staging. [56] In the USA, it is estimated that there is a screening deficit for 9.4 million individuals that is [57•] associated with the pandemic. In another study, an estimated increase of more than 15% of avoidable CRC deaths in England was to be expected as a result of these diagnostic delays. [58] Encouragingly, although the COVID pandemic did lead to a sustained reduction in the number of people referred, diagnosed, and treated for CRC in England, by October 2020, care pathway targets had returned to 2019 levels suggesting that screening can be re-implemented with modifications to usual practice. [59] Restarting cancer screening activities in the post-COVID era will be difficult and will likely require a well-coordinated effort to more proactively engage the community, encourage apparently healthy individuals to return to routine health care, and reorganize clinical services to minimize backlogs. A logbook for subjects who should have been previously but had their procedures delayed or cancelled should be made available in all endoscopy centers to ensure that these patients are followed up and offered timely endoscopy. Using CRC screening programs as an example, Basu et al. [60] discussed the possibility of allowing flexibility in screening programs such as switching to fecal immunochemical testing (FIT) in place of endoscopy-based screening. The application of a FIT test would allow the use of the mail system to send and return test kits to make it more convenient for subjects and to maintain a reasonably high participation rate. The authors also opined that the opportunity should be taken to replace qualitative FIT with quantitative FIT as an example of de-implementation of existing non-evidence-based

practices. [60] Facing an unprecedented outbreak with a global shortage of PPE, it was understandable that elective endoscopies were delayed or cancelled initially. However, this may also serve as a lesson if future pandemics arise where the real-world consequences of delaying diagnoses of time-dependent diseases such as malignancy need to be balanced carefully with infection prevention and control policies.

Adoption of Non-Endoscopy Alternatives

Capsule endoscopy possesses unique advantages such as single use, low risk of cross-infection, excellent tolerance with low risk of aerosol generation, minimal medical staff requirement, and separation of examination and reading/reporting which are particularly advantageous during the COVID pandemic. [61] These advantages will likely persist after the pandemic, and with wider availability and increased experience by clinicians, this may become a viable screening option for selected gastrointestinal conditions.

Another non-endoscopy modality that is non-invasive with negligible risk of aerosol generation and hence infective risk would be computed tomography (CT) colonography. These technologies are not new, but the previous uptake has been slow and they have yet to be fully integrated into current diagnostic algorithms. In light of the pandemic, utilizing these alternative methods may be a safe and effective way to offer suitable diagnostic tests and risk stratify patients, reserving invasive endoscopy which bears a higher infective risk for selected patients.

Telehealth

The pandemic has catalyzed the emergence of telehealth as a viable alternative to in-person clinic visits has been adopted in many clinical practices. [62] There is potential that current healthcare provision can be restructured to incorporate telehealth for improved patient convenience, satisfaction, and clinical efficacy. Patients with stable disease, and those who prefer to continue to use this modality should continue to have telehealth consultations arranged. In the future, it is possible that a certain proportion of clinical services and dedicated sessions are repurposed as telehealth visits. Though short-term data is encouraging, whether adopting telehealth is equally safe and effective in the long run requires further research.



Fig. 1 Full gear of personal protection equipment (including hair net, face shield/goggles, N95 respirators or equivalent, water)

Preparedness for Future Pandemics

Other measures to enhance preparedness for future outbreaks include a formal policy of stockpiling PPE, medical equipment, and crucial medications, developing surge capacity within healthcare systems, improving access to telehealth services, establishing pandemic preparedness committees to develop clinical frameworks for future outbreaks, strengthening collaboration and liaison with local public health officials, and developing better predictive models to enhance resource allocation [63] (Figs. 1 and 2).




Fig. 2 Workflow at Endoscopy Center during the COVID pandemic

Conclusion

Though far from over, the COVID pandemic will not be the only infectious disease facing humanity and endoscopists in the near future. The challenges we have faced have provided a litmus test on how endoscopy services can be safely and effectively delivered during pandemics. The experience gained and lessons learned from COVID will be crucial when we encounter inevitable challenges in the future.

Declarations

Conflict of Interest

Rashid N. Lui declares that he has no conflict of interest. Raymond S. Y. Tang declares that he has no conflict of interest. Philip W. Y. Chiu declares that he has no conflict of interest.

Human and Animal Rights and Informed Consent

This article does not contain any studies with human or animal subjects performed by any of the authors.

References

Papers of particular interest, published recently, have been highlighted as: • Of importance

1. • Sultan S, Lim JK, Altayar O, Davitkov P, Feuerstein JD, Siddique SM, Falck-Ytter Y, El-Serag HB, AGA. AGA Institute rapid recommendations for gastrointestinal procedures during the COVID-19 pandemic. *Gastroenterology* [Internet]. 2020 Mar 31; Available from: <https://doi.org/10.1053/j.gastro.2020.03.072>. (North American recommendations for endoscopy during COVID).
2. • Gralnek IM, Hassan C, Beilenhoff U, Antonelli G, Ebigbo A, Pellisè M, Arvanitakis M, Bhandari P, Bisschops R, Van Hooft JE, Kaminski MF, Triantafyllou K, Webster G, Pohl H, Dunkley I, Fehrke B, Gazic M, Gjergjek T, Maasen S, Waagenes W, de Pater M, Ponchon T, Siersema PD, Messmann H, Dinis-Ribeiro M. ESGE and ESGENA position statement on gastrointestinal endoscopy and the COVID-19 pandemic. *Endoscopy* [Internet]. 2020 Apr 17; <https://doi.org/10.1055/a-1155-6229>. (European recommendations for endoscopy during COVID).
3. • Chiu PWY, Ng SC, Inoue H, Reddy DN, Ling Hu E, Cho JY, Ho LK, Hewett DG, Chiu H-M, Rerknimitr R, Wang H-P, Ho SH, Seo DW, Goh K-L, Tajiri H, Kitano S, Chan FKL. Practice of endoscopy during COVID-19 pandemic: position statements of the Asian Pacific Society for Digestive Endoscopy (APSDE-COVID statements). *Gut*. 2020;69(6):991–6 (Asia-Pacific recommendations for endoscopy during COVID)
4. Lui RN, Wong SH, Sánchez-Luna SA, Pellino G, Bollipo S, Wong M-Y, Chiu PWY, Sung JY. Overview of guidance for endoscopy during the coronavirus disease 2019 pandemic. *J Gastroenterol Hepatol*. 2020;35(5):749–59.
5. Tse F, Borgaonkar M, Leontiadis GI. COVID-19: advice from the Canadian Association of Gastroenterology for endoscopy facilities, as of March 16, 2020. *J Can Assoc Gastroenterol*. 2020;3(3):147–9.
6. Soetikno R, Teoh AYB, Kaltenbach T, Lau JYW, Asokkumar R, Cabral-Prodigalidad P, Shergill A. Considerations in performing endoscopy during the COVID-19 pandemic. *Gastrointest Endosc*. 2020;92(1):176–83.
7. Repici A, Maselli R, Colombo M, Gabbiadini R, Spadaccini M, Anderloni A, Carrara S, Fugazza A, Di Leo M, Galtieri PA, Pellegatta G, Ferrara EC, Azzolini E, Lagioia M. Coronavirus (COVID-19) outbreak: what the department of endoscopy should know. *Gastrointest Endosc* [Internet]. 2020 Mar 14; Available from: <https://doi.org/10.1016/j.gie.2020.03.019>.
8. Irisawa A, Furuta T, Matsumoto T, Kawai T, Inaba T, Kanno A, Katanuma A, Kawahara Y, Matsuda K, Mizukami K, Otsuka T, Yasuda I, Tanaka S, Fujimoto K, Fukuda S, Iishi H, Igarashi Y, Inui K, Ueki T, Ogata H, Kato M, Shiotani A, Higuchi K, Fujita N, Murakami K, Yamamoto H, Ito T, Okazaki K, Kitagawa Y, Mine T, Tajiri H, Inoue H. Gastrointestinal endoscopy in the era of the acute pandemic of COVID-19: recommendations by Japan Gastroenterological Endoscopy Society (Issued on April 9 th , 2020.). *Dig Endosc* [Internet]. 2020 Apr 26; Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/den.13703>.
9. Furfaro F, Vuitton L, Fiorino G, Koch S, Allocca M, Gilardi D, Zilli A, D'Amico F, Radice S, Chevaux J-B, Schaefer M, Chaussade S, Danese S, Peyrin-Biroulet L. SFED recommendations for IBD endoscopy during COVID-19 pandemic: Italian and French experience. *Nat Rev Gastroenterol Hepatol*. 2020;17(8):507–16.
10. Crespo J, Fernández Carrillo C, Iruzubieta P, Hernández-Conde M, Rasines L, Jorquera F, Albillos A, Bañares R, Mora P, Fernández Vázquez I, Hernández-Guerra M, Turnes J, Calleja JL, COVID-19 SEPD/AEEH Group. Massive impact of coronavirus disease 2019 pandemic on gastroenterology and hepatology departments and doctors in Spain. *J Gastroenterol Hepatol*. 2021;36(6):1627–33.
11. Chan SM, Ma TW, Ka-Chun Chong M, Chan DL, Ng EKW, Chiu PWY. A proof of concept study: esophagogastroduodenoscopy is an aerosol-generating procedure and continuous oral suction during the procedure reduces the amount of aerosol generated. *Gastroenterology* [Internet]. 2020 Jul 7; Available from: <https://doi.org/10.1053/j.gastro.2020.07.002>.
12. Leung NHL, Chu DKW, Shiu EYC, Chan K-H, McDevitt JJ, Hau BJP, Yen H-L, Li Y, Ip DKM, Peiris JSM, Seto W-H, Leung GM, Milton DK, Cowling BJ. Respiratory virus shedding in exhaled breath and efficacy of face masks. *Nat Med*. 2020;26(5):676–80.
13. Rees CJ, East JE, Oppong K, Veitch A, McAlindon M, Anderson J, Hayee B, Edwards C, McKinlay A, Penman I. Restarting gastrointestinal endoscopy in the deceleration and early recovery phases of COVID-19 pandemic: guidance from the British Society of Gastroenterology. *Clin Med*. 2020;20(4):352–8.
14. Marín-Gabriel JC, de Santiago ER. AEG-SEED position paper for the resumption of endoscopic activity after the peak phase of the COVID-19 pandemic. *Gastroenterología y Hepatología (English Edition)* [Internet]. 2020 Jul 10; Available from: <http://www.sciencedirect.com/science/article/pii/S2444382420301188>.

15. • Bhandari P, Subramaniam S, Bourke MJ, Alkandari A, Chiu PWY, Brown JF, Keswani RN, Bisschops R, Hassan C, Raju GS, Muthusamy VR, Sethi A, May GR, Albéniz E, Bruno M, Kaminski MF, Alkhatry M, Almadi M, Ibrahim M, Emura F, Moura E, Navarrete C, Wulfson A, Khor C, Ponnudurai R, Inoue H, Saito Y, Yahagi N, Kashin S, Nikonov E, Yu H, Maydeo AP, Reddy DN, Wallace MB, Pochapin MB, Rösch T, Sharma P, Repici A. Recovery of endoscopy services in the era of COVID-19: recommendations from an international Delphi consensus. *Gut*. 2020;69(11):1915–24
(Expert views on how to safely resume endoscopy services during COVID)
16. Ménard C, Waschke K, Tse F, Borgaonkar M, Forbes N, Barkun A, Martel M. COVID-19: framework for the resumption of endoscopic activities from the Canadian Association of Gastroenterology. *Journal of the Canadian Association of Gastroenterology* [Internet]. [cited 2020 Aug 4]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7337808/>.
17. Antonelli G, Karsensten JG, Bhat P, Ijoma U, Osuagwu C, Desalegn H, Abera H, Guy C, Vilmann P, Dinis-Ribeiro M, Ponchon T, Sabbagh LC, Pausawasdi N, Makharia G, Hassan C, Veitch A, Aabakken L, Ebigbo A, Endoscopy and COVID-19 Cascade Working Group, European Society of Gastrointestinal Endoscopy, World Endoscopy Organization and World Gastroenterology Organization. Resuming endoscopy during COVID-19 pandemic: ESGE, WEO and WGO Joint Cascade Guideline for Resource Limited Settings. *Endosc Int Open*. 2021;9(4):E543–51.
18. Zhang S, Wu X, Feng Y, Wang Q, Jiang Q, Guo T, Wu D, Xu T, Li R, Tang S-J, Yang A. Resuming gastrointestinal endoscopy post-COVID-19 peak: focus on the guidance from international and national societies. *J Gastroenterol Hepatol*. 2021;36(2):526–33.
19. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, Perez JL, Pérez Marc G, Moreira ED, Zerbini C, Bailey R, Swanson KA, Roychoudhury S, Koury K, Li P, Kalina WV, Cooper D, Frenck RW Jr, Hammitt LL, Türeci Ö, Nell H, Schaefer A, Ünal S, Tresnan DB, Mather S, Dormitzer PR, Şahin U, Jansen KU, Gruber WC, C4591001 Clinical Trial Group. Safety and efficacy of the BNT162b2 mRNA COVID-19 vaccine. *N Engl J Med*. 2020;383(27):2603–15.
20. Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S, Novak R, Diemert D, Spector SA, Rouphael N, Creech CB, McGettigan J, Khetan S, Segall N, Solis J, Brosz A, Fierro C, Schwartz H, Neuzil K, Corey L, Gilbert P, Janes H, Follmann D, Marovich M, Masciola J, Polakowski L, Ledgerwood J, Graham BS, Bennett H, Pajon R, Knightly C, Leav B, Deng W, Zhou H, Han S, Ivarsson M, Miller J, Zaks T, COVE Study Group. Efficacy and safety of the mRNA-1273 SARS-CoV-2 vaccine. *N Engl J Med*. 2021;384(5):403–16.
21. Voysey M, Clemens SAC, Madhi SA, Weckx LY, Folegatti PM, Aley PK, Angus B, Baillie VL, Barnabas SL, Bhorat QE, Bibi S, Briner C, Cicconi P, Collins AM, Colin-Jones R, Cutland CL, Darton TC, Dheda K, Duncan CJA, Emary KRW, Ewer KJ, Fairlie L, Faust SN, Feng S, Ferreira DM, Finn A, Goodman AL, Green CM, Green CA, Heath PT, Hill C, Hill H, Hirsch I, Hodgson SHC, Izu A, Jackson S, Jenkin D, Joe CCD, Kerridge S, Koen A, Kwatra G, Lazarus R, Lawrie AM, Lelliott A, Libri V, Lillie PJ, Mallory R, Mendes AVA, Milan EP, Minassian AM, McGregor A, Morrison H, Mujadidi YF, Nana A, O'Reilly PJ, Padayachee SD, Pittella A, Plested E, Pollock KM, Ramasamy MN, Rhead S, Schwarzbold AV, Singh N, Smith A, Song R, Snape MD, Sprinz E, Sutherland RK, Tarrant R, Thomson EC, Török ME, Toshner M, Turner DPJ, Vekemans J, Villafana TL, Watson MEE, Williams CJ, Douglas AD, Hill AVS, Lambe T, Gilbert SC, Pollard AJ, Oxford COVID Vaccine Trial Group. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *Lancet*. 2021;397(10269):99–111.
22. Logunov DY, Dolzhikova IV, Shcheblyakov DV, Tukhvatulin AI, Zubkova OV, Dzharullaeva AS, Kovyrshina AV, Lubenets NL, Grousova DM, Erokhova AS, Botikov AG, Izhaeva FM, Popova O, Ozharovskaya TA, Esmagambetov IB, Favorskaya IA, Zrelkin DI, Voronina DV, Shcherbinin DN, Semikhin AS, Simakova YV, Tokarskaya EA, Egorova DA, Shmarov MM, Nikitenko NA, Gushchin VA, Smolyarchuk EA, Zyrjanov SK, Borisevich SV, Naroditsky BS, Gintsburg AL, Gam-COVID-Vac Vaccine Trial Group. Safety and efficacy of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine: an interim analysis of a randomised controlled phase 3 trial in Russia. *Lancet*. 2021;397(10275):671–81.
23. Jara A, Undurraga EA, González C, Paredes F, Fontecilla T, Jara G, Pizarro A, Acevedo J, Leo K, Leon F, Sans C, Leighton P, Suárez P, García-Escorza H, Araos R. Effectiveness of an inactivated SARS-CoV-2 vaccine in Chile. *N Engl J Med*. 2021;385(10):875–84.
24. • Hacısuleyman E, Hale C, Saito Y, Blachere NE, Bergh M, Conlon EG, Schaefer-Babajew DJ, DaSilva J, Muecksch F, Gaebler C, Lifton R, Nussenzweig MC, Hatzioannou T, Bieniasz PD, Darnell RB. Vaccine breakthrough infections with SARS-CoV-2 variants. *N Engl J Med*. 2021;384(23):2212–8
(Highlighting the problem posed by SARS-CoV-2 variants even in vaccinated populations)
25. Abdool Karim SS, de Oliveira T. New SARS-CoV-2 variants - clinical, public health, and vaccine implications. *N Engl J Med*. 2021;384(19):1866–8.
26. Han B, Song Y, Li C, Yang W, Ma Q, Jiang Z, Li M, Lian X, Jiao W, Wang L, Shu Q, Wu Z, Zhao Y, Li Q, Gao Q. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine (CoronaVac) in healthy children and adolescents: a double-blind, randomised, controlled, phase 1/2 clinical trial.

- Lancet Infect Dis [Internet]. 2021 Jun 28; Available from: [https://doi.org/10.1016/S1473-3099\(21\)00319-4](https://doi.org/10.1016/S1473-3099(21)00319-4).
27. Hayee B 'hussain, Thoufееq M, Rees CJ, Penman I, East J. Safely restarting GI endoscopy in the era of COVID-19. *Gut* [Internet]. 2020 Jun 5; Available from: <https://doi.org/10.1136/gutjnl-2020-321688>.
 28. • Hayee B 'hussain, SCOTS project group, East J, Rees CJ, Penman I. Multicentre prospective study of COVID-19 transmission following outpatient GI endoscopy in the UK. *Gut*. 2021;70(5):825–8.
- (Safe models that may be adopted for resuming endoscopy during COVID).
29. Valera E, Jankelow A, Lim J, Kindratenko V, Ganguli A, White K, Kumar J, Bashir R. COVID-19 point-of-care diagnostics: present and future. *ACS Nano*. 2021;15(5):7899–906.
 30. Sultan S, Siddique SM, Singh S, Altayar O, Caliendo AM, Davitkov P, Feuerstein JD, Kaul V, Lim JK, Mustafa RA, Falck-Ytter Y, Inadomi JM, American Gastroenterological Association. AGA rapid review and guideline for SARS-CoV2 testing and endoscopy post-vaccination: 2021 update. *Gastroenterology*. 2021;161(3):1011–29. e11.
 31. Sagami R, Nishikiori H, Sato T, Tsuji H, Ono M, Togo K, Fukuda K, Okamoto K, Ogawa R, Mizukami K, Okimoto T, Kodama M, Amano Y, Murakami K. Aerosols produced by upper gastrointestinal endoscopy: a quantitative evaluation. *Am J Gastroenterol*. 2021;116(1):202–5.
 32. Wong SH, Lui RN, Sung JJ. COVID-19 and the digestive system. *J Gastroenterol Hepatol* [Internet]. 2020 Mar 25; Available from: <https://doi.org/10.1111/jgh.15047>.
 33. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, Tamin A, Harcourt JL, Thornburg NJ, Gerber SI, Lloyd-Smith JO, de Wit E, Munster VJ. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med*. 2020;382(16):1564–7.
 34. Allen JG, Ibrahim AM. Indoor air changes and potential implications for SARS-CoV-2 transmission. *JAMA*. 2021;325(20):2112–3.
 35. Morawska L, Tang JW, Bahnfleth W, Bluyssen PM, Boerstra A, Buonanno G, Cao J, Dancer S, Floto A, Franchimon F, Haworth C, Hogeling J, Isaxon C, Jimenez JL, Kurnitski J, Li Y, Loomans M, Marks G, Marr LC, Mazzarella L, Melikov AK, Miller S, Milton DK, Nazaroff W, Nielsen PV, Noakes C, Peccia J, Querol X, Sekhar C, Seppänen O, Tanabe S-I, Tellier R, Tham KW, Wargoeki P, Wierzbicka A, Yao M. How can airborne transmission of COVID-19 indoors be minimised? *Environ Int*. 2020;142:105832.
 36. Johnston ER, Habib-Bein N, Dueker JM, Quiroz B, Corsaro E, Ambrogio M, Kingsley M, Papachristou GI, Kreiss C, Khalid A. Risk of bacterial exposure to the endoscopist's face during endoscopy. *Gastrointest Endosc*. 2019;89(4):818–24.
 37. Repici A, Aragona G, Cengia G, Cantù P, Spadacini M, Maselli R, Carrara S, Anderloni A, Fugazza A, Pace F, Rösch T, ITALIAN GI-COVID19 Working Group. Low risk of COVID-19 transmission in GI endoscopy. *Gut* [Internet]. 2020 Apr 22; Available from: <https://doi.org/10.1136/gutjnl-2020-321341>.
 38. Zorron Cheng Tao Pu L, Singh G, Rajadurai A, Terbah R, De Silva R, Vaughan R, Efthymiou M, Chandran S. Benefits of phone consultation for endoscopy-related clinics in the COVID-19 pandemic. *J Gastroenterol Hepatol*. 2021;36(4):1064–80.
 39. Lui RN. Safety in endoscopy for patients and healthcare workers during the COVID-19 pandemic. *Tech Innov Gastrointest Endosc* [Internet]. 2020 Oct 18; Available from: <https://doi.org/10.1016/j.tige.2020.10.004>.
 40. Srinivasan S, Sundaram S, Emura F, Reddy N, Faigel DO, Repici A, Parasa S, Sharma P. Ongoing global impact of the COVID-19 pandemic on endoscopy: a subsequent international survey of 121 centers from 35 countries (BRIEF REPORT). *Gastroenterology* [Internet]. 2021 Sep 23; Available from: <https://doi.org/10.1053/j.gastro.2021.09.042>.
 41. • Pawlak KM, Kral J, Khan R, Amin S, Bilal M, Lui RN, Sandhu DS, Hashim A, Bollipo S, Charabaty A, de-Madaria E, Rodríguez-Parra AF, Sánchez-Luna SA, Żorniak M, Walsh CM, Grover SC, Siau K. Impact of COVID-19 on endoscopy trainees: an international survey. *Gastrointest Endosc*. 2020;92(4):925–35.
- (International study showing the significant impact of COVID on endoscopy training)
42. Siau K, Iacucci M, Dunckley P, Penman I, EndoTrain Survey Collaborators. The impact of COVID-19 on gastrointestinal endoscopy training in the United Kingdom. *Gastroenterology*. 2020;159(4):1582–5. e3.
 43. Kumar S, Prenner S, Kochman ML. The impact of COVID-19 on endoscopic training. *Am J Gastroenterol*. 2020;115(7):1142–3.
 44. Khan R, Tandon P, Scaffidi MA, Bishay K, Pawlak KM, Kral J, Amin S, Bilal M, Lui RN, Sandhu DS, Hashim A, Bollipo S, Charabaty A, de-Madaria E, Rodríguez-Parra AF, Sánchez-Luna SA, Żorniak M, Siau K, Walsh CM, Grover SC. COVID-19 and Canadian gastroenterology trainees. *J Can Assoc Gastroenterol*. 2021;4(3): 156.
 45. Vignesh S, Butt AS, Alborai M, Martins BC, Piscoya A, Tran QT, Yew DTM, Ghazanfar S, Alavinejad P, Kamau E, Verma AM, Mendelsohn RB, Khor C, Moss A, Liao DWC, Huang CS, Tsai FC. Impact of COVID-19 on endoscopy training: perspectives from a global survey of program directors and endoscopy trainers. *Clin Endosc*. 2021;54(5):678–87.
 46. Koo CS, Siah KTH, Koh CJ. Endoscopy training in COVID-19: challenges and hope for a better age. *J Gastroenterol Hepatol*. 2021;36(10):2715–9.
 47. Barberio B, Massimi D, Dipace A, Zingone F, Farinati F, Savarino EV. Medical and

- gastroenterological education during the COVID-19 outbreak. *Nat Rev Gastroenterol Hepatol.* 2020;17(8):447–9.
48. Siddiqui UD, Aslanian HR. The new virtual reality: advanced endoscopy education in the COVID-19 era. *Dig Dis Sci.* 2020;65(7):1888–91.
 49. Koo CS, Siah KTH, Low HC, Koh CJ. A low-cost endoscopy trainer for novice endoscopy training in COVID-19. *Endoscopy.* 2020;52(12):E463–4.
 50. Keswani RN, Sethi A, Repici A, Messmann H, Chiu PW. How to maximize trainee education during the coronavirus disease-2019 pandemic: perspectives from around the world. *Gastroenterology.* 2020;159(1):26–9.
 51. Bilal M, Aby ES, Mahmood S, Chiang AL, Breu AC, Charabaty A. Standardized reporting of gastroenterology-related social media scholarship for career advancement. *Nat Rev Gastroenterol Hepatol.* 2021;18(8):519–20.
 52. FitzPatrick M, Clough J, Harvey P, Ratcliffe E. How can gastroenterology training thrive in a post-COVID world? *Frontline Gastroenterol.* 2021;12(4):338–41.
 53. Paleti S, Sobani ZA, McCarty TR, Gutta A, Gremida A, Shah R, Nutalapati V, Bazerbachi F, Jesudoss R, Amin S, Okwara K, Kathi PR, Ahmed A, Gessel L, Hung K, Masoud A, Yu J, Mony S, Akshintala V, Jamil L, Nasereddin T, Kochhar G, Vyas N, Saligram S, Garg R, Sandhu D, Benrajab K, Konjeti R, Agnihotri A, Trivedi H, Grunwald M, Mayer I, Mohanty A, Rustagi T. Impact of COVID-19 on gastroenterology fellowship training: a multicenter analysis of endoscopy volumes. *Endosc Int Open.* 2021;9(10):E1572–8.
 54. Ong J, Ong AML, Ong S, Xin X, Lee YY, Pausawasdi N, De Lusong MA, Makmun D, Chong VH, Ho SH, Lim W, Ong D, Dan YY, Khor C. The impact of the coronavirus disease 2019 pandemic on gastroenterologists in Southeast Asia: a mixed-methods study. *J Gastroenterol Hepatol* [Internet]. 2021 Jun 22; Available from: <https://doi.org/10.1111/jgh.15594>.
 55. Lau LHS, Wong SH, Yip TCF, Wong GLH, Wong VWS, Sung JJY. Collateral effect of coronavirus disease 2019 pandemic on hospitalizations and clinical outcomes in gastrointestinal and liver diseases: a territory-wide observational study in Hong Kong. *Gastroenterology* [Internet]. 2020 Jul 25; Available from: <https://doi.org/10.1053/j.gastro.2020.07.042>.
 56. Lui TKL, Leung K, Guo C-G, Tsui VWM, Wu JT, Leung WK. Impacts of the coronavirus 2019 pandemic on gastrointestinal endoscopy volume and diagnosis of gastric and colorectal cancers: a population-based study. *Gastroenterology.* 2020;159(3):1164–6.e3.
 57. Chen RC, Haynes K, Du S, Barron J, Katz AJ. Association of cancer screening deficit in the United States with the COVID-19 pandemic. *JAMA Oncol.* 2021;7(6):878–84.
- (Cancer screening deficits caused by COVID)
58. Maringe C, Spicer J, Morris M, Purushotham A, Nolte E, Sullivan R, Rachet B, Aggarwal A. The impact of the COVID-19 pandemic on cancer deaths due to delays in diagnosis in England, UK: a national, population-based, modelling study. *Lancet Oncol* [Internet]. 2020 Jul 20; Available from: [https://doi.org/10.1016/S1470-2045\(20\)30388-0](https://doi.org/10.1016/S1470-2045(20)30388-0).
 59. Morris EJA, Goldacre R, Spata E, Mafham M, Finan PJ, Shelton J, Richards M, Spencer K, Emberson J, Hollings S, Curnow P, Gair D, Sebag-Montefiore D, Cunningham C, Rutter MD, Nicholson BD, Rashbass J, Landray M, Collins R, Casadei B, Baigent C. Impact of the COVID-19 pandemic on the detection and management of colorectal cancer in England: a population-based study. *Lancet Gastroenterol Hepatol.* 2021;6(3):199–208.
 60. Basu P, Alhomoud S, Taghavi K, Carvalho AL, Lucas E, Baussano I. Cancer screening in the coronavirus pandemic era: adjusting to a new situation. *JCO Glob Oncol.* 2021;7:416–24.
 61. Liao Z, He S, Sha W, Feng Z, Sun X, Wan J, Pan J, Wang S, Linghu E, Li Z, Capsule Endoscopy Group of the Chinese Society of Digestive Endoscopy. Capsule endoscopy practice during the COVID-19 pandemic: recommendations from the Capsule Endoscopy Group of the Chinese Society of Digestive Endoscopy. *Endosc Int Open.* 2021;9(3):E280–3.
 62. Hennessy B, Vicari J, Bernstein B, Chapman F, Khaykis I, Littenberg G, Robbins D. Guidance for resuming GI endoscopy and practice operations after the COVID-19 pandemic. *Gastrointest Endosc* [Internet]. 2020 May 11; Available from: <https://doi.org/10.1016/j.gie.2020.05.006>.
 63. Kapuria D, Bollipo S, Rabiee A, Ben Yakov G, Kumar G, Siau K, Lee H-W, Congly S, Turnes J, Dhannasekaran R, Lui RN, Global Online Alliance for Liver Studies (GOAL). Roadmap to resuming care for liver diseases after COVID-19. *J Gastroenterol Hepatol* [Internet]. 2020 Jul 13; Available from: <https://doi.org/10.1111/jgh.15178>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.