

## Balkan endemic nephropathy

Ljubica DJUKANOVIĆ<sup>1</sup> and Zoran RADOVANOVIĆ<sup>2</sup>

<sup>1</sup>Clinical Centre of Serbia, Beograd, Serbia

<sup>2</sup>Kuwait University, Kuwait

|   |     |
|---|-----|
| Introduction  | 588 |
| Epidemiological features  | 588 |
| Distribution and frequency  | 588 |
| Demographic data  | 589 |
| Chronological characteristics   | 589 |
| Other epidemiological characteristics   | 590 |
| Overview of the descriptive epidemiological research                                    | 590 |
| Etiology  | 590 |
| Genetic factors   | 590 |
| Biological agents and their products  | 590 |
| Agents from the inanimate environment   | 591 |
| Overview of the etiological research  | 592 |
| Pathomorphological changes  | 592 |
| Macroscopic features  | 592 |
| Morphological studies of renal changes in post mortem material                          | 593 |
| Optic microscopic, immunofluorescent and electron microscopic studies of renal biopsies | 593 |
| Overview of morphological studies   | 593 |
| Clinical features, diagnostics and treatment  | 594 |
| Clinical picture and course   | 594 |
| Laboratory findings   | 594 |
| Imaging methods   | 596 |
| Diagnosis   | 596 |
| Prevention and treatment  | 597 |
| Overview of clinical and laboratory studies   | 597 |
| References  | 598 |

## Introduction

Balkan (or endemic) nephropathy is a chronic tubulo-interstitial disease of unknown, presumably exotoxic etiology. It has been shown to exist only in some parts of the southeastern Europe.

There were about 30 scientific meetings on Balkan nephropathy. Over 500 papers on the disease were published by the year 1970 [1]. By late 1980's, this number rose to 3000 [2]. Sociopolitical turmoil, including wars, and economical hardship prevented any serious research on the problem during the 1990's. In spite of numerous proceedings and a large number of publications on the subject, many features of Balkan nephropathy, its etiology and natural history in particular, remained nearly as mysterious as when described in the mid-fifties.

Meetings organized by international organizations [3-7] had a key role in informing the international scientific community on the disease. An updated source of information is a bilingual (in English and Serbian) monograph published in 2000 [8].

## Epidemiological features

### Distribution and frequency

Though exclusive geographical restriction of the agent(s) of Balkan nephropathy is not very likely, the disease has been diagnosed only among people living (or those who used to live) in more or less well defined areas of the Balkans. Along with Bulgaria and Romania, three republics of the former Yugoslavia have been affected: Bosnia, Croatia and Serbia (Figure 1).

As recently summarized [9], the affected territory has a shape of a rhomboid. Its longer diameter spreads over 500 km (from the Vratza municipality in Bulgaria to villages west of Slavonski Brod in Croatia), while its transversal diameter has about 300 km (from endemic foci in eastern Romania to Vitina municipality in Kosovo). The disease affects individuals who live (or used to live) in rural environment. There are some spared households even in the most affected areas, leading to frequently cited remarks on mosaic distribution of the disease.




Figure 1. Medical geography of Balkan nephropathy.

Topography of the terrain differs between endemic regions. All 14 endemic villages in Croatia are located in a single lowland municipality, at an altitude of about 100 m, while Bosnian foci are found up to 130 m. About 90% of all endemic settlements in Serbia are also situated at a low altitude, below 200 m [10], either in large plains, river valleys or, much more seldom, in hilly regions. There have been no studies of medical geography of Balkan nephropathy in Romania for the last 30-35 years and endemic localities are yet to be determined [11], but the original findings pointed to hilly areas, with endemic villages laying at the bottom of valleys eroded by flooding, at an altitude of 200-300 m. The endemic regions in Bulgaria are described as mountainous or semi-mountainous, without any relationship between endemicity and altitude. Hydrogeological features [12] and lack of floods differ (at least some) Bulgarian foci from other typical endemic regions.

Controversial data on the frequency of Balkan nephropathy are mainly the result of methodological shortcomings [9]. A main obstacle is the operational definition of the disease, leading to huge differences in estimated prevalence rates. The highest reported average annual incidence rate is 16.6 per 1000 in Cakonica, Bulgaria. The average cause-specific mortality rate over 15 years in one of the most affected Serbian foci was 3.3 per 1000 [13].

### Demographic data

During initial Balkan nephropathy research, patients were frequently in their thirties [14], and it was widely accepted that azotemia usually affects the age group 30-50 [3]. Later an apparent shift towards the older ages occurred, with most identified patients being above the age of 60 [9]. The diagnosis of clinical forms before the age of 20 was rare and never independently confirmed. Despite occasional statements on laboratory and bioptic abnormalities in the first decade of life among clinically healthy children from endemic areas, no follow up study ever confirmed that these children developed subsequent kidney disorder.

Both genders are similarly affected, especially considering mortality. As explained in details elsewhere [9], higher prevalence rates among women reported by some authors appears to be a consequence of unreliable diagnostic criteria.

A vast majority of experts believe that a link exist

between agricultural activity and exposure to the agent(s) of Balkan nephropathy. There is also near consensus on the absence of ethnic and/or religious differences as a risk factor for developing the disease. The most convincing data come from Croatia, where the large group of Ukrainians who settled a century ago had the same odds of being affected as the indigenous population. In addition, the first generation immigrants also develop Balkan nephropathy, usually a couple of decades after moving into an endemic region.

Initial studies of affected households show a low standard of living, poor hygienic level, and insufficient nutrition. However, socio-economic factors, including living conditions and well water quality, did not differ between contiguous affected and non-affected households, or between endemic and neighboring non-endemic villages.

### Chronological characteristics

The initial description of Balkan nephropathy emanated from Bulgaria [15, 16] and Serbia [17-19]. By 1957, the disease was recognized in Bosnia and Croatia, and by 1958 in Romania [20].

On retrospect, it was not a newly emerging condition but rather recognition of an already existing endemic process, a previous epidemic wave having occurred in the early forties. Unfortunately, attempts to trace the disease prior to World War 2 are speculative, due to the absence of reliable data and a high frequency of the competing causes of morbidity, notably tuberculosis and malaria.

Data on the secular trend of the disease are contradictory. The earliest affected Romanian endemic village, Ergevitz, now appears disease-free; however, the village was almost deserted and later repopulated by relatively young people (C. Tatu – personal communication). The duration of exposure is too short to make any judgement. The other extreme is the population of two best-known Serbian foci, Sopic and Petka, where the intensity of endemic process is unchanged (D. Bukvic – personal communication). In spite of such controversies, two facts, common to all endemic areas, are crucial in assessing any future trends. These are, an apparent shift of the age distribution of the incidence towards the older age groups, and a much longer natural history of the condition compared to previous data [9].

### Other epidemiological characteristics

Clustering of cases within a household is one of the most conspicuous features of the disease. It is generally agreed that the disease affects both blood related and non-blood related family members. The "phenomenon of simultaneous deaths" (dying of parents and their children within a short interval) has also been observed.

Between 1/3 and 1/2 of patients with Balkan nephropathy develop urothelial tumors [21]. An exceptionally high frequency of these tumors was also observed in the general population of endemic regions [22]. The attributive risk of developing upper urothelial tumors in inhabitants of endemic foci amounts to several dozen [23] or even to as much as 100-200.

There is no evidence that domestic and/or wild animals in endemic regions develop a similar condition.

### Overview of the descriptive epidemiological research

There is general agreement on the following descriptive-epidemiological characteristics of Balkan nephropathy [9]: The disease is known to exist only in some parts of the southeastern Europe, with Central Serbia as the most affected region. Balkan nephropathy has not spread beyond its already defined foci; the disease is distributed mosaically: non-endemic villages exist in the most affected regions, and there are spared families and households in the most affected settlements. There are clustering of cases in families and households. Children and adolescents are spared the clinical disease. Incidence is proportional to age, except for the oldest age groups. There are no clear differences in the cause-specific mortality rates. The excess risk of developing transitional cell urothelial tumors is expressed by two- or even three-digit numbers.

The large majority of researchers support the following statements [9]: autochthonous urban population is spared; rural way of life, i.e., agricultural activity is needed for exposure to the agent(s). Separation from an endemic focus early may prevent the disease, while immigration to an endemic area provides risk of disease development, providing that the exposure was sufficient. Prevalence of the disease has been stable over many years, but now appears to decline in most affected settlements. Incidence rates are shifting towards

the older age groups, and the clinical course is much more protracted, suggesting a less intensive contact with the agent(s) and, consequently, possible future spontaneous disappearance of Balkan nephropathy (providing, of course, that standard of living does not deteriorate too much).

## Etiology

### Genetic factors

The most elaborate and, seemingly consistent, hypotheses regarding etiology come from proponents of heredity as an explanation of the disease occurrence. These authors assume that the risk of developing the disease is restricted only to specific, ethnically distinct, population groups, irrespectively of their place of birth and residence history. Wider acceptance of these hypotheses has been hampered by the different perception of descriptive epidemiology of Balkan nephropathy by a majority of researchers on the topic.

Currently one research group argues in favor of a crucial role of genetic factors. They identified a specific chromosome marker (3q25) in Balkan nephropathy patients [24-26]. Another research team saw some aberrations of the X chromosome, but they resembled changes occurring after exposure to ochratoxin A [27].

Some authors described major anomalies of urinary organs in a high percentage of otherwise healthy children from affected households. "Foetal" and abnormal glomeruli were also observed among these children. Unfortunately no explanation exists for how severe anomalies, notably renal agenesis or double ureters, might disappear in adult life.

Genetic epidemiological approach suggest two possibilities, either polygenic type of inheritance with an insufficient expression of the main gene [28], or monofactorial model with a crucial role of a single gene of incomplete penetrance [29]. In both cases, a contributing environmental factor is needed.

There is no evidence supporting an immunological mechanism in Balkan nephropathy.

### Biological agents and their products

Unspecified viral particles [30], an unidentified cytopathogenic agent, serially propagated slow viruses [31], and an unknown virus associated with foci of

natural infection [32] have been mentioned in the context of Balkan nephropathy etiology. Several specific viruses, notably West Nile [33], coronavirus [34], and papova virus [35], were also suggested as a causative agent. A common feature of all these hypotheses was the unimpressive supporting evidence and ignorance of basic epidemiological features of the disease, in particular its absence of spreading [36].

Bacteria received particular attention in initial stages of the Balkan nephropathy research but, for the last 20 years, their possible etiological importance has been unanimously ruled out [2]. Protozoa have never attracted any attention.

Toxic fungal products are the principle and prime potential culprits. Most efforts have concentrated on ochratoxin A, a mycotoxin responsible for porcine (swine) nephropathy [37]. The substance is found in endemic foci but is also present in neighboring non-endemic areas, and the differences are not statistically significant [38, 39]. Still, the consistent isolating of ochratoxin A in greater frequency and higher concentrations from food and sera samples obtained from endemic, compared to control villages, provides strong support for this hypothesis.

Association of ochratoxin A with chronic interstitial nephropathy in Tunisia [40] and its relation to renal tumors [41] provides additional support for the idea of the etiological role of this mycotoxin. Other fungal toxins such as zerealenone, citrinin [42] and aflatoxin were also isolated in endemic foci. Experimental models suggested that a combination of mycotoxins, rather than a single one, might be involved in the etiology of Balkan nephropathy [43].

Aristolochic acid and its salts, originated from a weed, *Aristolochia clematitis*, have toxic and carcinogenic effect to the kidneys and urothelium [44], respectively. Ivic [45] postulated that this plant may be a cause of Balkan nephropathy, but failed to provide convincing evidence from field surveys. Considerable evidence that *A. clematitis* plays a central role in the etiology of Chinese herb nephropathy [46-48], a condition similar to Balkan nephropathy, requires a second look at this previously abandoned hypothesis.

No local practice, in terms of the use of teas or folk medicine, could be implicated. No one has ever studied flora of the local wells.

### Agents from the inanimate environment

Chronologically, lead poisoning was first offered as an explanation for the occurrence of Balkan nephropathy [17-19]. The idea on lead-contaminated flour led to abandonment of water mills in a part of Central Serbia. This energetic public health action had no impact on the disease frequency.

Effects of non-occupational exposure to cadmium [49], itai-itai disease in particular [50, 51], were occasionally compared with kidney damage seen in Balkan nephropathy patients. In spite of some resembling features, the idea of a common etiology between cadmium nephropathy (including itai-itai disease) and Balkan nephropathy was refuted [50, 52].

Many other metals, including radioactive ones such as uranium [53], were also suggested as possible causative agents of the disease. Results were both non-convincing and non-reproducible.

Inability to identify a single toxic effect of any metal or metalloid as a cause of Balkan nephropathy led researchers to two alternative approaches. First, deficiency, rather than abundance of such a chemical element was proposed [54], with selenium as the most likely candidate [55]. Second, attention was paid to a combined adverse effect of several elements. Synergism of uranium and some other elements, none of which exceeding maximal allowed levels, was proposed [56]. It was also noted that criteria used in occupational medicine (exposure only during working hours) have been applied to an ecological problem (constant exposure) and that concentrations of lead or cadmium within formally acceptable levels, combined with other factors, such as selenium deficiency, might lead to the disease [56]. All these suggestions remained speculative.

As for non-metals, there were attempts to relate Balkan nephropathy to silicon [57-60]. However, when affected and non-affected households were compared, there was even an inverse relationship between the silica content and endemicity. On one occasion, small differences in silica content happened to reach the level of statistical significance but the association was explained as a result of confounding variables [61].

A common hydrogeological characteristic of endemic foci [12] and an inverse relationship to altitude of wells and disease frequency in a longitudinal (cohort) study [61], point to potable water as a vehicle of

the agent(s). However, none of the previously mentioned or several dozen other non-organic substances were associated with the disease [62].

Organics in water have been investigated and provided some interesting data [63]. Except for nitrites [64], chemically unstable substances have not been studied. Wells associated with the disease are situated on alkali soil [65], but the finding was restricted to a single endemic area and has not been reproduced.

Based on chronological data, it is clear that no pesticides, fertilizers or chemicals introduced during the last few decades can be blamed for the occurrence of Balkan nephropathy. Except for exposure to agricultural activities, no occupation, habit (e.g., smoking, alcohol consumption), or hobby (e.g., hunting, fishing) has been shown to precede the disease onset.

### Overview of the etiological research

Genetic factors may play a role, given the same exposure, in developing Balkan nephropathy, upper urothelial tumors, both diseases, or none of them [66]. However, epidemiological data indicate that one or more external, environmental factors are crucial for the occurrence of both Balkan nephropathy and excessive frequency of these tumors in endemic areas.

Among biological agents and their products, probable candidates for etiological agents are mycotoxins and possibly viruses. Toxic role of a plant, notably *Aristolochia clematidis*, is less likely, but cannot be excluded.

As for inanimate environment, there is no chemical element that could be singled out but two facts appear likely. First, insufficient rather than abundant concentrations should be looked at, and second, a combination of factors rather than a single one should be searched for. Organics in water and/or soil are an interesting track to be followed. The same applies to chemically unstable substances.

### Pathomorphological changes


Balkan nephropathy is non-destructive and non-inflammatory tubulointerstitial renal disease [67]. The changes are non-specific and in the chronic, sclerotic phase they are similar to changes observed in other chronic interstitial diseases such as analgesic nephropathy [68], vascular nephrosclerosis [67] cyclosporine-induced nephropathy [69], radiation nephritis [70, 71]

and aging [70], intoxication with silicate, cadmium, lead, uranium [72], *Aristolochia clematidis* [45], mycotoxin ochratoxin A [73], and Chinese herbs [46, 48].

### Macroscopic features

Before introduction of hemodialysis in the treatment of chronic renal patients, the kidneys of patients who died of Balkan nephropathy used to be the smallest seen at post mortem examinations, weighing 14.8-80 g each (Figure 2A), the differences between the left and right kidneys being small (5-20 g) [72, 74-76]. Surface of the kidneys is smooth, occasionally wavy, but never granulated or roughly nodular. The section shows markedly narrowed cortex, pyramid and Bertin's columns are fairly well preserved, and corticomedullary border is well differentiated. Papillary necrosis of the pyramids has not been found.

Small, papillary, usually multiple tumors of the renal pelvis and ureters are also one of the characteristic findings (Figure 2B). In post-mortem studies tumors were reported in 8-50% of cases [72, 77].



**Figure 2.** A. Macroscopic appearance of the right kidney weighing 35 g in a case of BN, surface is smooth, occasionally mildly wrinkled (autopsy, a man aged 48). B. Multifocal papillary tumor of the right ureter (autopsy).

### Morphological studies of renal changes in post mortem material

Diffuse fibrosis of cortical interstitium and tubular atrophy may be observed in the absence of significant cellular interstitial infiltration. In contrast to the cortex, Bertin's columns are less markedly affected. Even in severe tubulointerstitial cortical changes, glomeruli are well preserved, partially collapsed, and subsequently subjected to focal or generalized sclerosis causing collapse. Glomeruli in Bertin's column are occasionally compensatorily enlarged. Pyramids are preserved or less markedly affected [72, 74, 77]

Blood vessels, arcuate or interlobular arteries, and arterioles are affected in the form of intimal sclerosis and thickening of lamina elastica interna. In addition, the blood vessels are compressed and torsioned [77].

In cases of tumors of the renal pelvis and ureters, morphological signs of pyelonephritis are recognized [72].

### Optic microscopic, immunofluorescent and electron microscopic studies of renal biopsies

In oligosymptomatic clinical cases, rare disseminated foci of interstitial fibrosis and tubular atrophy with preserved glomeruli are seen. These changes have no special predilection of distribution and are not inflammatory. They tend to be triangular, with the base oriented toward the renal surface [67, 77]. In cases with initial renal failure, the fields of acellular interstitial fibrosis are larger and diffuse.

The striking atrophic process observed in Balkan nephropathy suggests that apoptosis may play a role in this disease. In this context it is of interest that Savin et al. observed an increased apoptosis to proliferation ratio at the level of the tubuli [78].

The glomeruli are usually affected by generalized [80%] or segmental sclerosis [10%] and only in 8% is hyalinosis recorded. Double contour glomerular basement membrane occurs in 22% of the cases. In 2.7-6% of cases fetal-like glomeruli can be seen in the kidneys, while glomerular hypercellularity was recorded in 4% [70, 77, 79].

The most interesting changes are present in pre- and postglomerular blood vessels. In about 50% of cases PAS positive proteins are deposited in vasa afferens walls in a focal segmental or circumferential manner

in the form of droplets, bands or granules [70, 77, 79]. No tissue reaction is seen around these arterioles.

Interlobular capillaries are filled with thick, proteinaceous substances that are also deposited below the capillary endothelium and may even be found free in the interstitium. These changes are described as capillary sclerosis [68]. Although renal vascular changes in Balkan nephropathy have been pointed out as very important, they are not specific and can be encountered in other renal diseases. Ferluga et al. [70] and Sindjic [77] point out to similarities in early changes encountered in Balkan nephropathy with cyclosporine-induced nephrotoxic changes.

Immunofluorescence revealed irregular and scarce deposits of C3, fibrin and IgM, and occasionally IgA, C1q and C4, mainly on the vascular walls, Bowman's capsule and some sclerotic glomeruli [67, 80, 81].

Electron microscopic findings are either normal or correspond to degenerative and sclerotic changes. While some authors describe virus-like particles [34, 82, 83], others point out that such particles were not found [70, 71].

Despite these findings, some authors described Balkan nephropathy as a form of glomerulonephritis [79, 84]. However, the lack of reliable evidence supporting glomerulonephritis has led to it being discarded [71, 75] and abandoned even by its advocates [85].

Optic microscopic, immunofluorescent and electron microscopic studies of renal biopsies in children aged 5-15 from affected families in endemic regions failed to detect any Balkan nephropathy related changes [77].

### Overview of morphological studies

It is generally agreed that the morphologic changes of Balkan nephropathy are not specific and correspond to non-destructive, non-inflammatory kidney disease accompanied by marked changes on the blood vessels in both early and late stages of the disease, interstitial, multifocal fibrous expansion and severe tubular atrophy mainly in the upper cortex [67, 70, 71, 77, 79].

Changes on kidney arterioles have been described suggesting that the changes in early stage of the disease may be responsible for the development of multifocal, ischemic, vascular nephrosclerosis encountered in chronic stages of the disease [67, 70]. On the other hand, close similarity of Balkan nephropathy with an-

algesic and cyclosporine-induced nephropathy has been recognized [69, 70, 77]. All this leads to a suggestion that Balkan nephropathy develops following a model of toxic nephropathy, targeting primarily the vascular endothelium where the tubular epithelium is affected either directly or indirectly due to accompanying ischemia.

## Clinical features, diagnostics and treatment

### Clinical picture and course

Balkan nephropathy is a chronic tubulointerstitial disease with occult, insidious onset, usually progressing slowly without apparent signs and symptoms. After a long asymptomatic period, the disease is manifested as chronic renal failure. Less commonly blunt lumbar pain or renal colic may develop or, occasionally, dysuric symptoms induced by urinary tract infection. If hematuria exists, urothelial tumor should be suspected. In an advanced case polyuria and nocturia are present due to impaired concentrating ability of the kidneys. The disease is tolerated well and the patients preserve their working ability until advanced stages of renal failure [18, 74, 86, 87].

Objective examination reveals a characteristic skin tan of Balkan nephropathy patients: a pale yellow with copperish glow on the cheeks has been recognized since the augural reports on the disease [18, 86]. Besides, xanthochromia of the palms and soles is also frequently observed.

Patients with Balkan nephropathy do not suffer from edema, and their blood pressure is usually normal [18, 86-88]. In the advanced phase of the disease physical examination detects signs of dysfunction of other organs or systems as a consequence of chronic renal failure [19].

As Balkan nephropathy is characterized by a slow asymptomatic course, most authors identify two main stages of the disease: the first, asymptomatic (latent, subclinical) and second, manifest (symptomatic). The latter is usually subdivided into the stage without renal failure (early, compensated Balkan nephropathy, with no azotemia) and chronic renal failure (decompensated Balkan nephropathy, uremia) [19, 86, 87].

An important feature of Balkan nephropathy is its association with a high incidence of tumors of the re-

nal pelvis and ureters [22, 89]. Controversially, urinary bladder tumors are not more frequent in the regions of Balkan nephropathy compared to non-endemic ones [90]. Upper urothelial tumors of patients originating from the region with Balkan nephropathy differ from the same tumors identified in patients from other regions because of their similar incidence in both sexes, bilateral occurrence, slower evolution and more common association with chronic renal failure [90].

### Laboratory findings


Appearance and urine color are unchanged in most patients with Balkan nephropathy. Urine sediment is usually scarce; when microhematuria or leukocyturia are encountered they are usually associated with the occurrence of tumors or urinary tract infection [86, 87].

Bacteriological studies usually reveal sterile urine, but in 8.3-31.8% significant bacteriuria was confirmed and considered due to superimposed urinary tract infection [86, 87].


Proteinuria is a common finding in patients with Balkan nephropathy [18, 86]. It is usually intermittent, less than 1 g per day but becomes permanent in advanced renal failure [91]. Although proteinuria is one of the criteria for diagnosis of Balkan nephropathy, it has been reported in healthy members of endemic families [29, 91, 92]. Tubular proteinuria is most common and increased excretion of low-molecular weight proteins such as beta2-microglobulins, lysozyme, ribonuclease, light chains of immunoglobulin, retinol-binding protein has been reported [93-98]. Beside tubular proteinuria, smaller numbers of patients manifest mixed proteinuria, while in patients with renal failure, glomerular proteinuria may be encountered [96, 98].

Anemia has been noted in patients with Balkan nephropathy in early studies [18] and described as normocytic and normochromic or mildly hypochromic [86]. It has been suggested that anemia occurs earlier in the course of the disease progression than is the case in other renal diseases and that it precedes azotemia [86, 99], however, recent studies have failed to substantiate this claim [91, 100]. Also, there is no evidence that the anemia in Balkan nephropathy differs from anemia accompanying other renal diseases in either features [100] or rate of deterioration in the progression of renal failure [101]. Nevertheless, anemia in Balkan nephropathy patients treated with hemodialysis is







**Figure 3.** Interstitial fibrosis and tubular atrophy; glomerulus with mild mesangial hypercellularity and another with incomplete hyalinosis. PAS, x120.



**Figure 4.** Interlobar artery showing intimal fibrosis. PAS, x240.



**Figure 5.** Reduced number of tubules; fibrotic interstitium; few infiltrating cells. Masson's trichrome, x120.



**Figure 6.** Extensive interstitial scarring associated with severe arterio-arteriosclerosis. PAS, x120.

more severe than in patients with other renal diseases [101].

The leukocyte count in the peripheral blood of patients with Balkan nephropathy is normal and without pathological changes in the differential count and bone marrow [19, 86].

Investigation of renal function in patients with endemic nephropathy has revealed tubular dysfunction in the earliest stage of the disease: renal glycosuria, increased uric acid and amino acid excretion [94], as well as increased excretion of low molecular weight proteins [97]. Significantly higher activity of cellular enzymes in the urine and increased urinary excretion of Tamm-Horsfall protein have been described in patients with Balkan nephropathy, as well as in healthy members of endemic families [102]. Findings of a distal tubular disorder (impaired urinary acidification, impaired urine concentrating ability) were described in earlier studies [86, 87] but could not be confirmed in studies conducted in larger groups of patients with normal or mildly impaired glomerular filtration rate [91, 103]. The occurrence of certain disorders of the tubular function recorded in the course of chronic renal failure (increased natriuria, phosphaturia) can be considered as the result of kidney adaptation to the lost nephron mass, instead of Balkan nephropathy properties [103].

The immunological studies have failed to indicate that immune disorders participate in the pathogenesis of Balkan nephropathy, with some of the detected changes having been attributed to advanced renal failure [104].

### Imaging methods

Different methods of kidney imaging have shown that Balkan nephropathy patients with chronic renal failure have symmetrically shrunken kidneys with smooth surface and no calcifications [88]. The time at which the shrinking occurs remains to be determined. While some authors suggest that the size of the kidneys remains normal in patients in the latent phase of the disease and with normal renal function, others report cases of shrunken kidneys in patients in an early phase with normal glomerular filtration rate, and it was even proposed that the disease was characterized with primarily small kidneys [91, 103, 105].

Excretory urography does not reveal changes in the

pyelocaliceal system, except in cases with secondary infection or urothelial tumors.

Radionuclide methods have shown that renal plasma flow impairment is the first sign of the early phase of renal failure. Glomerular and tubular functions correspond to the severity of the disease.

### Diagnosis

The most commonly used criteria for the diagnosis of Balkan nephropathy are those proposed by Danilovic et al [99]. They include: (1) farmers in the endemic villages, (2) familial history positive for endemic nephropathy, (3) mild proteinuria, (3) low specific gravity of the urine, (4) anemia, early occurrence, normochromic or hypochromic, (5) retention of nitrous compounds in the blood (urea > 50 mg/dl, creatinine > 1.5 mg/dl), (6) symmetrically shrunken kidneys. Using these criteria, Danilovic classified patients into the following groups:

1. *potential*, a group with intermittent proteinuria and positive familial history, including those that fulfill at least the first three criteria;
2. *suspected*, a group that in addition to the first three fulfill at least one of the remaining three criteria;
3. *affected*, a group that fulfill at least 5 out of 6 criteria;
4. *decompensated* patients that fulfill at least 5 out of 6 criteria and have urea values > 150 mg% and manifested signs of uremia.

Analysis of these criteria leads to the conclusion that they enable detection only of patients with overt disease, and the criteria are not sufficiently specific to enable a reliable diagnosis. Therefore, numerous studies have focused on developing sufficiently sensitive and specific criteria to allow diagnosis in the early phase. Although indications of tubular disorders, particularly tubular proteinuria, may be used as sufficiently specific diagnostic criteria, so far no single clinical or laboratory finding is considered pathognomonic for Balkan nephropathy when differentiate it from other tubulointerstitial diseases.

The diagnosis of Balkan nephropathy is now established according to epidemiological criteria suggested by Danilovic (criteria 1-3) [99], presence of tubular proteinuria and ruling out other renal diseases.

Histopathological analysis makes the diagnosis of



Balkan nephropathy much easier [70, 77], and it is considered indispensable in classifying the following groups of patients with urinary abnormalities suggestive of endemic nephropathy:

1. Patients from families that were not previously been affected with endemic nephropathy, but live in an endemic village;
2. In cases of nephropathy of unknown etiology in villages close to endemic foci;
3. In immigrants to endemic regions and in emigrants from these regions [104].

Differential diagnosis of Balkan nephropathy should include all chronic, slowly progressive renal diseases, primarily chronic tubulointerstitial diseases. Although no specific indicators of Balkan nephropathy have been recognized, epidemiological data, familial history as well as clinical characteristics of the disease enable differential diagnosis. Thus, shrunken kidneys with smooth surface are characteristic of Balkan nephropathy and they differentiate it from analgesic nephropathy, pyelonephritis or reflux nephropathy that are characterized by shrunken kidneys with uneven surface. Pyelocaliceal system of the kidneys remains unaffected in patients with Balkan nephropathy, unlike the characteristic changes observed in pyelonephritis or obstructive nephropathy. Absence of papillary calcifications also enables differentiation of Balkan nephropathy from analgesic, obstructive, reflux nephropathy [103, 106].

Recently similarity of Balkan nephropathy and nephropathy induced by Chinese herbs used in slimming diets have been suggested [47]. Nevertheless, Chinese herb nephropathy is a rapid progressive tubulointerstitial disease with pronounced fibrosis and progression towards end-stage renal disease within few years, clearly different from the protracted clinical course of Balkan nephropathy.

### Prevention and treatment

Balkan nephropathy is a disease of unknown etiology, so that recommendations regarding effective prevention are not possible. Efforts have been made to improve the living conditions, bring high quality drinking water to endemic villages and undertake other hygienic measures. Treatment is planned according to the stage of renal impairment. In principle, the treatment involves treating any reversible factors,

which deteriorate the renal function and measures applied in chronic renal failure [107].

End-stage renal disease is treated with dialysis and kidney transplantation. Hypertension and cardiovascular diseases affect the Balkan nephropathy patients less frequently, so they tolerate hemodialysis rather well compared to patients with other renal diseases. The Balkan nephropathy patients on long-term hemodialysis frequently develop upper urothelial or urinary bladder carcinoma.

Although the number of reported cases with kidney transplant is small, neither specific post-transplantation problems nor disease recurrence on the transplanted kidney have been described.

### Overview of clinical and laboratory studies

Balkan nephropathy is a chronic tubulointerstitial disease with insidious occult onset which progresses without symptoms. Agreement as to how to define the early asymptomatic phase of the disease is lacking, since no specific indicators for the diagnosis have been recognized. The diagnosis is established according to epidemiological criteria (farmers in endemic villages, familial history positive for endemic nephropathy), presence of tubular proteinuria, findings of symmetrically shrunken kidneys with smooth surface, without calcifications and ruling out of other renal disease. Renal biopsy may make the diagnosis easier, although the changes are non-specific. One important feature of Balkan nephropathy is the associated high incidence of tumors of the renal pelvic and ureters.

So far, laboratory studies have failed to detect any disorder as a specific marker for early detection of the disease or a reliable indicator for differential diagnosis. Laboratory studies have confirmed that Balkan nephropathy is a tubulointerstitial disease so that tubular disorders precede impairment of glomerular filtration. Although anemia is one of the criteria for the diagnosis of the disease, it has not been evidenced that pathogenesis and features of this anemia differ from that observed in other chronic renal diseases. It is only more severe in end-stage Balkan nephropathy patients than in patients with other kidney diseases.

### Acknowledgement

Photos of all tissue specimens were provided by Dr. Jovan Dimitrijevic.

## References

1. Astrug A. (editor). Endemic Nephropathy – Bibliography 1956-1970, Sofia: Ministry of Public Health; 1971.
2. Radovanovic Z. Aetiology of Balkan nephropathy: a reappraisal after 30 years. *Eur J Epidemiol* 1989; 5: 372-377.
3. World Health Organization. The "Endemic Nephropathy" of South-Eastern Europe. *Bull Wld Hlth Org* 1965; 32(3): 431-448.
4. Wolstenholme GEW, Knight J (editors). The Balkan Nephropathy. Ciba Foundation Study Group No. 30. London: Churchill; 1967.
5. World Health Organization. WHO Meeting of Investigators on Endemic Nephropathy, Belgrade and Lazarevac, November 1974, NCD/WP/74.
6. Castegnaro M, Plestina R, Dirheimer G, Chernozemsky IN, Bartsch H (editors). Mycotoxins, Endemic Nephropathy and Urinary Tract Tumours. IARC Scientific Publications No. 115. Lyon: International Agency for Research on Cancer; 1991.
7. Hall P, Batuman V (guest editors). *Kidney Int* 1991; 40: Suppl 34 (Proc IUPAB meetings, Belgrade, 1989 and 1990).
8. Radovanovic Z, Sindjic M, Polenakovic M, Djukanovic Lj, Petronic V. Endemic Nephropathy. Belgrade: Office for Textbooks and Teaching Aids, 2000.
9. Radovanovic Z. Epidemiology and aetiology of endemic nephropathy. In: Radovanovic Z, Sindjic M, Polenakovic M, Djukanovic Lj, Petronic V. Endemic Nephropathy. Belgrade: Office for Textbooks and Teaching Aids, 2000.
10. Radovanovic Z. Topographical distribution of the Balkan endemic nephropathy in Serbia (Yugoslavia). *Trop geogr Med* 1979; 31: 185-189.
11. Gluhovschi G, Stefanovic V, Dimitrov Tz, et al. Endemic Balkan Nephropathy. Helicon, Timisoara, 1994.
12. Radovanovic Z, Peric J. Hydrogeological characteristics of endemic nephropathy foci. *Public Health* 1979; 93: 76-81.
13. Radovanovic Z, Danilovic V, Velimirovic D et al. Beta<sub>2</sub>-microglobulinuria as a predictor of death in a population exposed to Balkan endemic nephropathy. *Kidney Int* 1991; 40(S34): 32-34.
14. Danilovic V. Effet des recherches des néphrites endémiques en Yougoslavie. *Arch. Union Méd Balkan* 1964; 2(1): 31-40.
15. Tancev I, Evstatijev P, Dorosiev D, Penceva Z, Cvetkov G. Proucvanija na nefritite v'v Vracanska okoliija. *S'vr Med* 1956; 7(9): 14-29.
16. Ivanov L. Proucvanija v'rhu etiologijata na b'brecnite zaboljavanija v'v Vracanska okoliija. *S'vr Med* 1956; 7(9): 30-34.
17. Danilovic V, Djuriscic M, Mokranjac M, Stojimirovic B, Zivojinovic J, Stojakovic P. Porodicna oboljenja bubrega u selu Sopic izazvana hronicnom intoksikacijom olovom. *Srp arh celok lek* 1957; 85(10): 1115-1125.
18. Danilovic V, Djuriscic M, Mokranjac M, Stojimirovic B, Zivojinovic J, Stojakovic P. Néphrites chroniques provoquées par l'intoxication au plomb par voie digestive (farine). *Presse Méd* 1957; 65(90): 2039-2040.
19. Danilovic V. Chronic nephritis due to ingestion of lead-contaminated flour. *Brit Med J* 1958; 1: 27-28.
20. Milojevic B. Epidemic of chronic nephritis of unknown aetiology. *Brit Med J* 1960; i: 244-245.
21. Radovanovic Z, Jankovic S, Jevremovic I. Incidence of tumors of urinary organs in a focus of Balkan endemic nephropathy. *Kidney Int* 1991; 40(S34): 75-76.
22. Petkovic S, Mutavdzic M, Petronic V, Markovic V. Les tumeurs du bassin et de l'uretère. *Recherches cliniques et étiologique. J Urol Néphrol* 1971; 77(6): 429-439.
23. Radovanovic Z, Krajcinovic S, Petkovic S, Hall PW. Papillary transitional cell tumours, Balkan nephropathy, and  $\beta_2$ -microglobulin. *Lancet* 1981; 2: 689.
24. Toncheva D, Dimitrov Ts, Tzoneva M. Cytogenetic studies in Balkan endemic nephropathy. *Nephron* 1988; 48: 18-21.
25. Toncheva D, Gergov T, Tzoneva M, Bouchakliev Z. Spontaneous and induced chromosome aberrations in Balkan endemic nephropathy. *Kidney Int* 1991; 40 (Suppl 34): 97-101.
26. Toncheva D, Dimitrov T. Genetic predisposition to Balkan endemic nephropathy. *Nephron* 1996; 72(4): 564-569.
27. Manolova Y, Manolov G, Parvanova L, Petkova-Bocharova T, Castegnaro M, Chernozemsky IN. Induction of characteristic chromosomal aberrations, particularly X-trisomy, in cultured human lymphocytes treated by ochratoxin A, a mycotoxin implicated in Balkan endemic nephropathy. *Mutat Res* 1990; 231(2): 143-149.
28. Tucic N, Marinkovic D. The inheritance of liability of Balkan nephropathy. *Genetika* 1979; 11(3): 213-220.
29. Banjevic D, Tucic N. The monofactorial model for inheritance of liability to disease and its implications for relatives at risk. *Genetika* 1979; 11(3): 221-230.
30. Georgescu L, Litvac B, Manescu N, Petrovici A, Schwartzkopf A, Zosin C. Particules virales dans le rein de la néphropathie endémique Balcanique. *Sem Hôp Paris* 1970; 46: 3526.
31. Draganescu N, Antipa C, Girjabu E, Melencu M. Endemic Balkan nephropathy, a slow virus disease? *Rev Roum Med Virol.* 1983; 34(1): 3-6.
32. Birtasevic B, Vukovic B, Drndarevic D, Seguljev Z, Obradovic M, Pokorni D, Cobeljic M, Stojic P, Stefanovic S, Stankovic A, et al. Endemska (Balkanska) nefropatija kao prirodno-zariska infekcija? *Vojno-sanitetski pregled* 1983; 40: 319-324.
33. Kraus N. Studien über die endemische Nephrozirrhose der Balkanhalbinsel. *Zeitschrift für ärztliche Fortbildung* 1966; 60: 829-834.

34. Apostolov K, Spasic P, Bojanic N. Evidence of a viral etiology in endemic (Balkan) nephropathy. *Lancet* 1975; 2: 1271-1273.
35. Nastac E, Stoian M, Hozoc M, Iosipenco M, Melencu M. Further data on the prevalence of serum antibodies to papova viruses (BK and SV40) in subjects from the Romanian area with Balkan endemic nephropathy. *Rev Roum Med-Virol* 1984; 35(1): 65-67.
36. Radovanovic Z. Epidemiological evidence on Balkan nephropathy as a viral disease. *Medical Hypotheses* 1987; 22: 171-175.
37. Krogh P. Causal association of mycotoxic nephropathy. *Acta Path Microbiol Scand Sect A (Suppl)* 1978; 269: 1-28.
38. Petkova-Bocharova T, Castegnaro M. Ochratoxin A contamination of cereals in an area of high incidence of Balkan endemic nephropathy in Bulgaria. *Food Addit Contam* 1985; 2: 267-270.
39. Cvetnic Z, Pepeljnjak S. Ochratoxinogenicity of *Aspergillus ochraceus* strains from nephropathic and non-nephropathic areas in Yugoslavia. *Mycopathologia* 1990; 110(2): 93-99.
40. Maaroufi K, Achour A, Betbeder AM, Hammami M, Ellouz F, Creppy EE, Bacha H. Foodstuffs and human blood contamination by the mycotoxin ochratoxin A: correlation with chronic interstitial nephropathy in Tunisia. *Arch Toxicol* 1995; 69: 552-558.
41. Maaroufi K, Pfohl-Leszkowicz A, Achour A, el May M, Grosse Y, Hammami M, Ellouz F, Creppy EE, Bacha H. Ochratoxin A genotoxicity, relation to renal tumors. *Arch. Inst. Pasteur Tunis* 1994; 71(1-2): 21-31.
42. Yeulet SE, Mantle PG, Rudge MS, Greig JB. Nephrotoxicity of *Penicillium aurantiogriseum*, a possible factor in the aetiology of Balkan endemic nephropathy. *Mycopathologia* 1988; 102: 21-30.
43. Bach PH, Gregg NJ, Delacruz L. Relevance of a rat model of papillary necrosis and upper urothelial carcinoma in understanding the role of ochratoxin A in Balkan endemic nephropathy and its associated carcinoma. *Food Chem. Toxicol* 1992; 30(3): 205-211.
44. Mengs U, Lang W, Poch JA. The carcinogenic action of aristolochic acid in rats. *Arch Toxicol* 1982; 51: 107-119.
45. Ivic M. Problem etiologije endemske nefropatije. *Acta Fac Med Naiss* 1970; 1: 29-38.
46. Vanherweghem JL, Depierreux M, Tielemans C, Abramowicz D, Dratwa M, Jadoul M, Richard C, Vandervelde D, Verbeelen D, Vanhaelen-Fastre R, et al. Rapidly progressive interstitial renal fibrosis in young women: association with slimming regimen including Chinese herbs. *Lancet* 1993; 341:387-391.
47. Cosyns JP, Jadoul M, Squifflet JP, De Plaen JF, Ferluga D, van Ypersele de Strihou C. Chinese herbs nephropathy: a clue to Balkan endemic nephropathy? *Kidney Int* 1994; 45(6): 1680-1688.
48. Cosyns JP, Dehoux JP, Guiot Y, Goebbels RM, Robert A, Bernard AM, van Ypersele de Strihou C. Chronic aristolochic acid toxicity in rabbits: a model of Chinese herbs nephropathy? *Kidney Int.* 2001; 59(6): 2164-2173
49. Nogawa K, Kobayashi E, Honda R, Ishizaki A, Kawano S, Matsuda H. Renal dysfunctions of inhabitants in a cadmium-polluted area. *Environ Res* 1980; 23(1): 13-23.
50. Danilovic V (editor). *Bulletin, T. LXIX, Classe des sciences medicales No. 12.* Beograd: Academie Serbe des Sciences et des Arts, 1980.
51. Frieberg L, Elinder C-G, Kjellstrom T, Nordberg GF (editors). *Cadmium and health: a toxicological and epidemiological appraisal.* CRC Press, Boca Raton 1986.
52. Piscator M, Kostial K, Plestina R. Urinary excretion of cadmium and zinc among women with Balkan endemic nephropathy. *Trace Elements Med* 1984; 1: 134-138.
53. Ignjatovic B. Prilog proucavanju etiologije hronicnog nefrita. *Narodno zdravlje* 1962; 18: 329-334.
54. Maksimovic Z, Radovanovic Z. Balkan endemic nephropathy in Yugoslavia and geochemical studies. In: *Trace Substances in Environmental Health (18).* Hemphill DD (editor). Univ Missouri 1984; p 230-236.
55. Maksimovic Z. Selenium deficiency and Balkan endemic nephropathy. *Kidney Int* 1991; 40 (suppl 34): S12-S14.
56. Wedeen RP. Environmental renal disease. Lead, cadmium and Balkan endemic nephropathy. *Kidney Int* 1991; 40 (suppl 34): S4-S8.
57. Duancic V. Essai d'explication de la manifestation de l'affection rénale de masse à Slavonski Kobas. *Biol glasnik* 1960; 3: 43-50.
58. Markovic B, Lebedev S. Role étiopathogénique des silicates érosifs dans la néphrite endémique. *Presse méd.* 1965; 73(8): 401-404.
59. Markovic B, Arambasic M. Experimental chronic interstitial nephritis compared with endemic human nephropathy. *J Pathol* 1971; 103: 35-40.
60. Goldsmith JR, Goldsmith DF. Fiberglass or silica exposure and increased nephritis or ESRD (end-stage renal disease). *Am J Ind Med* 1993; 23: 873-881.
61. Radovanovic Z, Markovic-Denic Lj, Marinkovic J, Jevremovic I, Jankovic S. Well water characteristics and the Balkan nephropathy. *Nephron* 1991; 57: 52-54.
62. Radovanovic Z, Edmunds WM. Groundwater chemistry and the incidence of Balkan endemic nephropathy. *Environ Geochem Health* 1991; 13(2): 43-49.
63. Finkelman RB, Feder GL, Orem WH, Radovanovic Z. Relationship between low-rank coal deposits and Balkan endemic nephropathy. *AGID News* 1991; 65: 23.

64. Radovanovic Z, Stevanovic D. Different forms of mineral nitrogen in drinking water and the Balkan nephropathy. *Arch Environm Contam Toxicol* 1988; 17: 813-815.
65. Filipovic A, Abinun A, Nijemcevic S, Lekovic G, Popovic N. Ispitivanje ovisnosti pojave endemske nefropatije i sastava zemljišta iz kojeg stanovništvo uzima vodu za pice u opštini Modrica. *Medicinski arhiv* 1977; 31(5): 277-284.
66. Radovanovic Z, Velimirovic D, Naumovic T. Upper urothelial tumours and the Balkan nephropathy - Inference from the study of a family pedigree. *Eur J Cancer* 1990; 26: 391-392.
67. Sindjic M. Pathomorphology of the kidneys in endemic nephropathy. *Academie Serbe des Sciences et des Arts Bulletin* 1980; 69: 127-140.
68. Zollinger HU, Mihatsch MJ, Torhost J. Nephropathy due to analgesics. Morphological aspects. XIII Inter. Congr Inter Acad Pathology, Paris, 1980; 17 (Abstract).
69. Mihatsch MJ, Thuel G, Ryffel N. Morphology of cyclosporin nephropathy. *Prog Allergy* 1986; 38: 447-465.
70. Ferluga D, Hvala A, Vizjak A, Trnacevic S, Halilbasic A. Renal function, protein excretion and pathology of Balkan endemic nephropathy. III Light and electron microscopy studies. *Kidney Int* 1991; 40 (Suppl 34): S54-S67.
71. Hall PW III, Dammin GJ. Balkan nephropathy. In: *Renal pathology with clinical and functional correlations*. Fisher G, Brenner BM (editors). JB Lippincott Co, Philadelphia, 1989; p. 913-924.
72. Sindjic M. Resultats des recherches patho-anatomiques, faites jusqu'a present, concernant la nephropathie endemique. In: I Symposium sur la nephropathie endemique. 1973 Beograd. Danilovic V (redacteur). Academie Serbe des Sciences et des Arts, Beograd 1973; 187-202.
73. Krogh P, Hald B, Pleština R, Ceovic S. Balkan (endemic) nephropathy: evidence for support of a causal relationship. In: *Endemic (Balkan) Nephropathy. Proceedings of the 4th Symposium on Endemic (Balkan) Nephropathy*. Strahinjac S, Stefanovic V (editors). University Press, Niš, 1979; 35-43.
74. Radonic M, Radošević Z, •upanic V. Endemic nephropathy in Yugoslavia. In: *The Kidney*. Mostby S, David E (editors). Williams & Wilkins Co, Baltimore 1966; p. 503-522.
75. Zosin L et al. Aspects anatomo-cliniques de la néphropathie endémique balkanique. *Sém Hôp*, 1966; 42(3/1): 194-201.
76. Hall PW et al. Investigation of chronic endemic nephropathy in Yugoslavia. II *Renal Pathology*. *Am J Med* 1965; 39(2): 210-217.
77. Sindjic M. Morphological changes in kidneys affected by endemic nephropathy. In: Radovanovic Z, Sindjic M, Polenakovic M, Djukanovic LJ, Petronic V. *Endemic nephropathy*. Belgrade: Office for Textbooks and Teaching Aids, 2000.
78. Savin M, Bumbaširevic V, Djukanovic LJ, Petronic V. The significance of apoptosis for early diagnosis of Balkan nephropathy. *Nephrol Dial Transplant* 2001; 16 (Suppl 6): 30-32.
79. Doichinov D, Todorov D, Dinev I, Milev B. Biopsy specimen examination during early stage of endemic nephropathy. In: II Symposium sur la nephropathie endemique. 1977 Beograd-Lazarevac. Danilovic V (redacteur). Academie Serbe des Sciences et des Arts, Beograd 1979; p. 189-194
80. Polenakovic M, Hrizoho D. Endemska nefropatija – imunofluorescentna mikroskopska ispitivanja. In: II Simpozijum o endemskoj nefropatiji, Beograd-Lazarevac 1977; Danilovic V (editor) SANU, Beograd 1979; p. 137-150.
81. Vizjak A, Trnacevic S, Ferluga D, Halilbašić A. Renal function, protein excretion and pathology of Balkan endemic nephropathy. IV-Immunohistology. *Kidney Int* 1991; 40(34): 68-74.
82. Georgescu L, Litvac B, Diosi P, Plavosin L, Herzog G. Viruses in endemic (Balkan) nephropathy. *Lancet* 1976; i: 1086.
83. Uzelac-Keserovic B, Vasic D, Ikonomovski J, Bojanic N, Apostolov K. Isolation of a coronavirus from urinary tract tumours of endemic Balkan nephropathy patients. *Nephron* 2000; 86: 93-94
84. Pantic VR, Suša S. Ultrastructure of nephrons in kidney of endemic nephropathy patients. In: *Endemic (Balkan) Nephropathy. Proc 4th Symp*, Nis 1979. Strahinjac S, Stefanovic V (editors). Inst Nephro Haemod, Nis 1981; p. 123-132.
85. Doitchinov D. Immunohistochemical studies in Balkan endemic nephropathy. In: *Endemic (Balkan) Nephropathy. Proc 4th Symp*, Nis 1979. Strahinjac S, Stefanovic V (editors). Inst Nephro Haemod, Nis 1981; p. 265.
86. Puchlev A, Popov N, Astrug A, Dotchev D. Clinical studies on endemic nephropathy in Bulgaria. In: *Int Symp on End Nephro*, Sofia, 1963, Bulg Acad Sc Press, Sofia 1965; p. 17-24.
87. Bruckner I, Zosin C, Lazarescu R, et al. A clinical study of nephropathy of an endemic character in the People's Republic of Rumania. In: *Int Symp on Endemic Nephropathy*, Sofia 1963. Bulg Sofia: Acad Sc Press; 1965; p. 25-35.
88. Djukanovic LJ, Velimirovic D, Sindjic M. Balkan Nephropathy. In: *Clinical nephrotoxins - renal injury from drugs and chemicals*. De Broe ME, Porter GA, Bennett WM, Verpooten GA (editors). Kluwer Academic Publishers, Dordrecht 1988; p. 426-436.
89. Petkovic S. Epidemiology and treatment of renal pelvic and ureteral tumours. *J Urol* 1971; 114: 858.
90. Petronic V. Tumors of the upper urothelium and endemic nephropathy. In: *Endemic nephropathy*. Radovanovic Z, Sindjic M, Polenakovic M, Djukanovic LJ, Petronic V (editors). Office for Textbooks and Teaching Aids, Beograd, 2000.
91. Trnacevic S, Halilbašić A, Ferluga D, Plavljanic D, Vizjak A, Durakovic H, Habul V, Mesic E, Imamovic G, Hranisavljevic J, et al. Renal function, protein excretion and pathology of Balkan endemic nephropathy. I. Renal function. *Kidney Int (Suppl)* 1991; 34: S49-S51.

92. Cvorišec D, Radonic M, Ceovic S, Aleraj B. Characteristics of proteinuria in endemic nephropathy. *J Clin Chem Clin Biochem* 1983; 21: 569-571.
93. Bruckner I, Stoica G, Serban M. Studies on urinary proteins. In: *The Balkan Nephropathy*. Wolstenholme GEW and Knight J (editors). Ciba Foundation Study Group No. 30. Churchill, London 1967; p. 15-20.
94. Hall P W, III, Piscator M, Vasiljevic M, Popovic N. Renal function studies in individuals with the tubular proteinuria of endemic Balkan nephropathy. *Q J Med* 1972; 41: 385-393.
95. Radošević Z, Traeger J, Radonic M, Manuel Y, Revillard J P. Etude electrophoretique de la proteinurie de 31 sujet crates vivant en pays de nephropathie endemique. *J Urol Nephrol* 1968; 74: 703-710.
96. Bruckner I, Stoica Gh. Urinary proteins in renal diseases. II. Alpha<sub>2</sub>-microglobulins in tubular protienuria with special reference to endemic nephropathy. *Rev Roum Med Med Int* 1978; 16: 2-8.
97. Hall PW, Vasiljevic M. Beta2-microglobulin excretion as an index of renal tubular disorders with special reference to endemic Balkan nephropathy. *J Lab Clin Med* 1973; 81: 897-904.
98. Raicevic S, Trnacevic S, Hranisavljevic J, Vucelic D. Renal function, protein excretion, and pathology of Balkan endemic nephropathy. II. Protein excretion. *Kidney Int* 1991; 40 (Suppl 34): S52-S56.
99. Danilovic V. Endemic nephropathy in Yugoslavia. In: *Endemic (Balkan) Nephropathy*. Proc 4th Symp, Niš 1979. Strahinjic S, Stefanovic V (editors). *Inst Nephro Haemod, Niš* 1981; p. 1-5.
100. Pavlovic-Kentera V, Djukanovic LJ, Clemons GK, Trbojevic S, Dimkovic N, Slavkovic A. Anaemia in Balkan endemic nephropathy. *Kidney Int* 1991; 40 (Suppl 34): S46-S48.
101. Pavlovic-Kentera V, Clemons GK, Trbojevic S, Dimkovic N, Djukanovic LJ. Erythropoietin and anemia in the progression of Balkan endemic nephropathy and other renal diseases. *Nephron* 1990; 54: 139-143.
102. Cvorišec D, Stavljenic A, Radonic M. Relationship between tubular and Tamm-Horsfall proteinuria in Balkan endemic nephropathy. *Nephron* 1998; 42: 152-155.
103. Djukanovic LJ. Laboratory studies. In: *Endemic nephropathy*. Radovanovic Z, Sindjic M, Polenakovic M, Djukanovic LJ, Petronic V (editors). Office for Textbooks and Teaching Aids, Beograd 2000.
104. Polenakovic M, Stefanovic V. Balkan Nephropathy. In: *Oxford textbook of clinical nephrology*, Oxford medical Publications, Sec. Davison A, Cameron JS, Grunfeld JP, Kerr DNS, Ritz E, Winearls Ch (editors). Oxford University Press, Oxford 1998; p. 1202-1210.
105. Hrisoho D T. Balkan Endemic Nephropathy. Study of pathogenesis based on genetic factors. In: *Endemic (Balkan) Nephropathy*. Proc 4th Symp, Niš 1979. Strahinjic S, Stefanovic V (editors). *Inst Nephro Haemod, Niš* 1979; 1981; p. 147-154.
106. Elseviers MM, De Schepper A, Corthouts R, Bosmans JL, Cosyn L, Lins RL, Lornoy W, Matthys E, Roose R, Van Caesbroeck D, et al. High diagnostic performance of CT scan for analgesic nephropathy in patients with incipient to severe renal failure. *Kidney Int* 1995; 48(4): 1316-1323
107. Polenakovic M, Djukanovic LJ. Clinical picture, diagnostics and treatment of endemic nephropathy. In: *Endemic nephropathy*. Radovanovic Z, Sindjic M, Polenakovic M, Djukanovic LJ, Petronic V (editors). Belgrade: Office for Textbooks and Teaching Aids, 2000.