

Long Alternating Paths in Bicolored Point Sets

Jan Kynčl¹, János Pach^{2,*}, and Géza Tóth^{3,**}

¹ Department of Applied Mathematics and
Institute for Theoretical Computer Science,
Charles University, Prague, Czech Republic
jankyncl@centrum.cz

² City College, CUNY and Courant Institute of Mathematical Sciences,
New York University, New York, NY, USA
pach@cims.nyu.edu

³ Rényi Institute, Hungarian Academy of Sciences, Budapest, Hungary
geza@renyi.hu

Abstract. Given n red and n blue points in convex position in the plane, we show that there exists a noncrossing alternating path of length $n + c\sqrt{\frac{n}{\log n}}$. We disprove a conjecture of Erdős by constructing an example without any such path of length greater than $\frac{4}{3}n + c'\sqrt{n}$.

1 Introduction

It is a basic problem in geometric graph theory to decide which graphs can be drawn on a given point set with noncrossing straight-line edges. For instance, it is known that every outerplanar graph (i.e., triangulated cycle) G of n vertices can be drawn on any set of n points in general position in the plane [GMPP91]. Moreover, if G is a rooted tree, one can find such an embedding even if the image of the root is specified [IPTT94,T96]. An unsolved problem of this kind is to find the size of the smallest “universal” set in the plane, on which one can draw every planar graph of n vertices with noncrossing straight-line edges [dFPP90,CK89].

We obtain many interesting new questions by considering *colored* point sets; see [KK04] for a survey. It is a well known mathematics contest problem to prove that between any set R of n red and any set B of n blue points in general position in the plane there is a noncrossing matching, i.e., a one-to-one correspondence between their elements so that the segments connecting the corresponding point pairs are pairwise disjoint. Moreover, if R and B are separated by a line, one can also find an *alternating Hamilton path*, i.e., a noncrossing polygonal path passing through every element of $R \cup B$ such that any two consecutive vertices have opposite colors [AGH97]. If we do not assume that R and B are separated, then the last statement is known to be false for $n \geq 8$, even if $R \cup B$ is in *convex position*, i.e., its elements form the vertex set of a convex $2n$ -gon. The following problem was communicated to the second named author by Erdős around 1989.

* János Pach has been supported by NSF Grant CCR-00-98246, by PSC-CUNY Research Award 65392-0034, and by OTKA T-032452.

** Géza Tóth has been supported by OTKA-T-038397.

Problem. *Determine or estimate the largest number $\ell = \ell(n)$ such that, for every set of n red and n blue points on a circle, there exists a noncrossing alternating path consisting of ℓ vertices.*

Of course, the condition that the points are on a circle is equivalent to the assumption that they are in convex position.

Erdős and others conjectured that the asymptotically extremal configuration was the following. Suppose n is divisible by four. Cut the circle into four intervals, and place in them $\frac{n}{2}$ red, $\frac{n}{4}$ blue, $\frac{n}{2}$ red, and $\frac{3}{4}n$ blue points, respectively. It is easy to see that in this construction the number of vertices in the longest noncrossing alternating path is $\frac{3}{2}n + 2$. That is, we have $\ell(n) \leq \frac{3}{2}n + 2$. The main aim of this note is to disprove Erdős's conjecture by exhibiting a better construction in Section 2. A similar construction was found independently and at about the same time by Abellanas et al. [AGHT03].

From the other direction, it is easy to argue that $\ell(n) \geq n$. Indeed, divide the circle into two arcs, each containing n points. At least half of the points belonging to the first arc are of the same color, say, red. Then the second arc must contain the same number of blue points. Enumerate the red (resp. blue) points of the first (resp. second) arc in clockwise (resp. counterclockwise) order. Starting with the first red point on the first arc, and connecting each point with the next available element of opposite color on the other arc, we obtain a noncrossing alternating path of length $2\lceil \frac{n}{2} \rceil \geq n$. In Section 3, we improve this bound by a term that tends to infinity. Our results can be summarized as follows.

Theorem 1. *There exist constants $c, c' > 0$ such that*

$$n + c\sqrt{\frac{n}{\log n}} < \ell(n) < \frac{4}{3}n + c'\sqrt{n}.$$

It is an annoying feature of this problem that it is not clear whether the assumption that the points are in convex position plays any significant role. In particular, the above argument for finding an alternating path of length n easily generalizes to arbitrary 2-colored sets, on the other hand, our proof for the lower bound in Theorem 1 relies heavily on the fact that the points are in convex position. We conjecture that the upper bound in Theorem 1 is asymptotically tight, that is,

$$|\ell(n) - \frac{4}{3}n| = o(n).$$

See also our Conjecture at the end of the paper.

The problem of covering a set of n red and n blue points with *several* noncrossing alternating paths was discussed by Kaneko, Kano, and Suzuki [KKS04]. Alternating Hamiltonian *cycles* with at most $n - 1$ crossings were found by Kaneko, Kano, and Yoshimoto [KKY00]. Their result cannot be improved. Many other interesting questions about partitioning the plane into a given number of convex pieces, each containing roughly or exactly the same number of red and blue points, were studied in [BKS00, BM01, S02]. Analogous questions can be

asked when we color by red and blue all the $\binom{n}{2}$ segments between n points in general position in the plane. Furthermore, instead of long alternating paths, we may be interested in finding long *monochromatic* ones [KPTV98]. Merino et al. [MSU05] studied alternating paths in k -colored point sets for $k \geq 3$.

2 Upper Bound

Consider a ‘2-equicolored’ set C of $2n$ points in convex position in the plane. That is, let half of the elements of C be colored red and half of them blue. An *uninterrupted run* (or, in short, *run*) is a maximal set of consecutive points of C that have the same color. The *length* of a run is the number of its elements. We say that C is a *k-configuration* if it consists of k red and k blue runs.

A set of pairwise disjoint segments, each of which connects two points of different colors, is called a *matching*. The *size* of a matching is defined as the total number of points participating in it, that is, twice the number of segments. A matching is said to be *separated* if there is a straight-line that intersects the interior of each of its segments.

Lemma 2.1. *Let C be a k -configuration for some $k > 0$, which has a noncrossing alternating path of length l . Then C has a separated matching whose size is at least $l - 4k - 1$.*

Proof. Suppose without loss of generality that all elements of C lie on a circle. Consider a noncrossing alternating path p of length l . Fix a chord c of the circle that crosses the first and last segments along p , but does not pass through any point of C . Let M_1 denote the matching consisting of all odd-numbered segments of p . Clearly, the size of M_1 is at least $l - 1$. Let $M_2 \subseteq M_1$ be the set of all segments in M_1 that cross c . By definition, M_2 is a separated matching.

To establish the lemma, it is enough to show that the number of elements of M_1 that do not cross c is at most $2k$. Let us call these segments *outer* segments. For each pair of consecutive points of C that have different colors, pick a point between them on the circle. Any two consecutive runs of C are separated by at least one such point, so the number of points we selected is precisely $2k$. Every outer segment s divides the circle into two (closed) arcs. One of them, $I(s)$, contains both endpoints of c ; let the other one be denoted by $J(s)$. Since s connects two points of different colors, $J(s)$ must contain at least one of the selected points. On the other hand, both endpoints of the alternating path p belong to $I(s)$, so $J(s)$ cannot contain the endpoints of any outer segment other than the endpoints of s . Thus, for any two outer segments, s and s' , $J(s)$ and $J(s')$ are disjoint, so the selected points lying in the corresponding arcs $J(s)$ and $J(s')$, resp., are different. Hence, the number of outer segments cannot exceed the total number of selected points, which is $2k$. See Fig. 1. \square

Represent any k -configuration with runs S_1, S_2, \dots, S_{2k} by the sequence $(|S_1|, |S_2|, \dots, |S_{2k}|)$. We assume that the odd-numbered runs are red and the even-numbered are blue.




Fig. 1. $J(s)$ and $J(s')$ are disjoint.




Fig. 2. The k -configuration $(k \frac{n}{3k-2}, (2k-1) \frac{n}{3k-2}, k \frac{n}{3k-2}, \frac{n}{3k-2}, \dots, \frac{n}{3k-2})$.

Lemma 2.2. *Let $k \geq 2$ and assume that n is divisible by $3k - 2$. Then the size of any separated matching in the k -configuration $(k \frac{n}{3k-2}, (2k-1) \frac{n}{3k-2}, k \frac{n}{3k-2}, \frac{n}{3k-2}, \dots, \frac{n}{3k-2})$ is at most $2n \frac{2k-1}{3k-2}$.*

Proof. Let S_1, S_2, \dots, S_{2k} denote the consecutive runs of the k -configuration $(k \frac{n}{3k-2}, (2k-1) \frac{n}{3k-2}, k \frac{n}{3k-2}, \frac{n}{3k-2}, \dots, \frac{n}{3k-2})$ (see Fig. 2). Let M be a separated matching. We distinguish five cases, according to the set of runs that are connected to S_2 by at least one edge in M .

Case 1: No edge of M has an endpoint in S_2 .

Then M uses at most $n - n \frac{2k-1}{3k-2}$ blue points, so its size is at most $2n - 2n \frac{2k-1}{3k-2} = 2n \frac{k-1}{3k-2} < 2n \frac{2k-1}{3k-2}$.

Case 2: No edge of M runs between S_2 and $S_1 \cup S_3$.

Now the points in S_2 can be connected only to the elements of $S_5, S_7, \dots, S_{2k-1}$. Hence, at most $n \frac{k-2}{3k-2}$ points of S_2 are matched and at least $n \frac{k+1}{3k-2}$ are missed by M . The size of M is at most $2n - 2n \frac{k+1}{3k-2} < 2n \frac{2k-2}{3k-2}$.

Case 3: S_2 is connected by an edge of M to both S_1 and S_3 .

Since M is a separated matching, the blue points in S_4, S_6, \dots, S_{2k} are not matched, so M uses at most $n \frac{2k-1}{3k-2}$ blue points. Thus, the size of M is at most $2n \frac{2k-1}{3k-2}$.

Case 4: S_2 is connected by an edge of M to S_1 , but not to S_3 .

Suppose that the size of M exceeds $2n \frac{2k-1}{3k-2}$. Then M matches more than $n \frac{k}{3k-2}$ points of S_2 , so at least one edge of M must connect S_2 to a red run different from S_1 . Let $i > 1$ denote the smallest integer such that there is an edge of M between S_2 and S_{2i+1} . Then M matches at most $n \frac{k+(k-i)}{3k-2} = n \frac{2k-i}{3k-2}$ blue points from S_2 and misses the $n \frac{k-i}{3k-2}$ blue points in $S_{2i+2}, S_{2i+4}, \dots, S_{2k}$, because M is a separated matching. Therefore, it does not match at least $n \frac{(2k-1-(2k-i)+(k-i))}{3k-2} = n \frac{k-1}{3k-2}$ blue points, and its size is at most $2n \frac{2k-1}{3k-2}$.

Case 5: S_2 is connected by an edge of M to S_3 , but not to S_1 .

By symmetry, the same argument applies as in the previous case. □

Lemma 2.3. *For any positive integers k and $n \geq k$, there exists a k -configuration of $2n$ points with no alternating path longer than $2n \frac{2k-1}{3k-2} + 16k$.*

Proof. The statement is trivial for $k = 1$, and also for $n \leq 8k$. Suppose that $k \geq 2$, and $n > 8k$. Let $n_0 \leq n$ be the largest integer divisible by $3k - 2$. Let C_0 denote a k -configuration consisting of n_0 red and n_0 blue points, considered in Lemma 2.2. Add $n - n_0$ red points to S_1 and $n - n_0$ blue points to S_2 , and denote the resulting k -configuration by C .

We claim that C satisfies the requirement of the lemma. Let p be an alternating path of length $l(p)$ in C . By Lemma 2.1, there is a separated matching M_1 in C , whose size is $l(M_1) \geq l(p) - 4k - 1$. Remove from M_1 the $2n - 2n_0$ points that were added later and all edges in M_1 incident to them. We obtain a separated matching M_0 of C_0 of size $l(M_0) \geq l(M_1) - 4(n - n_0) \geq l(M_1) - 4(3k - 2) = l(M_1) - 12k + 8 \geq l(p) - 16k$. By Lemma 2.2, we have $l(M_0) \leq 2n_0 \frac{2k-1}{3k-2}$, so that $l(p) \leq 2n \frac{2k-1}{3k-2} + 16k$. □

The upper bound in Theorem 1 immediately follows from Lemma 2.3. For any n , set $k = \lfloor \sqrt{n} \rfloor$. Applying Lemma 2.3, we obtain a configuration of n red and n blue points in which the length of any noncrossing alternating path is at most $\frac{2k-1}{3k-2} \cdot 2n + 16k < \frac{4n}{3} + 20\sqrt{n}$, as required.

3 Lower Bound

As before, let C be the vertex set of a convex $2n$ -gon, with n red and n blue elements. Suppose without loss of generality that the elements of C lie on a

circle. A set of consecutive vertices of C (of not necessarily the same color) is said to be an *interval*. The *length* of an interval is its cardinality.

Assume that we can find a separated matching M of size $2l$, all of whose segments are crossed by a chord c . Then we can easily construct a noncrossing alternating path of length $2l$. To see this, enumerate the segments s_1, s_2, \dots, s_l of M according to the order of their intersection points with c . Let r_i and b_i be the red and blue endpoints of s_i , respectively. Then $p = (r_1 b_1, b_1 r_2, r_2 b_2, \dots, b_{l-1} r_l, r_l b_l)$ is a noncrossing alternating path of length $2l$.

Therefore, it is sufficient to establish a lower bound on the size of a separated matching in a k -configuration of $2n$ points. We divide the proof into two steps. Lemmas 3.1 and 3.2 provide reasonably good bounds when k is small and when k is large, respectively. Their combination implies the general lower bound in Theorem 1.

Lemma 3.1. *Let k, m, n be positive integers such that $k = 2^m$ divides n . Then every k -configuration C of $2n$ points contains a separated matching of size at least $n \left(1 + \frac{1}{k(m+1)}\right)$.*

Proof. Let S be the run of length at least $\frac{n}{k}$. Let I_0 denote a monochromatic interval in S , whose length is precisely $\frac{n}{k}$. For $1 \leq i \leq m+1$, let I_i be an interval of length $2^{i-1} \frac{n}{k}$ such that I_0, I_1, \dots, I_{m+1} are consecutive in the clockwise direction (see Fig. 3). These intervals form a partition of the underlying set C consisting of all $2n$ vertices. Assume without loss of generality that all elements of I_0 are blue. □

Suppose for contradiction that there is no separated matching whose size is at least $n \left(1 + \frac{1}{k(m+1)}\right)$.

Claim. *For every $0 \leq i \leq m+1$, the interval $J_i = I_0 \cup I_1 \cup \dots \cup I_i$ has at least $\left(2^{i-1} \frac{1}{k} + \frac{1}{2k} - \frac{i}{2k(m+1)}\right) n$ blue points. Moreover, strict inequality holds if $i > 0$.*

Note that Lemma 3.1 immediately follows from the Claim. Indeed, for $i = m+1$, we obtain that there are more than $\left(2^m \frac{1}{k} + \frac{1}{2k} - \frac{m+1}{2k(m+1)}\right) n = n$ blue points on the circle, which is a contradiction.

Proof of Claim. We proceed by induction on i . For $i = 0$, the statement obviously holds. Assume that for some $i \in \{0, 1, 2, \dots, m\}$, there are at least $\left(2^{i-1} \frac{1}{k} + \frac{1}{2k} - \frac{i}{2k(m+1)}\right) n$ blue points in the interval J_i . We show that there are more than $\left(2^{i-1} \frac{1}{k} - \frac{1}{2k(m+1)}\right) n$ blue points in J_{i+1} .




Fig. 3. I_0, I_1, \dots, I_{m+1} form a partition of the vertices.

Suppose this is not the case. Then there are at least $l = \left(2^{i-1}\frac{1}{k} + \frac{1}{2k(m+1)}\right)n$ red points in I_{i+1} . Since we have $2^{i-1}\frac{1}{k} + \frac{1}{2k} - \frac{i}{2k(m+1)} \geq 2^{i-1}\frac{1}{k} + \frac{1}{2k(m+1)}$, the number of blue points in J_i is at least l . Thus, there is a separated matching of size $2l$ between the blue points of J_i and the red points of I_{i+1} . Let M be the most ‘economical’ such matching of size $2l$. That is, if b_1, b_2, \dots, b_s ($s \geq l$) denote the blue points of J_i listed in counterclockwise order and r_1, r_2, \dots, r_t ($t \geq l$) denote the red points of I_{i+1} listed in clockwise order, then let M consist of the segments $b_1r_1, b_2r_2, \dots, b_lr_l$.

Let K denote the interval between b_l and r_l , oriented clockwise. All blue points from K not matched by M lie in I_{i+1} , so their number is at most $u_b = \left(2^{i-1}\frac{1}{k} - \frac{1}{2k(m+1)}\right)n$. All red points of K not matched by M lie in J_i , so their number is at most

$$\begin{aligned} u_r &= \left(2^i\frac{1}{k} - \left(2^{i-1}\frac{1}{k} + \frac{1}{2k} - \frac{i}{2k(m+1)}\right)\right)n \\ &= \left(2^{i-1}\frac{1}{k} - \frac{1}{2k} + \frac{i}{2k(m+1)}\right)n. \end{aligned}$$

Using the fact that $u_b - u_r = \left(\frac{1}{2k} - \frac{i+1}{2k(m+1)}\right)n \geq 0$, we obtain that $u_b \geq u_r$.

Let L denote the complement of K in the set C of all points. Clearly, L has at least $n_0 = n - l - u_b = \left(1 - 2^i\frac{1}{k}\right)n$ points of each color. Divide L into two intervals L_1, L_2 , each of length at least n_0 . Obviously, at least one of the following two conditions is satisfied:

- (a) There are at least $\frac{n_0}{2}$ blue points in L_1 and at least $\frac{n_0}{2}$ red points in L_2 .
- (b) There are at least $\frac{n_0}{2}$ red points in L_1 and at least $\frac{n_0}{2}$ blue points in L_2 .

We can assume without loss of generality that (a) is true. Then there exists a separated matching M' of size at least n_0 between intervals L_1 and L_2 . The union of M and M' is also a separated matching. (One endpoint of the corresponding chord lies between the intervals J_i and I_{i+1} , and the other between L_1 and L_2 .) The size of $M \cup M'$ is at least

$$2l + n_0 = \left(2^{i-1}\frac{1}{k} + \frac{1}{2k(m+1)}\right)2n + \left(1 - 2^i\frac{1}{k}\right)n = \left(1 + \frac{1}{k(m+1)}\right)n,$$

which is a contradiction. Hence, our assumption was wrong: there are more than $\left(2^{i-1}\frac{1}{k} - \frac{1}{2k(m+1)}\right)n$ blue points in I_{i+1} .

Consequently, the number of blue points in the interval $J_{i+1} = I_0 \cup I_1 \cup \dots \cup I_{i+1}$ is larger than

$$\begin{aligned} &\left(2^{i-1}\frac{1}{k} + \frac{1}{2k} - \frac{i}{2k(m+1)}\right)n + \left(2^{i-1}\frac{1}{k} - \frac{1}{2k(m+1)}\right)n \\ &= \left(2^i\frac{1}{k} + \frac{1}{2k} - \frac{i+1}{2k(m+1)}\right)n, \end{aligned}$$

completing the induction step, the proof of the Claim, and hence the proof of Lemma 3.1.

Lemma 3.2. *For $n \geq k \geq 1$, every k -configuration of $2n$ points admits an alternating path whose length is at least $n + k - 1$.*

Proof. Let v_1, v_2, \dots, v_{2n} be the vertices of a k -configuration, in clockwise direction. Assume that v_1 is red. For any $1 \leq i < 2n$, if v_i is red and v_{i+1} is blue, then v_i and v_{i+1} are called special vertices. There are $2k$ special and $2n - 2k$ non-special vertices. Moreover, exactly half of the special and exactly half of the non-special vertices are red, and blue. Let m be the smallest number such that there are $n - k$ non-special vertices in the set $\{v_i \mid 1 \leq i \leq m\}$. Assume that $t \geq \lceil \frac{n-k}{2} \rceil$ of them are red. (The other case can be settled analogously.) Denote those red points by u_1, u_2, \dots, u_t , in clockwise direction. Then the set $\{v_i \mid m+1 \leq i \leq 2n\}$ also contains $n - k$ non-special vertices, and t of them are blue. Denote those blue points by w_1, w_2, \dots, w_t , in counterclockwise direction.

The vertices u_1, u_2, \dots, u_t divide the set $\{v_i \mid 1 \leq i \leq m\}$ into $t + 1$ intervals of consecutive vertices, denote them by I_0, I_1, \dots, I_t , in clockwise direction. For $0 \leq l \leq t$, if I_l contains some special vertices, denote them by $u_{i,1}, u_{i,2}, \dots, u_{i,\alpha_i}$, in clockwise direction. Since u_1, u_2, \dots, u_t are non-special, $u_{i,j}$ is red if j is odd, and blue if j is even.

Similarly, the vertices w_1, w_2, \dots, w_t divide the set $\{v_i \mid m+1 \leq i \leq 2n\}$ into $t + 1$ intervals of consecutive vertices, denote them by J_0, J_1, \dots, J_t , in counterclockwise direction. For $0 \leq l \leq t$, if J_l contains some special vertices, denote them by $w_{i,1}, w_{i,2}, \dots, w_{i,\beta_i}$, also in counterclockwise direction. Now $w_{i,j}$ is blue if j is odd, and red if j is even.

Finally, consider the following path:

$$u_{0,1}, u_{0,2}, \dots, u_{0,\alpha_0}, u_1, w_{0,1}, w_{0,2}, \dots, w_{0,\beta_0}, w_1, u_{1,1}, u_{1,2}, \dots, u_{1,\alpha_1}, u_2, w_{1,1}, w_{1,2}, \dots, w_{1,\beta_1}, w_2, \dots, u_{t,1}, u_{t,2}, \dots, u_{t,(\alpha_t-1)}, w_{t,1}, w_{t,2}, \dots, w_{t,\beta_t}.$$

It is a noncrossing, alternating path of length $2t + 2k - 1 \geq n - k + 2k - 1 = n + k - 1$. This concludes the proof of the lemma. \square

Now we are ready to prove the lower bound in Theorem 1. Suppose that we have a k -configuration of $2n$ points. We distinguish two cases. If $k \geq \frac{1}{10} \sqrt{\frac{n}{\log n}}$, then, by Lemma 3.2, there exists an alternating path of length at least $n + \frac{1}{10} \sqrt{\frac{n}{\log n}} - 1$.

We are left with the case $k < \frac{1}{10} \sqrt{\frac{n}{\log n}}$. Let m be the least positive integer such that $k \leq 2^m$. Then $m < 1 + \log k$. Let $n' = 2^m \lfloor \frac{n}{2^m} \rfloor \geq n - 2^m$ and choose any subconfiguration C' of n' red and n' blue points from C . C' is a k_0 -configuration for some $k_0 \leq k$. So it has a run of length at least $\frac{n'}{k_0} \geq \frac{n'}{k} \geq \frac{n'}{2^m}$. Now, according to Lemma 3.1, C has a separated matching (and also an alternating path) whose size is at least

$$\begin{aligned} n' + \frac{n'}{2^m(m+1)} &\geq n - 2k + \frac{n}{2^{m+1}(m+1)} \geq \\ n - 2k + \frac{n}{4k(\log k + 2)} &\geq n - 2k + \frac{n}{2k \log n} \geq \\ n - \frac{1}{5} \sqrt{\frac{n}{\log n}} + \frac{5n}{\log n} \sqrt{\frac{\log n}{n}} &\geq n + 4 \sqrt{\frac{n}{\log n}}. \end{aligned}$$

This completes the proof of the lower bound in Theorem 1.

Conjecture. *For any fixed k and large n , every k -configuration of $2n$ points admits a separated matching of size at least $2n \frac{2k-1}{3k-2} + o(n)$.*

Acknowledgement

We are very grateful to the anonymous referee for his useful comments.

References

- [AGH97] M. Abellanas, J. Garcia, G. Hernández, M. Noy, and P. Ramos: Bipartite embeddings of trees in the plane, in: *Graph Drawing* (S. North, ed.), *Lecture Notes in Computer Science* **1190**, Springer-Verlag, Berlin, 1997, 1–10. Also in: *Discrete Appl. Math.* **93** (1999), 141–148.
- [AGHT03] M. Abellanas, A. Garcia, F. Hurtado, and J. Tejel: Caminos alternantes, in: *X Encuentros de Geometría Computacional* (in Spanish), Sevilla, 2003, 7–12.
- [BM01] I. Bárány and J. Matoušek: Simultaneous partitions of measures by k -fans, *Discrete Comput. Geom.* **25** (2001), 317–334.
- [BKS00] S. Bespamyatnikh, D. Kirkpatrick, and J. Snoeyink: Generalizing ham sandwich cuts to equitable subdivisions, *Discrete Comput. Geom.* **24** (2000), 605–622.
- [dFPP90] H. de Fraysseix, J. Pach, and R. Pollack: How to draw a planar graph on a grid, *Combinatorica* **10** (1990), 41–51.
- [CK89] M. Chrobak and H. Karloff: A lower bound on the size of universal sets for planar graphs, *SIGACT News* **20**/4 (1989), 63–86.
- [GMPP91] P. Gritzmann, B. Mohar, J. Pach, and R. Pollack: Embedding a planar triangulation with vertices at specified points (solution to Problem E3341), *Amer. Math. Monthly* **98** (1991), 165–166.
- [IPTT94] Y. Ikeba, M. Perles, A. Tamura, and S. Tokunaga: The rooted tree embedding problem into points on the plane, *Discrete Comput. Geom.* **11** (1994), 51–63.
- [KK04] A. Kaneko and M. Kano: Discrete geometry on red and blue points in the plane – a survey, in: *Discrete and Computational Geometry* (B. Aronov et al., eds.), Springer-Verlag, Berlin, 2004, 551–570.
- [KKS04] A. Kaneko, M. Kano, and K. Suzuki: Path coverings of two sets of points in the plane, in: *Towards a Theory of Geometric Graphs* (J. Pach, ed.), *Contemporary Mathematics* **342** (2004), 99–111.
- [KKY00] A. Kaneko, M. Kano, and K. Yoshimoto: Alternating Hamiltonian cycles with minimum number of crossings in the plane, *Internat. J. Comput. Geom. Appl.* **10** (2000), 73–78.
- [KPTV98] Gy. Károlyi, J. Pach, G. Tóth, and P. Valtr: Ramsey-type results for geometric graphs II, *Discrete Comput. Geom.* **20** (1998), 375–388.
- [MSU05] C. Merino, G. Salazar, and J. Urrutia: On the length of longest alternating paths for multicolored point sets in convex position, manuscript.
- [S02] T. Sakai: Balanced convex partitions of measures in R^2 , *Graphs and Combinatorics* **18** (2002), 169–192.
- [T96] S. Tokunaga: On a straight-line embedding problem of graphs, *Discrete Math.* **150** (1996), 371–378.