

17. Gonnella JD, Hornbrook MC, Louis DZ: Staging of disease: a case-mix measure. *Journal of the American Medical Association* 1984; 251: 637-644.
18. Special Committee on PTSD: *Outpatient Mental Health Treatment of Vietnam Era Veterans with Post-traumatic Stress Disorder*. Final Report. 1987. Washington, DC: Department of Veterans Affairs.
19. Kessler RC, Price RH, Wortman CB: Social factors in psychopathology: stress, social support and coping processes. *Annual Review of Psychology* 1986; 36:531-572.
20. Emrick CD, Hansen J: Assertions regarding effectiveness of treatment for alcoholism: fact or fantasy? *American Psychologist* 1983; 38:1078-1088.
21. Ogborne AC: Patient characteristics as predictors of treatment outcomes for alcohol and drug abusers. In: Israel V, Glaser F, Kalant H, et al. (eds.) *Research Advances in Alcohol and Drug Problems*. New York: Plenum Press, 1978, Vol. 4.
22. Gibbs L, Flanagan J: Prognostic indicators of alcoholism treatment outcome. *The International Journal of the Addictions* 1977; 12:1097-1141.
23. Hayes WL: *Statistics* (3rd ed.). New York: Holt, Rinehart and Winston, 1983.
24. Singer M: Evaluating alcoholism treatment programs: considerations and caveats. *The International Journal of the Addictions* 1983; 18:1145-1155.

Erratum

In: Boerstler H, de Figueiredo JM: Prediction of use of psychiatric services: application of the CART algorithm. *Journal of Mental Health Administration* 1991; 18(1):27-34. The percentage of high use for those whose most recent psychiatric treatment used was outpatient (Table 1) should be 47.1 and not 97.1.