

**Correction to ‘The Boolean algebra of spectra’**

**Commentarii Mathematici Helvetici 54/3 (1979) pp. 368–377**

A. K. BOUSFIELD

J. M. Boardman and others have pointed out an error in my proof of Proposition 1.5 [1]. Namely, the presentation of the CW-spectrum  $B_\lambda/A$  as the homotopy cofibre of  $1 - g$  on p. 371 is incorrect for a general limit ordinal  $\lambda$ . The error arose when I wrongly simplified an earlier proof, and the proposition remains valid. As suggested by Boardman, the proof can be repaired by using the equivalence of  $B_\lambda/A$  with the homotopy colimit  $C$  of the transfinite sequence  $\{B_s/A\}_{s < \lambda}$ . The required theory of homotopy colimits can be found in [3], [4], [5]. In more detail,  $C$  can be obtained by imposing appropriate face identifications on the wedge of the  $B_{s_0}/A \wedge (\Delta^n \cup *)$  running over all  $(n + 1)$ -tuples of ordinals  $s_0 < s_1 < \dots < s_n < \lambda$  for all  $n \geq 0$ . Thus,  $C$  is a CW-spectrum with an increasing filtration by closed subspectra  $\{F_n C\}$  such that  $F_n C/F_{n-1} C$  is the wedge of the  $B_{s_0}/A \wedge S^n$  running over all  $(n + 1)$ -tuples of ordinals  $s_0 < s_1 < \dots < s_n < \lambda$ . The associated spectral sequence for  $\pi_* C$  has  $E_{n,t}^2 \approx \text{colim}^n \{\pi_t B_s/A\}$ , and this derived colimit vanishes for  $n > 0$  because it is indexed by a directed set. Thus there is an edge isomorphism  $\text{colim}_{s < \lambda} \pi_* B_s/A \approx \pi_* C$  and the canonical map  $C \rightarrow B_\lambda/A$  is a weak equivalence of CW-spectra. Consequently  $C \approx B_\lambda/A$ . This equivalence can also be shown by using the isomorphisms  $\text{colim}_T \pi_* C_T \approx \pi_* C$  and  $\pi_* C_T \approx \pi_* B_{m(T)}/A$  where  $T$  runs over all finite nonempty sets of ordinals less than  $\lambda$ , where  $C_T \subset C$  is the homotopy colimit of the finite sequence  $\{B_s/A\}_{s \in T}$ , and where  $m(T)$  is the largest ordinal in  $T$ . Having shown  $C \approx B_\lambda/A$  one uses the Milnor cofibered  $\bigvee_{n \geq 0} F_n C \rightarrow \bigvee_{n \geq 0} F_n C \rightarrow C$  together with the above wedge decomposition of  $F_n C/F_{n-1} C$  to deduce that  $B_\lambda/A$  is  $[E, ]_*$ -colocal and belongs to Class-E as required for the proof of Proposition 1.5 [1] and for subsequent applications. A similar error appeared in the proof of Lemma 1.13 [2] and can be repaired similarly.

REFERENCES

[1] A. K. BOUSFIELD, *The Boolean algebra of spectra*. Comment. Math. Helvet. 54 (1979), 368–377.  
 [2] A. K. BOUSFIELD, *The localization of spectra with respect to homology*. Topology 18 (1979), 257–281.

- [3] A. K. BOUSFIELD and D. M. KAN, *Homotopy limits, completions and localizations*. Lecture Notes in Mathematics Vol. 304, Springer-Verlag, New York, 1972.
- [4] G. SEGAL, *Classifying spaces and spectral sequences*. Publ. Math. I.H.E.S. 34 (1968), 105–112.
- [5] R. M. VOGT, *Homotopy limits and colimits*. Math. Z. 134 (1973), 11–52.

*University of Illinois at Chicago*  
*Dept of Mathematics*  
*P.O. Box 4348*  
*Chicago Ill 60680/USA*

Received April 8, 1983