

A Simple and Relatively Efficient Triangulation of the n -Cube*

Mark Haiman

Department of Mathematics, Massachusetts Institute of Technology,
 77 Massachusetts Avenue, Cambridge, MA 02139, USA

Abstract. The only previously published triangulation of the n -cube using $o(n!)$ simplices, due to Sallee, uses $O(n^{-2}n!)$ simplices. We point out a very simple method of achieving $O(\rho^n n!)$ simplices, where $\rho < 1$ is a constant.

1. Introduction

This short note is intended to point out a simple method of triangulating the n -cube I^n using significantly fewer simplices than in previous constructions.

Various authors [2], [3], [5]–[8] have considered the problem of triangulating I^n into fewer than the easily achievable maximum of $n!$ simplices. Since the volume of a simplex with vertices in \mathbf{Z}^n is an integral multiple of $1/n!$, it is clear that $n!$ is in fact the maximum number of simplices in any triangulation. A lower bound can also be derived from volume considerations as follows. I^n can be inscribed in a sphere of diameter \sqrt{n} . The maximum volume of a simplex contained in this sphere is $(n+1)^{(n+1)/2}/2^n n!$, attained by the equilateral simplex. This shows that any triangulation of I^n uses at least $2^n(n+1)^{-(n+1)/2}n!$ simplices. This lower bound is very much less than $n!$, being $O(c^n(n!)^{1/2})$. By replacing the cube with an “ideal cube” in hyperbolic space and using hyperbolic instead of Euclidean volume it is possible to improve the lower bound [8], but only by a factor of $O((3/2)^{n/2})$.

In view of the large gap between the lower and upper bounds it is perhaps surprising that all triangulations of I^n proposed so far use nearly $n!$ simplices. In fact, only Sallee [7] achieves $o(n!)$ simplices. Sallee’s triangulation, however, uses more than $2A(n-1, \lfloor (n-1)/2 \rfloor)$ simplices, where the *Eulerian number* $A(n, k)$ is the number of permutations of n having k “descents” [10]. Since $A(n, k)$ is unimodal as a function of k , $A(n-1, \lfloor (n-1)/2 \rfloor) \geq (n-1)!/(n-1) = (n-2)!$

* This research was supported in part by N.S.F. Grant No. DMS-8717795.

and hence Sallee uses at least $O(n^{-2}n!)$ simplices. (Readers of [7] should observe that the function $g(n, m)$ of that paper is actually $A(n - 1, m - 1)$, a fact not noted there.)

The construction given below triangulates I^n with $O(\rho^n n!)$ simplices, where $\rho < 1$. In fact, given a triangulation of I^n into $T(n)$ simplices for any particular value of n , we can take $\rho = (T(n)/n!)^{1/n}$. The smallest value for ρ obtainable from triangulations published to date is $\rho = (13,248/40,320)^{1/8} \approx 0.870$ from Sallee's triangulation of I^8 . We remark that Todd [11] proposed the quantity $R(n) = (T(n)/n!)^{1/n}$ as a measure of a triangulation's efficiency. Previous constructions have $\lim_{n \rightarrow \infty} R(n) = 1$, whereas our results show that any value of $R(n)$ achievable for one triangulation is achievable asymptotically. This is still far from $R(n) = o(1)$, let alone the lower bound of $R(n) = O(n^{-1/2})$.

2. Construction

Definition. A polyhedral decomposition of an n -dimensional polytope P is a union $P = T_1 \cup T_2 \cup \dots \cup T_k$ of n -dimensional polytopes T_i such that for all i, j the vertices of T_i are vertices of P and $T_i \cap T_j$ is a (possibly empty) face of both T_i and T_j . If each T_i is a simplex, then $\{T_i\}$ is a triangulation of P .

Lemma 1. Every polyhedral decomposition of P can be refined to a triangulation.

Proof. We require triangulations θ_i of the T_i such that θ_i and θ_j induce the same triangulation on $T_i \cap T_j$ considered as a face of T_i and as a face of T_j . Now there are well-known constructions [4], [6], [9], [12] whereby we associate to a total ordering α of the vertices of a polytope T a triangulation θ in such a way that the triangulation induced on each face $F \subseteq T$ is the one associated to the restriction of α to the vertices of F . Hence fixing any total ordering α_0 of all the vertices of P and triangulating each T_i in accordance with α_0 we obtain compatible triangulations θ_i as required. □

Lemma 2 [1]. Every triangulation of $\Delta_k \times \Delta_l$ uses exactly $(k + l)!/k!l!$ simplices, where Δ_n denotes an n -dimensional simplex.

Proof. Realize Δ_k in \mathbf{R}^k as the convex hull of 0 and the unit coordinate vectors $e_j = (0, \dots, 0, 1, 0, \dots, 0)$. Likewise Δ_l . Then $\Delta_k \times \Delta_l \subseteq \mathbf{R}^{k+l}$ has vertices 0, e_i ($1 \leq i \leq k + l$), and $e_i + e_j$ ($1 \leq i \leq k < j \leq k + l$). Its volume is $v(\Delta_k)v(\Delta_l) = 1/k!l!$. We claim every nondegenerate $(k + l)$ -simplex Δ spanned by vertices of $\Delta_k \times \Delta_l$ has volume $1/(k + l)!$. Note that there are affine-linear symmetries of $\Delta_k \times \Delta_l$ acting transitively on the vertices. These preserve volume, so we can assume 0 is a vertex of Δ . Then $\pm(k + l)!v(\Delta)$ is the determinant of the matrix M whose rows are the coordinates of the other vertices; we are to show that this determinant is ± 1 . If some e_i is a vertex of Δ , then expanding by minors on the corresponding row gives the result by induction. If not, then M is the edge-vertex incidence matrix of a (k, l) -bipartite graph with $k + l$ edges. Having one too many

edges to be a tree, this graph must contain a cycle, necessarily even. But then M is singular, contrary to hypothesis. \square

Theorem 1. *Given a triangulation $\{S_1, \dots, S_s\}$ of a k -dimensional polytope P and a triangulation $\{T_1, \dots, T_t\}$ of an l -dimensional polytope Q , there exists a triangulation of $P \times Q$ using $s \cdot t \cdot (k + l)!/k! l!$ simplices.*

Proof. It is easy to see that $\{S_i \times T_j\}$ is a polyhedral decomposition of $P \times Q$. Refine it to a triangulation by Lemma 1. Each $S_i \times T_j$ will then contain $(k + l)!/k! l!$ simplices by Lemma 2, establishing the result. \square

Corollary 1. *If I^n can be triangulated into $T(n)$ simplices, then I^{kn} can be triangulated into $[(kn)!/(n!)^k]T(n)^k = \rho^{kn}(kn)!$ simplices, where $\rho = (T(n)/n!)^{1/n}$.*

Proof. Immediate from Theorem 1 by induction on k . \square

References

1. L. J. Billera, R. Cushman, and J. A. Sanders, The Stanley decomposition of the harmonic oscillator, *Nederl. Akad. Wetensch. Proc. Ser. A* **91** (1988), 375–393.
2. R. W. Cottle, Minimal triangulations of the 4-cube, *Discrete Math.* **40** (1982), 25–29.
3. C. Lee, Triangulating the d -cube, in *Discrete Geometry and Convexity*, J. E. Goodman, E. Lutwak, J. Malkevitch, and R. Pollack, eds., New York Academy of Sciences, New York (1985), pp. 205–211.
4. C. Lee, Some notes on triangulating polytopes, in *Proc. 3. Kolloquium über Diskrete Geometrie*, Institut für Mathematik, Universität Salzburg (1985), pp. 173–181.
5. P. S. Mara, Triangulations of the Cube, M. S. Thesis, Colorado State University (1972).
6. J. F. Sallee, A triangulation of the n -cube, *Discrete Math.* **40** (1982), 81–86.
7. J. F. Sallee, Middle-cut triangulations of the n -cube, *SIAM J. Algebraic Discrete Methods* **5**, no. 3 (1984), 407–419.
8. W. D. Smith, Polytope triangulations in d -space, improving Hadamard's inequality and maximal volumes of regular polytopes in hyperbolic d -space. Manuscript, Princeton, NJ (September 1987).
9. R. P. Stanley, Decompositions of rational convex polytopes, *Ann. Discrete Math.* **6** (1980), 333–342.
10. R. P. Stanley, *Enumerative Combinatorics*, Vol. I, Wadsworth & Brooks/Cole, Monterey, CA (1986).
11. M. J. Todd, *The Computation of Fixed Points and Applications*, Lecture Notes in Economics and Mathematical Systems, Vol. 124, Springer-Verlag, Berlin (1976).
12. B. Von Hohenbalken, How To Simplicially Partition a Polytope, Research Paper No. 79–17, Department of Economics, University of Alberta, Edmonton (1979).

Received September 21, 1989, and in revised form November 27, 1989.