

39. Harper, J. F.; Brooker, G. Femtomole sensitive radioimmunoassay for cyclic AMP and cyclic GMP after 2'0 acetylation by acetic anhydride in aqueous solutions. *J. Cyclic Nucleotide Res.* 1: 207-218; 1975.
40. Steiner, A. L.; Pagliara, A. S.; Chase, L. R.; Kipnis, D. M. Radioimmunoassay for cyclic nucleotides. II. Adenosine 3', 5'-monophosphate and guanosine 3', 5'-monophosphate in mammalian tissues and body fluids. *J. Biol. Chem.* 247: 1114-1120; 1972.
41. Johnson, A. R. Human pulmonary endothelial cells in culture: Activities of cells from arteries and cells from veins. *J. Clin. Invest.* 65: 841-850; 1980.
42. Murray, J. F. *The normal lung: The basis for diagnosis and treatment of pulmonary disease.* Philadelphia, PA: W. B. Saunder Co.; 1976.
43. Gimbrone, M. A., Jr.; Alexander, R. W. Angiotensin II stimulation of prostaglandin production in cultured human vascular endothelium. *Science* 189: 219-220; 1975.
44. Alexander, R. W. Hormone receptors in cultured vascular cells. Bevan, J. A.; Maxwell, R. A.; Godfraind, T.; Vanhoutte, P. M. eds. *Vascular neuroeffector mechanisms.* New York: Raven Press; 1980: 15-16.

This work was supported by a Young Investigator Grant HL-21189 from the National Institutes of Health, United States Public Health Service.

A preliminary report of this study appeared in *In Vitro* 16: 259; 1980.

Note Added in Proof: During the time the present paper was being considered for publication two other reports have appeared on the stimulation of cyclic AMP levels by catecholamines and prostaglandins (prostaglandin I₂) in cultures of human umbilical vein and bovine aortic endothelial cells.

Schafer, A. I.; Gimbrone, M. A., Jr.; Handin, R. I. Endothelial cell adenylate cyclase: Activation by catecholamines and prostaglandin I₂. *Biochem. & Biophys. Res. Commun.* 96: 1640-1647; 1980.

Hopkins, N. K.; Gorman, R. R. Regulation of endothelial cell cyclic nucleotide metabolism by prostacyclin. *J. Clin. Invest.* 67: 540-546; 1981.

ERRATUM

The cell line described in the manuscript by Meyskens et al., *Biological and Biochemical Properties of the Human Uveal Melanocyte Derived Cell Line (In Vitro* 16: 775-780; 1980) is most likely of rat origin. The source and time of this contamination is not presently known since we do not carry rat lines in our laboratory. The exact identification of this cell line is continuing.