

- d. Math. 24 (1978), pp. 427-436.
- [15] J. R. Moschovakis: A topological interpretation of second order intuitionistic arithmetic, *Compositio Math.* 26 (1973), pp. 261-275.
- [16] J. Myhill: Formal systems for intuitionistic analysis I, in: *Proceedings of 3<sup>rd</sup> International Congress of Logic, Methodology and Philosophy of Science*, ed. by B. van Rootselaer and J. F. Staal, North-Holland, Amsterdam, 1968.
- [17] J. Myhill: Formal systems for intuitionistic analysis II, in: *Intuitionism and Proof Theory*, ed. by A. Kino, J. Myhill, R. E. Vesley, North-Holland, Amsterdam, 1970.
- [18] J. Myhill: Constructive set theory, *JSL* 40 (1975), pp. 347-382.
- [19] J. Myhill; A. Scedrov: Second-order intuitionistic propositional calculus, to appear.
- [20] D. S. Scott: Extending the topological interpretation to intuitionistic analysis, I. *Compositio Math.* 20 (1968), pp. 194-210.
- [21] D. S. Scott: Extending the topological interpretation to intuitionistic analysis, II. in: *Intuitionism and Proof Theory*, ed. by A. Kino, J. Myhill, R. E. Vesley, North Holland, Amsterdam, 1970.
- [22] W. C. Powell: *Seminar Notes*, Ohio State University, 1977.
- [23] M. Tierney: *Sheaf Theory and Continuum Hypothesis*, in: *Toposes, Algebraic Geometry and Logic*, ed. by F. W. Lawvere, Springer Lecture Notes in Mathematics 274, Berlin, 1972.

#### ERRATA

In the proof of Theorem 1.9., validity of RDC should be established as follows: Let  $f, g, h$  be mutually disjoint rare sequences, where  $G_f \subseteq \text{stab}(z_i)$  for all parameters  $z_i$  in an instance of RDC (including  $X$ ). Let  $k_i$  ( $i \geq 1$ ) be mutually disjoint (rare) subsequences of  $h$ , and let  $h_i = k_1 * k_2 * \dots * k_i$ . Then in the antecedent,  $x$  ranges over  $\text{dom}(X)$  so that  $G_{f * g} \subseteq \text{stab}(x)$ , but  $y$  ranges over  $\text{dom}(X)$  so that  $G_{f * g * h_1} \subseteq \text{stab}(y)$ . Also,  $G_{f * g * h_i} \subseteq \text{stab}(z_{\langle j_1, \dots, j_i \rangle}) \cap \text{stab}(\delta_{\langle j_1, \dots, j_i \rangle}^i)$ , and  $G_{f * g * h} \subseteq \text{stab}(f_s^X)$ , so  $f_s^{X \in V(\Gamma)}$ .