

A comprehensive framework for studying W' and Z' bosons at hadron colliders with automated jet veto resummation

Benjamin Fuks^{a,b,c} and Richard Ruiz^d

^a*Sorbonne Universités, UPMC Univ. Paris 06, UMR 7589, LPTHE, F-75005, Paris, France*

^b*CNRS, UMR 7589, LPTHE, F-75005, Paris, France*

^c*Institut Universitaire de France, 103 boulevard Saint-Michel, 75005 Paris, France*

^d*Institute for Particle Physics Phenomenology (IPPP),*

Department of Physics, Durham University,

Durham, DH1 3LE, U.K.

E-mail: fuks@lpthe.jussieu.fr, richard.ruiz@durham.ac.uk

ABSTRACT: The production of high-mass, color-singlet particles in hadron collisions is universally accompanied by initial state QCD radiation that is predominantly soft with respect to the hard process scale Q and/or collinear with respect to the beam axis. At TeV-scale colliders, this is in contrast to top quark and multijet processes, which are hard and central. Consequently, vetoing events with jets possessing transverse momenta above p_T^{Veto} in searches for new color-singlet states can efficiently reduce non-singlet backgrounds, thereby increasing experimental sensitivity. To quantify this generic observation, we investigate the production and leptonic decay of a Sequential Standard Model W' boson at the 13 TeV Large Hadron Collider. We systematically consider signal and background processes at next-to-leading-order (NLO) in QCD with parton shower (PS) matching. For color-singlet signal and background channels, we resum Sudakov logarithms of the form $\alpha_s^j(p_T^{\text{Veto}}) \log^k(Q/p_T^{\text{Veto}})$ up to next-to-next-to-leading logarithmic accuracy (NNLL) with NLO matching. We obtain our results using the MADGRAPH5_AMC@NLO and MADGRAPH5_AMC@NLO-SCET frameworks, respectively. Associated Universal FEYNRULES Output model files capable of handling NLO+PS- and NLO+NNLL-accurate computations are publicly available. We find that within their given uncertainties, both the NLO+PS and NLO+NNLL(veto) calculations give accurate and consistent predictions. Consequently, jet vetoes applied to color-singlet processes can be reliably modeled at the NLO+PS level. With respect to a b -jet veto of $p_T^{\text{Veto}} = 30$ GeV, flavor-agnostic jet vetoes of $p_T^{\text{Veto}} = 30 - 40$ GeV can further reduce single top and $t\bar{t}$ rates by a factor of 2-50 at a mild cost of the signal rate. Jet vetoes can increase the signal-to-noise ratios by roughly 10% for light W' boson masses of 30 – 50 GeV and 25%-250% for masses of 300-800 GeV.

KEYWORDS: Jets, NLO Computations

ARXIV EPRINT: [1701.05263](https://arxiv.org/abs/1701.05263)

Contents

1	Introduction	1
2	An effective framework for W' and Z' studies at colliders	3
2.1	A simplified model for extra gauge boson searches	3
2.2	Computational setup	8
3	Jet veto resummation	9
3.1	Jet veto resummation at next-to-next-to-leading logarithmic accuracy with next-to-leading order matching	9
3.2	Non-perturbative corrections to cross sections with jet vetoes	12
3.3	Scale uncertainties in resummed jet veto rates from varying jet definitions	14
4	Signal and background process modeling with jet vetoes	16
4.1	W' production and decay	16
4.2	SM Drell-Yan continua	18
4.3	Top quark background	20
5	Missing transverse energy and jet modeling	22
5.1	Exclusive and inclusive missing transverse momentum	22
5.2	Exclusive and inclusive jet veto	24
6	Observability of W_{SSM} with jet vetoes at hadron colliders	27
6.1	Signal definition and event selection	28
6.2	Extended discovery potential and sensitivity at 13 TeV	29
7	Summary and conclusion	31

1 Introduction

The existence of new massive, colorless vector bosons is a key prediction of many theories that address the empirical and theoretical shortcomings of the Standard Model (SM) of particle physics. This, for instance, includes dark photons and Z'_D bosons in dark matter models, W_R^\pm and Z_R gauge bosons in left-right symmetric models, Z'_{B-L} bosons in neutrino mass models, or W_{KK}^\pm , Z_{KK} and γ_{KK} Kaluza-Klein excitations in extra-dimension models. These bosons are generically referred to as W'^\pm and Z' bosons [1]. Searches for these particles are and will continue to be an integral part of the Large Hadron Collider (LHC) physics program. Subsequently, an ability to categorically increase the experimental sensitivity of such searches is desirable.




Figure 1. Born-level Feynman diagrams for W' boson (left) and Z' -boson (right) production and decay into leptons in hadronic collisions. All figures are drawn with JAXODRAW [8].

A powerful and robust LHC test of these models consists of reinterpreting searches for leptonic decays of Sequential Standard Model (SSM) W_{SSM}^{\pm} and Z_{SSM} bosons [2–7], which proceed through the Drell-Yan (DY) processes

$$q\bar{q}' \rightarrow W_{\text{SSM}}^{\pm} \rightarrow \ell^{\pm}\nu_{\ell} \quad \text{and} \quad q\bar{q} \rightarrow Z_{\text{SSM}} \rightarrow \ell^{+}\ell^{-} \quad \text{with } q, q' \in \{u, d, c, s, b\} \text{ and } \ell \in \{e, \mu, \tau\}, \quad (1.1)$$

and whose leading order (LO) diagrams are shown in figure 1. SSM bosons couple to SM fermions in the same manner as the SM W^{\pm} and Z bosons up to overall coupling normalizations and may thus possess couplings that radically differ from any of the aforementioned models. It is nonetheless straightforward to reinterpret the SSM collider limits on masses and couplings within another theoretical framework.

The collider signatures relevant for the benchmark processes of eq. (1.1) consist of final states made of one or two charged leptons (ℓ^{\pm}) with a large transverse momentum (p_T), and additionally accompanied by a large amount of missing transverse energy (\cancel{E}_T) in the W_{SSM} case. The dominant (and irreducible) backgrounds are thus the charged current and neutral current DY continua. However, as in most hadron collider searches, SSM searches are *inclusive* with respect to jet and soft QCD activity, so that high- p_T multijet and top quark processes, with cross sections many orders of magnitude larger than the SSM boson production rates, also contribute to the SM background.

Intriguingly, s -channel W_{SSM} and Z_{SSM} production are color-singlet $q\bar{q}$ annihilation processes, meaning that QCD radiation off the initial-state quarks is favorably soft with respect to the hard process scale $Q \sim M_{W'/Z'}$ and/or collinear with respect to the beam axis. This implies that the corresponding jet activity is inherently softer and more collinear than for the leading QCD backgrounds. In particular, top quark decay products characteristically feature large momenta scales of $p_T \sim 50 - 60$ GeV, which suggests that inclusive, i.e., flavor-independent, jet vetoes, even loose ones, can enhance signal-to-noise ratios in SSM boson searches.

Historically, such arguments were made first for the vector boson fusion (VBF) process [9]. More recently, they have also been discussed in DY, non-VBF Higgs and multi-boson production channels [10–26], as well as in several searches for physics beyond the Standard Model (BSM) [27–30]. We argue, however, that due to the universality of QCD radiation in the soft and collinear limits, jet vetoes are in fact generically applicable to *any* color-singlet process that results in colorless final-state particles and in which QCD processes are a non-negligible fraction of the background.

As learned from measurements of the (inclusive) $W^+W^- + 0j$ cross section at the LHC [15–18, 31–34], reliable predictions for color-singlet processes initiated by $q\bar{q}$ annihilation and gluon fusion on which a jet veto of scale p_T^{Veto} is applied require the resummation of Sudakov logarithms of the form $\alpha_s(p_T^{\text{Veto}}) \log(Q/p_T^{\text{Veto}})$. These logarithms originate from t -channel propagators corresponding to initial-state gluon radiation and spoil the convergence of the perturbative series when the hard process mass scale Q is much greater than the veto scale p_T^{Veto} . Indeed, the lowest order at which the p_T spectrum of any color-singlet system is even qualitatively accurate everywhere is at next-to-leading order (NLO) in QCD for the inclusive process matched to the leading logarithmic (LL) resummation of the recoiling radiation momenta [35]. Formally, this is the same accuracy as inclusive NLO calculations matched with presently available parton shower (PS). Recently, jet veto resummation at next-to-next-to-leading (NNLL) logarithmic accuracy has been automated [18]. In the latter case, resummed computations for generic color-singlet processes are matched with the NLO fixed order results within the MADGRAPH5_aMC@NLO (MG5aMC) platform [36] and the Soft-Collinear Effective Theory [37–39] (SCET) formalism.

In this study, we investigate the impact of (resummed) jet vetoes on current and future searches for generic W' and Z' bosons both at the LHC and at hypothetical very large hadron colliders [40, 41], such as the proposed Future Circular Collider (FCC) at CERN [42] or the Circular pp Collider (CppC) at IHEP [43]. We focus, as a benchmark scenario, on the case of a W_{SSM} boson, and perform our study on the basis of the automated resummation technology implemented within the MG5aMC+SCET framework. We compare our veto resummed results to those obtained at NLO+PS accuracy. Our work includes the construction of a new general FEYNRULES [44] model for extra gauge boson studies that can be interfaced with NLOCT [45] and FEYNARTS [46]. The associated Universal FEYNRULES Output (UFO) [47] model is publicly available from the FEYNRULES model database [48] and can be used to simulate hadronic and leptonic collisions up to NLO+PS accuracy.

The remainder of this study continues in the following order: in section 2, we present the theoretical framework for extending the SM field content extra gauge boson, summarize current LHC constraints on new vector bosons, and provide details on our computational setup. In section 3, we briefly review jet veto resummation within the SCET formalism, and discuss rate uncertainties associated with mass, p_T^{Veto} , and jet radius scale choices in W' boson production. Signal and background modeling with jet vetoes is described in section 4, and we focus particularly in section 5 on the modeling of the missing energy and jet properties in $W' \rightarrow e/\mu + \cancel{E}_T$ searches. We then dedicate section 6 to estimating the improved discovery for W_{SSM} boson searches gained by applying jet vetoes and finally summarize and conclude in section 7.

2 An effective framework for W' and Z' studies at colliders

2.1 A simplified model for extra gauge boson searches

We take a simplified approach to modeling physics beyond the Standard Model. We do this by minimally extending the SM to construct a general effective framework that can be used for studying various models featuring extra colorless gauge bosons that couple to

Charge	u_L	d_L	ν_L	e_L	u_R	d_R	e_R
$T_L^{3,f}$	$+\frac{1}{2}$	$-\frac{1}{2}$	$+\frac{1}{2}$	$-\frac{1}{2}$	0	0	0
Q^f	$+\frac{2}{3}$	$-\frac{1}{3}$	0	-1	$+\frac{2}{3}$	$-\frac{1}{3}$	-1

Table 1. Weak isospin and electric charge assignments for the left-handed and right-handed chiral fermions f_L and f_R entering the Z_{SSM} vector and axial-vector couplings of eq. (2.3).

SM fermions. Specifically, we supplement the SM field content by two massive, colorless vector fields W'^{\pm} and Z' that are respectively electrically charged and neutral. To ensure model independence, the exact form of the W' and Z' chiral couplings to SM fermions is not specified, and any interaction of the new vector bosons with other gauge or scalar bosons is omitted. Following refs. [49, 50], the Lagrangian parameterizing the new vector bosons' couplings to up-type and down-type quark fields u_i and d_j is given by

$$\begin{aligned} \mathcal{L}_{\text{NP}}^q = & -\frac{g}{\sqrt{2}} \sum_{i,j} \left[\bar{u}_i V_{ij}^{\text{CKM}} W_{\mu}^{\prime+} \gamma^{\mu} \left(\kappa_L^q P_L + \kappa_R^q P_R \right) d_j + \text{H.c.} \right] \\ & - \frac{g}{\cos \theta_W} \sum_{q=u,d} \sum_i \left[\bar{q}_i Z'_{\mu} \gamma^{\mu} \left(\zeta_L^q P_L + \zeta_R^q P_R \right) q_i \right], \end{aligned} \quad (2.1)$$

where i and j denote flavor indices, $P_{L/R} = \frac{1}{2}(1 \mp \gamma_5)$ are the usual left/right-handed chirality projectors, V^{CKM} is the Cabbibo-Kobayashi-Maskawa (CKM) matrix, and g and θ_W are the weak coupling constant and mixing angle respectively. We choose coupling normalizations facilitating the mapping to the reference SSM Lagrangian \mathcal{L}_{SSM} [1]. The real-valued quantities $\kappa_{L,R}^q$ and $\zeta_{L,R}^q$ serve as overall normalization of the new interactions relative to the strength of the weak coupling constant. We do not assume additional sources of flavor violation beyond the SM CKM matrix.

Similarly, the interactions involving charged lepton ℓ and massless neutrino ν_{ℓ} fields are parametrized by [49, 50]

$$\begin{aligned} \mathcal{L}_{\text{NP}}^{\ell} = & -\frac{g}{\sqrt{2}} \sum_i \left[\bar{\nu}_{\ell i} W_{\mu}^{\prime+} \gamma^{\mu} \kappa_L^{\ell} P_L \ell_i^{-} + \text{H.c.} \right] \\ & - \frac{g}{\cos \theta_W} \sum_{f=\ell, \nu_{\ell}} \sum_i \left[\bar{f}_i Z'_{\mu} \gamma^{\mu} \left(\zeta_L^f P_L + \zeta_R^f P_R \right) f_i \right]. \end{aligned} \quad (2.2)$$

The quantities κ_L^{ℓ} are real-valued and serve as normalizations for leptonic coupling strengths. As no right-handed neutrinos are present in the SM, the corresponding right-handed leptonic new physics couplings are omitted ($\zeta_R^{\nu} = \kappa_R^{\ell} = 0$). We assume that leptonic interactions are flavor diagonal.

From our general Lagrangians of eq. (2.1) and eq. (2.2), the SSM limit is obtained by imposing the coupling strengths to be equal to the SM weak couplings up to an overall normalization factor,

$$\zeta_{R,L}^f = \zeta_{\text{SSM}}^f \left(g_V^f \pm g_A^f \right) \quad \text{with} \quad g_V^f = \frac{1}{2} T_L^{3,f} - Q^f \sin^2 \theta_W \quad \text{and} \quad g_A^f = -\frac{1}{2} T_L^{3,f}, \quad (2.3)$$

Gauge group	Charge	u_L	d_L	ν_L	e_L	u_R	d_R	N_R	e_R
$SU(2)_L$	$T_L^{3,f}$	$+\frac{1}{2}$	$-\frac{1}{2}$	$+\frac{1}{2}$	$-\frac{1}{2}$	0	0	0	0
$SU(2)_R$	$T_R^{3,f}$	0	0	0	0	$+\frac{1}{2}$	$-\frac{1}{2}$	$+\frac{1}{2}$	$-\frac{1}{2}$
$U(1)_{EM}$	Q^f	$+\frac{2}{3}$	$-\frac{1}{3}$	0	-1	$+\frac{2}{3}$	$-\frac{1}{3}$	0	-1

Table 2. Weak isospin and electric charge assignments for the left-handed and right-handed chiral fermions f_L and f_R entering the Z_R couplings of eq. (2.5). Right-handed neutrinos N_R are included for completion.

where the quantum number assignments are listed in table 1. In the canonical SSM, the overall normalizations are further trivially fixed as

$$\kappa_L^{q,\ell} = 1, \quad \kappa_R^q = 0 \quad \text{and} \quad \zeta_{Z_{SSM}}^f = 1. \quad (2.4)$$

This parameterization can be used to describe any model featuring extra colorless, massive vector bosons, provided there is no new source of flavor violation with respect to the SM. For instance, right-handed W_R and Z_R boson interactions can be obtained by enforcing

$$\begin{aligned} \zeta_L^f &= \frac{\kappa_R^f \cos \theta_W}{\sqrt{1 - \frac{\tan^2 \theta_W}{(\kappa_R^f)^2}}} \frac{\tan^2 \theta_W}{(\kappa_R^f)^2} \left[T_L^{3,f} - Q^f \right] \\ \zeta_R^f &= \frac{\kappa_R^f \cos \theta_W}{\sqrt{1 - \frac{\tan^2 \theta_W}{(\kappa_R^f)^2}}} \left[T_R^{3,f} - \frac{1}{\kappa_R^f} \tan^2 \theta_W Q^f \right], \end{aligned} \quad (2.5)$$

with $\kappa_R^{q,\ell}$ being the free parameters entering the interactions of eq. (2.1) (in which $\kappa_L^{q,\ell} = 0$) and where the electric and isospin charges are shown in table 2. Right-handed neutrino couplings could be easily added in our effective framework, following the minimal parameterization of ref. [51].

As a function of the vector boson mass, we show in figure 2 the total inclusive $pp \rightarrow W_{SSM}$ (solid fill) and $pp \rightarrow Z_{SSM}$ (hatch fill) production rates evaluated at NLO in QCD, assuming the inputs listed in section 2.2. We set the collision center-of-mass energy to (a) $\sqrt{s} = 13$ TeV and (b) 100 TeV, and use both the benchmark coupling normalizations given in eq. (2.4) (circle) as well as the much smaller choice (diamond)

$$\kappa_L^{q,\ell} = 0.01, \quad \kappa_R^q = 0 \quad \text{and} \quad \zeta_{Z_{SSM}}^f = 0.01. \quad (2.6)$$

We set as central factorization (μ_f) and renormalization (μ_r) scales half the sum of the transverse energies of all final-state particles,

$$\mu_f, \mu_r = \mu_0 = \frac{1}{2} \sum_{k \in \{\text{final states}\}} E_T^k \quad \text{with} \quad E_T^k = \sqrt{M_k^2 + p_T^k}. \quad (2.7)$$




Figure 2. Total NLO $pp \rightarrow W_{\text{SSM}}$ and $pp \rightarrow Z_{\text{SSM}}$ production cross section at a center-of-mass energy of $\sqrt{s} = 13$ TeV (a) and 100 TeV (b). The thickness of the curves corresponds to the residual scale uncertainty obtained by independently varying the central renormalization and factorization scales by a factor of two up and down.

The thickness of each curve in the main panel of figure 2 corresponds to the residual scale uncertainty as evaluated when varying μ_f and μ_r independently by a factor of two up and down with respect to the central scale μ_0 . We do not include uncertainties associated with parton distribution functions (PDF). At 13 TeV (100 TeV), the canonical SSM production rates for a boson mass lying in the [10 GeV, 5 TeV] ([10 GeV, 30 TeV]) range span approximately

$$\begin{aligned}
 W_{\text{SSM}} : & \quad 1.0^{+6.3\%}_{-6.8\%} - 55 \times 10^9 \quad {}^{+3.9\%}_{-21\%} \text{ fb} \quad \left(200 \times 10^{-3} \quad {}^{+3.7\%}_{-4.2\%} - 320 \times 10^9 \quad {}^{+16\%}_{-31\%} \text{ fb} \right), \\
 Z_{\text{SSM}} : & \quad 0.7^{+5.3\%}_{-5.7\%} - 25 \times 10^9 \quad {}^{+3.8\%}_{-21\%} \text{ fb} \quad \left(86 \times 10^{-3} \quad {}^{+3.2\%}_{-3.7\%} - 150 \times 10^9 \quad {}^{+18\%}_{-32\%} \text{ fb} \right),
 \end{aligned}
 \tag{2.8}$$

where the largest rates and residual scale uncertainties correspond to the smallest SSM boson masses. For the coupling scenario of eq. (2.6), the cross sections reduce precisely by a factor of 10^{-4} . As the same mass scales are probed, the uncertainties for both large and small SSM coupling scenarios are essentially the same. For electroweak (EW)- and TeV-scale boson masses, the residual scale uncertainties reaches the few-to-several percent level. However, unlike NNLO contributions, threshold resummation effects for $M_{W_{\text{SSM}}/Z_{\text{SSM}}}/\sqrt{s} \gtrsim 0.3$ greatly exceed the NLO uncertainty band [52].

In the lowest panel of the figures, we show NLO K -factor defined as the ratio

$$K^{\text{NLO}} \equiv \frac{\sigma^{\text{NLO}}(pp \rightarrow A + X)}{\sigma^{\text{LO}}(pp \rightarrow A + X)}, \quad (2.9)$$

for the standard scale choice of eq. (2.7). For $M_V > \mathcal{O}(10^2 - 10^3)$ GeV, we observe for both collider energies that NLO QCD corrections are, as expected [53], modest, with K^{NLO} remaining below ~ 1.4 . At smaller masses, NLO corrections are large and $K^{\text{NLO}} \gtrsim 2$ for $M_V \sim 10$ GeV. In the middle panel of the figures, we evaluate again eq. (2.9) but instead with a central scale choice of the partonic center-of-mass energy,

$$\mu_0 = \sqrt{\hat{s}}. \quad (2.10)$$

We observe the same qualitative dependence of K^{NLO} on the gauge boson mass M_V , which suggests that the K -factor is mostly independent of the scale choice. For $M_V = 10 - 50$ GeV, the large $\mathcal{O}(\alpha_s)$ correction is interpreted as the dominance of the $gq \rightarrow V'q'$ channel where the final-state quark p_T satisfies $p_T^q > \mu_f$. For the inclusive NLO V' production cross section, this is a LO-accurate contribution, and hence suffers from large scale uncertainties. The largeness of the $\mathcal{O}(\alpha_s)$ corrections and residual scale uncertainties highlight the importance of computing QCD corrections for processes sensitive to the deep low- x region of the gluon PDF, even for EW processes.

In the rest of this work, we focus on the canonical SSM parameterization, although our results can be easily generalized to any framework featuring extra gauge bosons. For arbitrary $\kappa_{L,R}^{q,\ell}$ parameters, the LO W_{SSM} partial decay widths to fermions are given by [49, 50, 54]

$$\begin{aligned} \Gamma(W_{\text{SSM}}^+ \rightarrow u_i \bar{d}'_j) &= N_c \left[\kappa_L^{q2} + \kappa_R^{q2} \right] |V_{ij}^{\text{CKM}}|^2 \frac{g^2 M_{W_{\text{SSM}}}}{48\pi}, \\ \Gamma(W_{\text{SSM}}^+ \rightarrow t \bar{b}) &= N_c \left[\kappa_L^{q2} + \kappa_R^{q2} \right] |V_{tb}^{\text{CKM}}|^2 \frac{g^2 M_{W_{\text{SSM}}}}{48\pi} \left(1 - r_t^{W_{\text{SSM}}} \right)^2 \left(1 + \frac{1}{2} r_t^{W_{\text{SSM}}} \right), \\ \Gamma(W_{\text{SSM}}^+ \rightarrow \ell^+ \nu_\ell) &= \left[\kappa_L^{\ell2} + \kappa_R^{\ell2} \right] \frac{g^2 M_{W_{\text{SSM}}}}{48\pi}, \end{aligned} \quad (2.11)$$

with $r_i^{W_{\text{SSM}}} = m_i^2/M_{W_{\text{SSM}}}^2$. Subsequently, the W_{SSM} total width reads, after summing over all final-state flavors,

$$\Gamma_{W_{\text{SSM}}} = \Gamma(W_{\text{SSM}} \rightarrow u \bar{d}) + \Gamma(W_{\text{SSM}} \rightarrow c \bar{s}) + \Gamma(W_{\text{SSM}} \rightarrow t \bar{b}) + 3\Gamma(W_{\text{SSM}} \rightarrow \ell \nu_\ell). \quad (2.12)$$

In the canonical SSM where the overall W_{SSM} coupling strengths are fixed as in eq. (2.4), the W_{SSM} branching fraction to a single lepton species is

$$\text{BR}(W_{\text{SSM}} \rightarrow \ell \nu_\ell) \approx \frac{g^2 M_{W_{\text{SSM}}}/48\pi}{g^2 M_{W_{\text{SSM}}}(N_c + 1)/16\pi} = \frac{1}{3(N_c + 1)} \approx 8.3\%, \quad (2.13)$$

where the approximation holds in the limit where the W_{SSM} boson mass $M_{W_{\text{SSM}}}$ is much larger than the top-quark mass m_t and where the CKM matrix is assumed to be an identity matrix. In the above expression, the factor of three corresponds to three generations with

Mass	30 GeV	300 GeV	500 GeV	3 TeV	5 TeV
$\Gamma(W_{\text{SSM}})$	0.760 GeV	8.92 GeV	16.1 GeV	101 GeV	169 GeV
$\Gamma(Z_{\text{SSM}})$	0.802 GeV	8.02 GeV	14.3 GeV	89.6 GeV	149 GeV

Table 3. LO W_{SSM} and Z_{SSM} total widths for representative $M_{W_{\text{SSM}}}$ and $M_{Z_{\text{SSM}}}$ mass values.

universal couplings, and $(N_c + 1)$ to the respective triplet and singlet color representations of quarks and leptons. The small branching fraction shows that W_{SSM} searches relying on leptonic final-state signatures may lack sensitivity in the high-mass region.

For arbitrary $\zeta_{Z_{\text{SSM}}}^f$ values, the LO Z_{SSM} partial widths to fermion-antifermion pairs $f\bar{f}$ are universally given by

$$\Gamma(Z_{\text{SSM}} \rightarrow f\bar{f}) = N_c^f \frac{\zeta_{Z_{\text{SSM}}}^{f2} g}{12\pi} M_{Z_{\text{SSM}}} \sqrt{1 - 4r_f^{Z_{\text{SSM}}}} \left[g_A^{f2} \sqrt{1 - 4r_f^{Z_{\text{SSM}}}} + g_V^{f2} (1 + 2r_f^{Z_{\text{SSM}}}) \right], \quad (2.14)$$

with $r_f^{Z_{\text{SSM}}} = m_f^2/M_{Z_{\text{SSM}}}^2$, N_c^f being respectively equal to 1 and to 3 for leptonic and quark final states, and where the vector and axial-vector couplings g_V^f and g_A^f are defined in eq. (2.3). The branching ratio into a specific leptonic final state in the canonical SSM and in the heavy Z_{SSM} limit is about 4%.

We evaluate in table 3 the total widths of canonical W_{SSM} and Z_{SSM} bosons for representative masses, using the EW input parameter values shown below in eq. (2.16). In such a setup, the bosons are always narrow, so that they could be discovered by several LHC searches for heavy resonances.

High-mass dijet resonance search results have constrained, at the 95% confidence level (CL), charged SSM boson masses to be above 2.6 TeV after analyzing CMS and ATLAS collision data at a center-of-mass energy of 13 TeV [55, 56], whereas low-mass dijet resonance searches are capable of excluding leptophobic Z_{SSM} with a mass in the [350, 500] GeV mass window for $\zeta_{Z_{\text{SSM}}}^q > 0.25 - 0.26$ [57]. Extra gauge boson searches in the leptonic channels currently constrain the neutral canonical Z_{SSM} boson to have a mass greater than 4.05 TeV (in the dileptonic mode) [2, 3] and the charged W_{SSM} boson to be heavier than 4.74 TeV (in the single leptonic mode) [4–7]. In terms of couplings, $W' \rightarrow \mu + \cancel{E}_T$ searches at 13 TeV imply W' couplings to fermions must obey [7]

$$\kappa_L^{q,\ell} = \frac{g'}{g_{\text{SM}}} \lesssim 2.6 \times 10^{-2} \quad \text{for} \quad \kappa_R^q = 0 \quad \text{and} \quad M_{W'} = 300 \text{ GeV}. \quad (2.15)$$

2.2 Computational setup

For concreteness, we consider as a benchmark scenario a SSM model with five flavors of massless quarks and a diagonal CKM matrix V^{CKM} with unit entries. We fix the EW inputs as in the 2014 Particle Data Group review [58],

$$\alpha^{\overline{\text{MS}}}(M_Z) = \frac{1}{127.940}, \quad M_Z = 91.1876 \text{ GeV} \quad \text{and} \quad \sin_{\overline{\text{MS}}}^2(\theta_W; M_Z) = 0.23126. \quad (2.16)$$

Our phenomenological study relies on automated NLO predictions matched to NNLL jet veto resummation as computed using the MADGRAPH5_aMC@NLO-SCET framework [18, 36]. More precisely, within MG5aMC (v2.5.1), one-loop virtual contributions are numerically evaluated by the MADLOOP package [59] and combined with the real contributions using the Frixione-Kunszt-Signer (FKS) subtraction method [60] as implemented in MADFKS [61]. For a jet veto of p_T^{Veto} and a hard process scale Q , logarithms of the form $\alpha_s^k(p_T^{\text{Veto}}) \log^l(Q/p_T^{\text{Veto}})$ with $l \leq 2k$ are resummed up to the NNLL following the procedure detailed in section 3.1. To generate the necessary UFO model library [47], we design a model file based on the above Lagrangians for the FEYNRULES program [44] (v2.3.10) that is jointly used with NLOCT [45] and FEYNARTS [46] (v3.8) for the computation of the ultraviolet and R_2 counterterms required for numerical one-loop calculations. Associated UFO files are available publicly from the FEYNRULES model database [48]. Hard scattering events are showered and hadronized using the PYTHIA 8 (PY8) infrastructure [62] (v8.212) and passed to MADANALYSIS 5 [63] (v1.4) for particle-level clustering using the FASTJET library [64] (v3.20) and its implementation of the anti- k_T algorithm [65].

Our calculations rely on PDFs and the evaluation of the strong coupling constant $\alpha_s(\mu_r)$ extracted using the LHAPDF 6 libraries [66] (v6.1.6). We employ the NNPDF 3.0 NLO PDF sets for LO and NLO calculations, and the NNLO set for NLO-NNLL calculations [67]. The factorization and renormalization scales are dynamically set according to eq. (2.7). Following ref. [7], underlying events are modeled by making use of the PY8 CUETP8M1 tune, also known as the “Monash*” tune [68].

3 Jet veto resummation

3.1 Jet veto resummation at next-to-next-to-leading logarithmic accuracy with next-to-leading order matching

Historically, the first higher order jet veto resummations were carried out in refs. [22–24]. In particular, within the SCET framework, jet veto resummation was developed in parallel in refs. [10, 14, 25, 26]. To carry out our NNLL jet veto resummation with fixed order NLO matching, we employ the resummation formalism developed in refs. [10, 14] and implemented into MG5aMC [18]. Within SCET [37–39, 69], jet veto resummation for the production of a color-singlet, n -body final-state system X , i.e.,

$$a b \rightarrow X \quad \text{with} \quad a, b \in \{q, \bar{q}, g\}, \tag{3.1}$$

follows from the existence of the resummed and refactorized fully differential cross section [10, 14],

$$\begin{aligned} \frac{d\sigma^{\text{N}^j\text{LL}}(p_T^{\text{Veto}})}{dy dQ^2 d\text{PS}_n} &= \sum_{a,b=g,q,\bar{q}} \left[\bar{B}_a(\xi_1, p_T^{\text{Veto}}) \bar{B}_b(\xi_2, p_T^{\text{Veto}}) + (1 \leftrightarrow 2) \right] \\ &\times E_I(Q^2, p_T^{\text{Veto}}, \mu_h, \mu, R) \mathcal{H}_{ab}(Q^2, \mu_h) \frac{d\hat{\sigma}_{ab}^B(Q^2, \mu)}{d\text{PS}_n}. \end{aligned} \tag{3.2}$$

Starting from the far right, $\hat{\sigma}_{ab}^B$ is the Born, parton-level scattering rate for the hard process given in eq. (3.1) that occurs at a scale Q and with a rapidity y . The so-called hard function

\mathcal{H}_{ab} contains the finite virtual corrections to $\hat{\sigma}_{ab}^B$, and, as non-vanishing loop diagrams factorize in the soft and collinear limits, is given by the power series

$$\mathcal{H}_{ab}(Q^2, \mu_h) = \sum_{k=0} \left(\frac{\alpha_s(\mu_h)}{4\pi} \right)^k \mathcal{H}_{ab}^{(k)}(Q^2, \mu_h) = 1 + \frac{\alpha_s(\mu_h)}{4\pi} \mathcal{H}_{ab}^{(1)}(Q^2, \mu_h) + \mathcal{O}(\alpha_s^2). \quad (3.3)$$

The $\mathcal{H}_{ab}^{(k)}$ coefficients possess logarithms of (Q/μ_h) , where μ_h is the scale at which $\mathcal{H}_{ab}^{(k)}$ is regulated, that can spoil the perturbative convergence of eq. (3.2) if $\mu_h \ll Q$. To avoid this, one sets $\mu_h \sim Q$. Details on the efficient evaluation of the $\mathcal{O}(\alpha_s)$ coefficient $\mathcal{H}_{ab}^{(1)}$ numerically can be found in refs. [18, 59]. The evolution of the hard process down to scales $\mu, p_T^{\text{Veto}} \ll \mu_h, Q$ is governed by the evolution factor E_I ,

$$E_I(Q^2, p_T^{\text{Veto}}, \mu_h, \mu, R) = U_I(Q^2, \mu_h, \mu) e^{-2F_I(p_T^{\text{Veto}}, \mu, R) \log \frac{Q}{p_T^{\text{Veto}}}} e^{2h_I(p_T^{\text{Veto}}, \mu)}. \quad (3.4)$$

The renormalization group evolution function U_I consists of exponentiated Sudakov form factors and anomalous dimensions. The first exponential in the above expression is the collinear anomaly that arises from the breaking of the scale invariance of hadron momenta at the one-loop level in SCET [70]. At the classical level, the proton momenta are given by $P_i = E_{P_i}(1, 0, 0, \pm 1)$ and the momentum fractions ξ_1 and ξ_2 carried by the partons a and b remain unchanged with the scaling $P_i \rightarrow \tilde{P}_i = \lambda P_i$. One indeed has

$$\xi_i \rightarrow \tilde{\xi}_i = \tilde{p}_a^0 / \tilde{P}_i^0 = \lambda p_a^0 / \lambda P_i^0 = \xi_i. \quad (3.5)$$

In particular, simultaneously scaling both proton momenta by λ_i and $\lambda_j = \lambda_i^{-1}$ leaves the hard scale Q unchanged,

$$Q \rightarrow \tilde{Q}^2 = (4\tilde{p}_a^0 \tilde{p}_b^0) = (4p_a^0 p_b^0) \lambda_i \lambda_i^{-1} = Q^2. \quad (3.6)$$

In the SCET context, while the former invariance of eq. (3.5) is broken, the latter one of eq. (3.6) remains intact. In the context of perturbative QCD, the collinear anomaly, which arises first at NNLL, can be understood as the interference between soft virtual corrections and collinear emissions [71]. The second exponential in eq. (3.4) is an auxiliary evolution function that connects the scale μ to the veto scale p_T^{Veto} . Whereas the indices a and b appearing in $\hat{\sigma}_{ab}^B$, \mathcal{H}_{ab} and in the beam functions \bar{B}_a and \bar{B}_b (below) denote specific incoming partons, e.g., $a = b = g$ or $a (b) = u (\bar{d})$, the index $I \in \{g, \bar{q}\}$ in the evolution factor E_I refers to the color representations associated with the $q\bar{q}$ or gg initial state. This emphasizes the fact that eq. (3.2) only holds for color-singlet processes.

Lastly, the beam function for a parton species a in a proton p with a transverse momentum $p_T^a < p_T^{\text{Veto}}$ and carrying a longitudinal momentum fraction $\xi_i = E_a/E_{P_i} = e^{\pm y} Q/\sqrt{s}$ into the hard process is given by

$$\bar{B}_a(\xi, p_T^{\text{Veto}}) = \sum_{c=g, q, \bar{q}} \int_{\xi}^1 \frac{dz}{z} \bar{I}_{ac}(z, p_T^{\text{Veto}}, \mu_f) f_{c/p}\left(\frac{\xi}{z}, \mu_f\right). \quad (3.7)$$

The function $f_{c/p}(x, \mu_f)$ denotes the usual transverse-momentum-integrated density of a parton species c in the proton p carrying a longitudinal momentum fraction $x = (\xi/z)$ and




Figure 3. Schematic description of the factorization theorem with jet veto resummation in the SCET framework.

evolved to a collinear factorization scale μ_f . The $c \rightarrow a$ splitting kernel \bar{I}_{ac} accounts for the low- p_T (i.e., $p_T < p_T^{\text{Veto}}$) collinear splittings of partons that emerge from $f_{c/p}(x, \mu_f)$ and connects the factorization scale μ_f to the veto scale p_T^{Veto} . For $\mu_f \sim p_T^{\text{Veto}}$, \bar{I} can be expanded in powers of α_s with coefficients consisting of the Altarelli-Parisi splitting functions. Moreover, in the $(p_T^{\text{Veto}}/Q) \rightarrow 0$ limit, emission recoils can be neglected and the partons a and b in eq. (3.1) remain parallel to their parent protons.

The factorization theorem of eq. (3.2) is illustrated in figure 3 and represents the likelihood of the process $pp \rightarrow X$ to occur when X is accompanied by an arbitrary number of QCD emissions possessing a transverse momentum $p_T < p_T^{\text{Veto}}$. It is derived in the $(p_T^{\text{Veto}}/Q) \rightarrow 0$ limit and hence is valid up to (p_T^{Veto}/Q) power corrections. For even moderate values of p_T^{Veto} , such contributions are a source of sizable theoretical uncertainties. These corrections, however, are precisely those that are well-described by fixed order calculations, which follows from the usual Collins Soper Sterman (CSS) Collinear Factorization Theorem [35]. The problem can thus be resolved by an appropriate matching procedure that entails removing the double-counted regions of phase space. The use of eq. (3.2) is necessary due to the breakdown of the CSS formalism in the presence of a jet veto: a veto prematurely terminates a summation over all collinear, but potentially high- p_T , emissions that is otherwise necessary for ensuring the cancellation of long-range Glauber gluon exchanges [72].

Matching fixed order and resummed expressions usually involves Taylor expanding the resummed expression $d\sigma^{\text{N}^j\text{LL}}(p_T^{\text{Veto}})$ in powers of α_s to the same accuracy of the fixed order result $d\sigma^{\text{N}^k\text{LO}}|_{p_T < p_T^{\text{Veto}}}$. This quantity is then subtracted from the sum of the fixed order and resummed calculations. For instance, NNLO matching would require an $\mathcal{O}(\alpha_s^2)$ expansion. In the SCET framework, the presence of the hard and evolution functions marginally complicates the procedure. Extracting these functions, one can rewrite eq. (3.2) as

$$\frac{d\sigma^{\text{N}^j\text{LL}}(p_T^{\text{Veto}})}{dy dQ^2 d\text{PS}_n} = \sum_{a,b=g,q,\bar{q}} E_I(Q^2, p_T^{\text{Veto}}, \mu_h, \mu, R) \times \mathcal{H}_{ab}(Q^2, \mu_h) \times \left\{ \left[\bar{B}_a(\xi_1, p_T^{\text{Veto}}) \bar{B}_b(\xi_2, p_T^{\text{Veto}}) + (1 \leftrightarrow 2) \right] \frac{d\hat{\sigma}_{ab}^B(Q^2, \mu)}{d\text{PS}_n} + \Delta\tilde{\sigma}_{ab} \right\}, \quad (3.8)$$

where the $\Delta\tilde{\sigma}_{ab}$ term stands for the (p_T^{Veto}/Q) power corrections with E_I and \mathcal{H}_{ab} factored out. At the NNLL accuracy, the beam functions correspond to the Altarelli-Parisi splitting kernels \bar{I} expanded to $\mathcal{O}(\alpha_s)$, which means that the bracketed quantity represents low- p_T QCD emissions off the Born process up to $\mathcal{O}(\alpha_s)$. Physically, this is equivalent to the NLO calculation once a selection on the transverse momentum of the radiated jet of $p_T^j < p_T^{\text{Veto}}$ is imposed. After subtracting the resummed-fixed order overlap, the matched differential jet veto cross section at the NLO+NNLL accuracy is given, for the generic process introduced in eq. (3.1), by

$$\begin{aligned} \frac{d\sigma^{\text{NLO+NNLL}}(p_T^{\text{Veto}})}{d\text{PS}_n} &= \sum_{a,b=g,q,\bar{q}} E_I(Q^2, p_T^{\text{Veto}}, \mu_h, \mu, R) \times \left(1 + \frac{\alpha_s(\mu_h)}{4\pi} \mathcal{H}_{ab}^{(1)}(Q^2, \mu_h) \right) \\ &\times \left[\frac{d\sigma_{ab}^{\text{NLO}}}{d\text{PS}_{(n+1)}} \Big|_{p_T < p_T^{\text{Veto}}} - \frac{\alpha_s(\mu)}{4\pi} \left(\mathcal{H}_{ab}^{(1)}(Q^2, \mu) + E_I^{(1)}(Q^2, p_T^{\text{Veto}}, \mu) \right) \frac{d\sigma_{ab}^{\text{LO}}}{d\text{PS}_n} \right]. \end{aligned} \quad (3.9)$$

Numerically, the matched result is evaluated over an $(n + 1)$ -body phase space domain despite the process in eq. (3.1) being an n -body process. The extra emission is however soft by construction, so that each $(n + 1)$ -body phase space point is mapped to an n -body configuration following the FKS prescription [60]. With the exception of $d\sigma^{\text{NLO}}$, all terms are then evaluated within the n -body subspace for the Born process. The relevant analytic expressions for the ingredients contributing to the matched cross section of eq. (3.9) can be found in ref. [18] and the references therein. The NNLL resummation describes the likelihood of process eq. (3.1) being accompanied by up to two soft emissions, implying some overlap with the NNLO fixed order calculation. It is therefore more appropriate to use NNLO PDFs when performing NLO+NNLL computations as oppose to NLO PDFs, which are needed for NLO computations.

3.2 Non-perturbative corrections to cross sections with jet vetoes

A consequence of the collinear anomaly in eq. (3.2) is the emergence of logarithmically enhanced non-perturbative corrections that, following ref. [18], are expected to behave as

$$\frac{\delta\sigma^{\text{Non-Pert.}}}{\sigma^{\text{Born}}} \sim \frac{\Lambda_{\text{Non-Pert.}}}{p_T^{\text{Veto}}} \log \left(\frac{Q}{p_T^{\text{Veto}}} \right), \quad (3.10)$$

where the energy scale $\Lambda_{\text{Non-Pert.}} \sim \mathcal{O}(1 - 2) \text{ GeV}$ is the scale at which QCD becomes strongly coupled. Such uncertainties are distinct from non-perturbative corrections to jet observables [74], e.g., shifts in p_T of the hardest jet from out-of-jet emissions of hadrons. A study of this second class of corrections in the context of jet vetoes is beyond the scope of this report. However, some of these effects are included due to our use of a modern parton shower in our NLO+PS-accurate event simulations [74]. For $Q \gg p_T^{\text{Veto}}$, non-perturbative contributions can be sizable. To investigate the impact of these terms when employing jet vetoes in searches for new color-singlet states at hadron colliders, we present the relative magnitude of the non-perturbative contributions of eq. (3.10) as a function of Q and for representative p_T^{Veto} values in figure 4. We choose $\Lambda_{\text{Non-Pert.}} = \Lambda_{\text{Non-Pert.}}^{\text{Default}} = 1 \text{ GeV}$ as




Figure 4. Non-perturbative corrections to jet veto rates arising from the collinear anomaly [18, 73] in jet-veto resummed cross sections for low (a) and high (b) ranges of the hard process scale Q , and for representative p_T^{Veto} values. The thickness of the bands reflects the variation of the non-perturbative scale $\Lambda_{\text{Non-Pert.}}$ as shown in eq. (3.11).

the central value for the non-perturbative scale, and we include $\Lambda_{\text{Non-Pert.}}$ variation bands obtained by spanning

$$0.5 \times \Lambda_{\text{Non-Pert.}}^{\text{Default}} < \Lambda_{\text{Non-Pert.}} < 2 \times \Lambda_{\text{Non-Pert.}}^{\text{Default}}. \quad (3.11)$$

As the $\Lambda_{\text{Non-Pert.}}$ dependence in eq. (3.10) is linear, these arbitrarily chosen limits induce precisely a variation of a factor of two up and down around the central value extracted from eq. (3.10).

For low p_T^{Veto} scales of 10, 30 and 50 GeV, the relative size of the non-perturbative contribution (with respect to the Born process) respectively reaches $\delta\sigma^{\text{Non-Pert.}}/\sigma^{\text{Born}}$ values of about 20%, 4% and 2% for a hard scale of $Q = 100$ GeV. For a larger choice of $Q = 1$ TeV, the non-perturbative effects are expected to grow to 50%, 15% and 6%. At an even larger scale of $Q = 10$ TeV, the uncertainty originating from a choice of $p_T^{\text{Veto}} = 30$ GeV reaches the 20% level, whereas it drops to $\lesssim 5\%$ for $p_T^{\text{Veto}} \gtrsim 100$ GeV. Those results may suggest that the linear dependence on the ratio $(\Lambda_{\text{Non-Pert.}}/p_T^{\text{Veto}})$ in eq. (3.10) spoils perturbative predictability for overly aggressive p_T^{Veto} choices when probing mass scales well above the EW scale. On the other hand, equally aggressive p_T^{Veto} choices for EW-scale processes give rise to non-perturbative corrections that are comparable or within the current perturbative and experimental uncertainties [10, 20, 34].

For a potential next-generation hadron collider with a center-of-mass energy well above 13 TeV, and hence sensitivity to comparably larger hard scales Q , the necessity for choosing p_T^{Veto} at or above the EW scale to avoid large non-perturbative corrections raises the question of whether or not jet vetoes are practical for high-mass resonance searches. Standard Model processes, like $t\bar{t}$ production, dominantly occur near threshold, so associated final state momenta scale like the EW scale, and thereby evade such vetoes. It may be more advantageous to veto according to a different metric, such as jet mass. However, it may also be possible that further investigations into the non-perturbative corrections induced by the




Figure 5. Estimated uncertainty on jet-veto resummed predictions for different choices of the jet radius parameter R for pp collisions at a center-of-mass energy of 13 TeV (a) and 100 TeV (b). The results are presented as a function of the W_{SSM} boson mass and relatively to the NLO total rate $\sigma_{\text{Tot}}^{\text{NLO}}$. The uncertainties associated with the later are indicated by a black band.

collinear anomaly reveals a milder sensitivity to p_T^{Veto} than in eq. (3.10). In particular, one may find for perturbative choices of p_T^{Veto} , e.g., $p_T^{\text{Veto}} = 30 - 40$ GeV, where $\alpha_s(p_T^{\text{Veto}}) \ll 1$, that the non-perturbative contributions turn out to be negligible when probing multi-TeV hard process scales.

3.3 Scale uncertainties in resummed jet veto rates from varying jet definitions

The scale dependence of jet veto calculations on the jet definition is sizable but also intuitive: for a given hadron collision, a geometrically larger jet will contain more objects and hence will be associated with a larger mass scale. This can lead to a larger jet momentum implying that the corresponding event is more likely to be vetoed. Furthermore, the lowest order at which the p_T spectrum of any color-singlet system, which is necessary for calculating jet vetoes in perturbative QCD, is qualitatively accurate is at NLO for the inclusive process matched to $\text{LL}(k_T)$ resummation. This is also the formal accuracy of NLO+PS calculations used with present day general-purpose event generators. Similarly, for the veto-resummed calculation, an explicit dependence on the jet radius parameter R of the k_T -style jet algorithms appears first at the two-loop order, i.e., at the NNLL level [11, 14]. Therefore, predictions provided at NLO+NNLL(Veto) accuracy embeds the lowest order scale dependence on the choice of R .

To explore the jet veto scale uncertainty associated with different jet definitions, we consider the benchmark process

$$pp \rightarrow W_{\text{SSM}} \tag{3.12}$$

and focus on two collider energies of $\sqrt{s} = 13$ and 100 TeV. We restrict ourselves to only investigating the dependence on varying R and p_T^{Veto} as the veto resummation is identical for all k_T -style algorithms at NNLL. In our choice of representative R values, we are limited by two factors. First, the factorization theorem of eq. (3.2) assumes a jet radius R

satisfying [10]

$$\frac{p_T^{\text{Veto}}}{Q} \ll R \ll \log \frac{Q}{p_T^{\text{Veto}}}, \quad (3.13)$$

which indicates that R and p_T^{Veto} must obey the relationship

$$p_T^{\text{Veto}} \ll Q \times e^{-R} \approx Q \left(1 - R + \frac{R^2}{2} \right). \quad (3.14)$$

For respectively small, medium and large radius with $R = 0.1, 0.4$ and 1 , this translates to p_T^{Veto} scales much smaller than $0.9Q, 0.7Q$ and $0.5Q$. For larger p_T^{Veto} scales, matching to the fixed order calculation is necessary due to a breakdown of the factorization theorem of eq. (3.2), as derived in refs. [10, 14]. The second limitation stems from the logarithmic dependence on R of the evolution function E_I introduced in eq. (3.2). For very small jet radii, these logarithmic terms are large and need to be resummed [26, 75]. The study of the impact of these resummed small- R logarithms is beyond the scope of the present analysis and we refer to ref. [20] for more information. For large R , the expressions for the anomalous dimensions in eq. (3.4) break down [18].

In figure 5, we present, as a function of the W_{SSM} boson mass, the veto efficiency $\varepsilon^{\text{NLO+NNLL(Veto)}}$ associated with the process of eq. (3.12),

$$\varepsilon^{\text{NLO+NNLL(Veto)}}(p_T^{\text{Veto}}) \equiv \frac{\sigma^{\text{NLO+NNLL(Veto)}}(pp \rightarrow W_{\text{SSM}} \rightarrow \ell\nu_\ell; p_T^{\text{Veto}})}{\sigma_{\text{Tot.}}^{\text{NLO}}(pp \rightarrow W_{\text{SSM}} \rightarrow \ell\nu_\ell)}, \quad (3.15)$$

for representative jet radii of $R = 0.1, 0.4$ and 1 , with $p_T^{\text{Veto}} = 30$ (100) GeV, and a collider energy of 13 (100) TeV. Shaded bands correspond to the scale uncertainty; PDF uncertainties are omitted. At both colliders, we observe systematically smaller efficiencies for larger R values, in agreement with the argument above. For increasing W_{SSM} mass, we observe a monotonically decreasing veto efficiency, which follows from logarithmically-enhanced soft-gluon emissions that grow as $\alpha_s(p_T^j) \log(Q^2/p_T^{j,2})$ for $Q \sim M_{W_{\text{SSM}}}$. This tendency for higher mass color singlet processes to radiate more soft gluons is in addition the basic argument motivating threshold and recoil resummations.

As a function of R , the associated scale uncertainty shrinks (grows) with increasing (decreasing) jet radius due to the increasing (decreasing) inclusiveness of the observable $\varepsilon(p_T^{\text{Veto}})$. For $p_T^{\text{Veto}} = 30$ GeV at 13 TeV, the uncertainties on the veto efficiency are of $\delta\varepsilon \sim 10\%, 5\%$ and 1% for $R = 0.1, 0.4$ and 1 respectively. For $p_T^{\text{Veto}} = 100$ GeV at 100 TeV, they correspondingly drop to $\delta\varepsilon \sim 5\%, 1\%$ and 1% . The fleetingly small uncertainties associated with the $R = 1$ jet case are due to the R dependence in the evolution operator E_I being largely logarithmic, up to neglected power corrections. They are therefore minimized in the $R \rightarrow 1$ limit.

In figure 6, we show the scale dependence of the veto efficiency on the veto scale p_T^{Veto} for representative W_{SSM} masses and radius $R = 1$ at (a) $\sqrt{s} = 13$ and (b) 100 TeV. The results are consistent with the findings of figure 5: for a fixed p_T^{Veto} , the production of a heavier W_{SSM} bosons leads to events that are relatively enriched with high- p_T jets, which subsequently reduce the veto efficiency. As a function of collider energies, increasing




Figure 6. Dependence of the resummed jet-veto efficiencies on the veto scale p_T^{Veto} for the $pp \rightarrow W_{\text{SSM}} \rightarrow \mu\nu$ process at a collider energy of 13 TeV (a) and 100 TeV (b). Basic acceptance selections are included, and we consider a representative sample of W_{SSM} boson masses.

\sqrt{s} opens phase space for more jet activity, which again implies a smaller veto efficiency. Conversely, for a fixed W_{SSM} boson mass, increasing the veto scale increases the veto efficiency since events are allowed to feature harder jets. For increasing p_T^{Veto} , ε converges to 1 and the matched-resummed result converges to the fixed order result, as one would expect. However, as the NLO+NNLL result has been computed with NNLO PDFs whereas the NLO result with NLO PDFs, there exists a slight mismatch between the different central values that is within the (not shown) PDF uncertainties.

4 Signal and background process modeling with jet vetoes

Searches for $W' \rightarrow e/\mu + \cancel{E}_T$ are inundated with SM Drell-Yan continua and leptonic decays of top quarks. In this section, we describe our procedure for modeling both the signal and background processes with jet vetoes. For all processes, we use the computational setup described at the end of section 2.

4.1 W' production and decay

The benchmark BSM collider signature that we will ultimately simulate consists of the charged current process

$$pp \rightarrow W' \rightarrow \mu \nu, \tag{4.1}$$

where we consider a final-state muon for the sake of an example. We use the SSM coupling normalizations of eq. (2.4) and reinterpret our results for smaller coupling strengths introduced via a multiplicative scaling factor. We neglect any interference with the SM W boson due to a severe model dependence that prevents us from including these effects in a generic way. While necessary for any SSM-like scenario with a boson mass $M_{W'}$ of the order of the SM W boson mass, little or no such interference is present for right-handed W_R bosons in left-right symmetric models or for W' bosons that are odd under some discrete symmetry with respect to the SM W .

$M_{W'}$ [GeV]	$\sigma_{\text{Tot.}}^{\text{NLO}}$ [fb]	$\sigma_{(\text{Veto})}^{\text{NLO+PS}}$ [fb]	$\sigma_{(\text{Veto})}^{\text{NLO+NNLL}}$ [fb]	$K_{\text{NLO+PS}}^{\text{NLO+NNLL(Veto)}}$
30	$262_{-25\%}^{+16\%} \times 10^6$	$256_{-14\%}^{+9.5\%} \times 10^6$	$296_{-8.2\%}^{+4.5\%} \times 10^6$	1.16
50	$68.9_{-17\%}^{+9.4\%} \times 10^6$	$65.6_{-8.7\%}^{+5.7\%} \times 10^6$	$72.8_{-6.4\%}^{+3.4\%} \times 10^6$	1.11
300	$289_{-2.8\%}^{+2.1\%} \times 10^3$	$213_{-1.0\%}^{+2.4\%} \times 10^3$	$227_{-0.5\%}^{+0.8\%} \times 10^3$	1.07
500	$47.8_{-1.0\%}^{+1.4\%} \times 10^3$	$31.7_{-2.3\%}^{+1.8\%} \times 10^3$	$33.6_{-1.0\%}^{+1.1\%} \times 10^3$	1.06
1000	$3.58_{-1.5\%}^{+1.7\%} \times 10^3$	$2.04_{-1.5\%}^{+0.7\%} \times 10^3$	$2.19_{-2.2\%}^{+2.0\%} \times 10^3$	1.07
3000	$15.4_{-2.3\%}^{+1.2\%}$	$7.73_{-1.7\%}^{+0.1\%}$	$8.06_{-3.4\%}^{+0.6\%}$	1.04
5000	$446_{-1.7\%}^{+1.3\%} \times 10^{-3}$	$263_{-0.8\%}^{<0.1\%} \times 10^{-3}$	$258_{-1.8\%}^{+0.7\%} \times 10^{-3}$	0.98

Table 4. Cross sections [fb] for $pp \rightarrow W' \rightarrow \mu\nu_\mu$ at various accuracies with residual scale uncertainties [%] (no PDF uncertainties), at the 13 TeV LHC. The results are shown for representative W' boson masses and either without (second column) or with (third and fourth columns) a jet veto (for $p_T^{\text{Veto}} = 40$ GeV, $R = 1$). The K -factor defined in eq. (4.2) is also indicated (last column).

We first generate events at the NLO+PS accuracy for the 13 TeV LHC. At the analysis level, we impose a jet veto by rejecting events with $R = 1$ jets possessing $p_T^j > p_T^{\text{Veto}} = 40$ GeV. For several representative W' masses, table 4 summarizes the total inclusive cross section obtained at NLO ($\sigma_{\text{Tot.}}^{\text{NLO}}$, second column), as well as NLO+PS after applying the above jet veto selection ($\sigma_{(\text{Veto})}^{\text{NLO+PS}}$, third column). The resummed result $\sigma_{(\text{Veto})}^{\text{NLO+NNLL}}$ is given in the fourth column of the table and will be used for normalizing the generated NLO+PS events to the NLO+NNLL(Veto) cross section. We report residual scale uncertainties [%]; PDF uncertainties are omitted. To quantify the impact of this normalization, we define an appropriate K -factor as the ratio of the resummed rate to the NLO+PS rate once a jet veto event selection is applied,

$$K_{\text{NLO+PS}}^{\text{NLO+NNLL(Veto)}}(p_T^{\text{Veto}}) \equiv \frac{\sigma_{(\text{Veto})}^{\text{NLO+NNLL(Veto)}}(pp \rightarrow W' + X; p_T^{\text{Veto}})}{\sigma_{(\text{Veto})}^{\text{NLO+PS}}(pp \rightarrow W' + X; p_T^{\text{Veto}})}. \quad (4.2)$$

We give, in the last column of table 4, the corresponding values for this K -factor. For light W' bosons, the K -factors are of the order $K \gtrsim 1.1$ reduce to $K \sim 1.05$ for $M_{W'} > 100$ GeV, and drop below this for $M_{W'} > 1$ TeV. In most cases, the PS and resummed results agree within one or two widths of their scale uncertainty bands. Not shown PDF uncertainties contribute to an additional $\mathcal{O}(1-2)\%$ error. Below 5 TeV, the K -factors are greater than unity, indicating that the logarithmic corrections in the resummed calculation are positive-definite. Our K -factors are in agreement with the findings of ref. [17] for EW-scale masses and suggest that the PS and NNLL result converge at much larger mass scales. NNLO corrections to the NLO result are known to be comparable in size and negative, indicating that the NNLO+NNLL(Veto) result is in agreement with both the

NLO+PS and NLO+NNLL(Veto) calculations [17]. As the resummed corrections are essentially independent of the hard process, we expect this behavior to broadly extend to other color-singlet processes. Hence, within their given uncertainties, both the NLO+PS and NLO+NNLL(veto) calculations give accurate and consistent predictions. Consequently, jet vetoes applied to color-singlet BSM processes can be reliably modeled at the NLO+PS level. This is a main finding of our investigation.

We now briefly comment on whether normalization by eq. (4.2) is justified at a differential level. In short, particle kinematics for color-singlet processes in resummed calculations, which possess Born-like kinematics, and in NLO+PS calculations, which include recoil from soft and hard radiation, are largely the same after applying a jet veto. This follows from factorization in unbroken gauge theories: amplitudes containing QCD radiations in the soft/collinear limit factorize into a product of universal form factors and the (color-connected) Born amplitude. As a consequence, in this limit, $\mathcal{O}(\alpha_s)$ corrections to differential distributions for inclusive DY processes reduce to a multiplicative factor applied to the Born cross section. Furthermore, this holds analytically for arbitrary W'/Z' couplings and masses [76]. As the jet veto by definition removes hard QCD radiations and parton showers are based on collinearly factorized emissions, the kinematics of the two results should therefore exhibit differences only of the order of (p_T^{Veto}/Q) , which we assume to be vanishingly small for the validity of the jet veto factorization theorem in eq. (3.2).

To verify that this holds, we focus on the process in eq. (4.1) and present, in figure 7, the (a) p_T and (b) pseudorapidity η distribution of the muon at 13 TeV. We show results, for representative W' masses, both at LO (solid) and NLO+PS accuracy with a jet veto of $p_T^{\text{Veto}} = 30$ GeV (dash). At LO, the veto has no impact as no jets are present. In the lower panel of the figure, we depict the differential NLO+PS K -factor for each observable $\hat{\mathcal{O}}$,

$$K_{\hat{\mathcal{O}}}^{\text{NLO+PS}}(p_T^{\text{Veto}}) \equiv \frac{d\sigma^{\text{NLO+PS}}(pp \rightarrow W' + X; p_T^{\text{Veto}})/d\hat{\mathcal{O}}}{d\sigma^{\text{LO}}(pp \rightarrow W' + X)/d\hat{\mathcal{O}}}. \quad (4.3)$$

For both distributions, we observe that the bin-by-bin ratios of the LO and NLO+PS distributions are largely flat when away from resonant regions. This indicates that the NLO+PS result with a jet veto is dominated by soft gluon radiation, and therefore that the NLO+PS+ p_T^{Veto} kinematics approximate well the jet veto-resummed kinematics.

4.2 SM Drell-Yan continua

To model the SM charged and neutral current DY continua, we simulate at NLO+PS accuracy the processes

$$pp \rightarrow W^{(*)} \rightarrow \mu\nu \quad \text{and} \quad pp \rightarrow \gamma^*/Z^{(*)} \rightarrow \mu^+\mu^-. \quad (4.4)$$

For the neutral current channel, we impose a generator-level regulator on the dilepton invariant mass $M_{\ell\ell} > 10$ GeV. Due to their color-singlet nature, we treat the DY background much like the signal process, normalizing the cross sections after including a jet veto by a K -factor such as the one defined in eq. (4.2). For the neutral current background, the mass scale is naturally given by the invariant mass of the dilepton system, so that for each $M_{W'}$




Figure 7. Muon transverse-momentum (a) and pseudorapidity (b) distributions for the $pp \rightarrow W' \rightarrow \mu\nu_\mu$ process at a center-of-mass energy of 13 TeV. We show results at the LO accuracy (solid) and at the NLO+PS accuracy when a jet veto such that $p_T^{\text{Veto}} = 30$ GeV is included (dash). We have selected a few representative W' boson masses.

mass hypothesis, we derive the K -factor by additionally imposing the selection $M_{\ell\ell} > M_{W'}$. In the charged current case, constructing the $M_{\ell\nu}$ invariant mass is more subtle due to the (typical) inability to reconstruct the longitudinal momentum of final-state neutrinos at the PS level. However, as we discuss in section 6.1, we adopt as a discriminating variable sensitive to the W' mass scale the transverse mass m_T of the lepton- \cancel{E}_T system. Therefore, for each $M_{W'}$ mass hypothesis, we determine the K -factor after imposing the selection $m_T > M_{W'}$. Technically, this selection can be implemented in MG5aMC by identifying neutrinos as *charged* leptons in the `SubProcesses/cuts.f` and `SubProcesses/setcuts.f` files, and by replacing the $M_{\ell\ell}$ observable by an implementation of the transverse mass m_T in `SubProcesses/cuts.f`. The relevant selection parameter is thus `m11`, as in the neutral current case.

In figure 8 we present, as a function of the dilepton mass scale $M_{\ell\ell}$ and m_T for the neutral and charged current cases respectively, the veto efficiency for the DY processes given in eq. (4.4) for (a) a veto scale of $p_T^{\text{Veto}} = 30$ GeV at 13 TeV, and (b) $p_T^{\text{Veto}} = 100$ GeV at 100 TeV. As anticipated, the impact of the veto becomes more severe for increasing mass scales, just like the W' case treated in section 4.1. For both collider and veto setups, we find that the jet veto efficiencies are independent of the processes and span roughly

$$\begin{aligned}
 13 \text{ TeV} : \quad & \varepsilon^{\text{NLO+NNLL(Veto)}}(p_T^{\text{Veto}} = 30 \text{ GeV}) = 90 - 30\% \text{ for } M_{\ell X} \in [0.050, 5] \text{ TeV}, \\
 100 \text{ TeV} : \quad & \varepsilon^{\text{NLO+NNLL(Veto)}}(p_T^{\text{Veto}} = 100 \text{ GeV}) = 80 - 30\% \text{ for } M_{\ell X} \in [0.3, 30] \text{ TeV},
 \end{aligned}
 \tag{4.5}$$

with a residual scale uncertainty of about $\pm 1 - 5\%$.




Figure 8. Dependence of the resummed jet-veto efficiencies on the veto scale p_T^{Veto} for the neutral current and charged current DY processes at a collider energy of 13 TeV (a) and 100 TeV (b). Basic acceptance selections on the gauge boson decay product are included, and we also show the fixed order results.

In tables 5 and 6, we report, for several representative mass scales, the inclusive cross sections for the charged current and neutral current DY channels, respectively. The predictions are given at NLO ($\sigma^{\text{NLO}}_{\text{Tot}}$, second column), NLO+PS after applying a jet veto with $p_T^{\text{Veto}} = 40$ GeV ($\sigma^{\text{NLO+PS}}$, third column), and after resumming the jet veto effects ($\sigma^{\text{NLO+NNLL(Veto)}}$, fourth column). The veto K -factor defined as in eq. (4.2) is shown in the sixth column. Overall, we find a good agreement between the parton showered and resummed predictions given their few-percent-level uncertainties. For both channels, the K -factors are found to span approximately the 1.0 – 1.1 for mass scales above 30 GeV. Despite the three different scale choices, i.e., $M_{W'}$, $M_{\ell\ell}$, and m_T , we observe the K -factors for the signal and background processes to be very comparable in size and direction.

4.3 Top quark background

The top quark background for $W' \rightarrow \ell\nu_\ell$ searches contains both a top-antitop pair and single top component,

$$pp \rightarrow t\bar{t} \rightarrow \ell^\pm + \cancel{E}_T + X, \quad pp \rightarrow t j \rightarrow \ell^\pm + \cancel{E}_T + X \quad \text{and} \quad pp \rightarrow t W^* \rightarrow \ell^\pm + \cancel{E}_T + X, \quad (4.6)$$

where one or all top quarks decay leptonically for the first two processes, and where either the top quark or the W -boson (or both) proceeds via a leptonic decay in associated tW production. In the five-flavor scheme, the s -channel $t\bar{b}$ production mode is included in the tj process definition. We ignore additional channels, such as associated $t\bar{t}W/Z/\gamma^*$ production, as they are both coupling suppressed with respect to the above processes and possess similar kinematics.

We simulate inclusive $t\bar{t}$ and tj production at NLO+PS accuracy. For medium and high W' boson mass, we impose a generator-level selections on the top quark transverse momentum. For the tW channel, we simulate the $pp \rightarrow t\ell\nu_\ell$ process at LO+PS accuracy.

m_T [GeV]	$\sigma_{\text{Tot.}}^{\text{NLO}}$ [fb]	$\sigma_{(\text{Veto})}^{\text{NLO+PS}}$ [fb]	$\sigma_{(\text{Veto})}^{\text{NLO+NNLL}}$ [fb]	$K_{\text{NLO+PS}}^{\text{NLO+NNLL(Veto)}}$
0	$18.4^{+6.7\%}_{-12\%} \times 10^6$	$16.5^{+4.1\%}_{-4.7\%} \times 10^6$	$18.4^{+2.7\%}_{-5.0\%} \times 10^6$	1.12
30	$16.3^{+6.3\%}_{-11\%} \times 10^6$	$14.7^{+4.2\%}_{-4.0\%} \times 10^6$	$16.1^{+2.6\%}_{-4.8\%} \times 10^6$	1.10
50	$12.8^{+5.3\%}_{-10\%} \times 10^6$	$11.6^{+4.6\%}_{-3.7\%} \times 10^6$	$12.7^{+2.6\%}_{-4.8\%} \times 10^6$	1.09
100	$71.2^{+3.7\%}_{-6.2\%} \times 10^3$	$61.1^{+2.6\%}_{-1.5\%} \times 10^3$	$63.8^{+1.7\%}_{-3.0\%} \times 10^3$	1.04
300	$858^{+1.8\%}_{-1.6\%}$	$576^{+1.2\%}_{-1.1\%}$	$605^{+0.8\%}_{-0.7\%}$	1.05
500	$129^{+1.6\%}_{-1.3\%}$	$78.4^{+3.5\%}_{-0.3\%}$	$84.7^{+2.0\%}_{-1.9\%}$	1.08
1000	$7.49^{+2.0\%}_{-2.1\%}$	$4.01^{+3.5\%}_{-10\%}$	$4.229^{+2.4\%}_{-2.9\%}$	1.05
3000	$10.9^{+3.6\%}_{-4.5\%} \times 10^{-3}$	$5.05^{+3.3\%}_{-4.5\%} \times 10^{-3}$	$5.26^{+0.5\%}_{-5.3\%} \times 10^{-3}$	1.04

Table 5. Same as in table 4 but for the SM charged current DY process.

$M_{\mu\mu}$ [GeV]	$\sigma_{\text{Tot.}}^{\text{NLO}}$ [fb]	$\sigma_{(\text{Veto})}^{\text{NLO+PS}}$ [fb]	$\sigma_{(\text{Veto})}^{\text{NLO+NNLL}}$ [fb]	$K_{\text{NLO+PS}}^{\text{NLO+NNLL(Veto)}}$
10	$7.65^{+24\%}_{-32\%} \times 10^6$	$7.50^{+15\%}_{-21\%} \times 10^6$	$9.50^{+5.2\%}_{-9.3\%} \times 10^6$	1.27
30	$2.13^{+7.2\%}_{-13\%} \times 10^6$	$1.91^{+5.4\%}_{-4.6\%} \times 10^6$	$2.09^{+2.7\%}_{-5.0\%} \times 10^6$	1.09
50	$1.80^{+5.4\%}_{-11\%} \times 10^6$	$1.61^{+4.7\%}_{-3.5\%} \times 10^6$	$1.75^{+2.4\%}_{-4.5\%} \times 10^6$	1.09
100	$73.7^{+4.2\%}_{-8.2\%} \times 10^3$	$62.7^{+2.9\%}_{-3.2\%} \times 10^3$	$69.2^{+2.2\%}_{-3.9\%} \times 10^3$	1.10
300	$696^{+1.7\%}_{-1.8\%}$	$481^{+1.9\%}_{-1.2\%}$	$510^{+0.7\%}_{-0.5\%}$	1.06
500	$111^{+1.6\%}_{-1.2\%}$	$69.6^{+5.3\%}_{-0.4\%}$	$72.8^{+1.5\%}_{-1.4\%}$	1.05
1000	$6.97^{+1.6\%}_{-1.7\%}$	$3.87^{+0.1\%}_{-0.3\%}$	$3.99^{+2.1\%}_{-2.5\%}$	1.03

Table 6. Same as in table 4 but for the SM neutral current DY process.

The difference in accuracy with respect to the two other processes is necessary to avoid double counting of diagrammatic contributions that appear both in the NLO corrections to the tW process and in the LO contributions to top-antitop production when using the five-flavor scheme. Whilst a consistent matching of these two channels at the NLO+PS accuracy has recently been achieved [77, 78], such a precision is unnecessary for our purposes. The listed top processes are intrinsically finite at the Born level and thus do not need regulating selections in the collider signature definitions. For $t\bar{t}$ and single t production and our scale choices, we apply K -factors of $K = 1.2$ to account for NNLO and threshold resummation corrections beyond NLO [79–81].

Unlike the color-singlet signal and background processes, the top quark channels inherently give rise to jets that are well-described by fixed order perturbation theory. At the Born level, the final-state partons that evolve into jets possess p_T comparable to the hard process scale and are emitted at wide-angles with respect to the beam axis. Jet vetoes applied to the top quark background can thus be well-approximated without the need for resummation beyond the PS. Measurements of low jet multiplicities in $t\bar{t}$ production at 8 TeV for instance show good agreement with the theory once both experimental and theoretical uncertainties are accounted for [82]. We consequently model the application of jet vetoes to the top background by simply imposing a p_T selection on the hardest jet present within the detector fiducial volume after parton showering.

5 Missing transverse energy and jet modeling

In this section, we discuss the impact of missing energy and jet modeling in $W' \rightarrow \ell\nu$ searches with jet vetoes. In particular, we comment on the use of exclusive versus inclusive \cancel{E}_T definitions by ATLAS [4] and CMS [5] respectively, as well as exclusive vetoes (with e.g., anti- b -tagging) versus inclusive vetoes (i.e., which are flavor-summed). Exclusive \cancel{E}_T is noteworthy as it is potentially a large source of systematic uncertainty that has been previously neglected.

5.1 Exclusive and inclusive missing transverse momentum

At 13 TeV, the CMS collaboration uses inclusive \cancel{E}_T in its $W' \rightarrow \ell\cancel{E}_T$ search. It is defined in the usual sense as the norm of the transverse momentum imbalance of all visible particles [5],

$$\cancel{E}_T \equiv |\vec{\cancel{p}}_T| \quad \text{where} \quad \vec{\cancel{p}}_T = - \sum_{X \in \{\text{visible}\}} \vec{p}_T^X. \quad (5.1)$$

Invisible particles are not restricted to light neutrinos, but also include ultra-soft and ultra-collinear objects as well as anything absorbed by inactive detector material, like screws and nails. Furthermore, particle identification is based on the particle-flow technique [83, 84], which exploits the detector’s magnetic field and its tracker and electromagnetic calorimeter resolution. “Blocks” with known momentum are constructed from tracks and calorimeter clusters and then identified as particle candidates. In a loose sense, the \cancel{E}_T of a CMS event is known before its particle content.




Figure 9. Normalized distributions for the SM process $pp \rightarrow W \rightarrow \mu^\pm \nu_\mu$ in 13 TeV LHC collisions at the NLO+PS accuracy, with respect to (a) the ratio of reconstructed \cancel{E}_T to the neutrino p_T and (b) the transverse mass reconstructed from the $(\ell\cancel{p})$ -system, using inclusive and exclusive \cancel{E}_T definitions and assuming various jet radii and minimum jet p_T .

The ATLAS W' boson search of ref. [4] takes a complementary approach to defining \cancel{E}_T by building the $\vec{\cancel{p}}_T$ vector from reconstructed objects already satisfying kinematic and fiducial requirements,

$$\cancel{E}_T^{\text{Exclusive}} \equiv |\vec{\cancel{p}}_T^{\text{Exclusive}}|, \quad \text{where} \quad \vec{\cancel{p}}_T^{\text{Exclusive}} = - \sum_{X \in \{\text{visible leptons, high-}p_T \text{ jets, photons}\}} \vec{p}_T^X. \quad (5.2)$$

Specifically, the hadronic contribution includes only $R = 0.4$ anti- k_T jets with $p_T > 20$ GeV. Unlike the CMS procedure where the \cancel{E}_T is independent of additional QCD splittings (ignoring pathological regions of phase space that correspond, for example, to screws and nails), the definition of eq. (5.2) by construction does not sum over an arbitrary number of low energy jets, and hence is an exclusive quantity. We now investigate the phenomenological consequences of using eq. (5.2) as a missing energy definition.

The first impact of using exclusive \cancel{E}_T is the broadening of all \cancel{E}_T -based observables. At the LHC, a pp collision may produce up to a few high- p_T objects but is typically dominated by the high multiplicity production of lower energy hadrons, i.e., the underlying event and real emissions off the hard process. While on average particle production is uniform in the transverse plane, radiation is distributed asymmetrically on an event-by-event basis. Hence, when clustered with a separation scale of R , up to a few moderate-to-high- p_T jets are balanced transversely by many more low- p_T jets. Excluding the low- p_T jets from the missing energy definition, as done in eq. (5.2), thus injects additional missing energy that is weakly correlated with any real source of \cancel{E}_T that may originate from the hard process. The issue is exacerbated for smaller jet radii $R' < R$, which distributes the same momentum from the hard and underlying processes over a larger jet multiplicity, thereby decreasing the average jet p_T .

To demonstrate this phenomenon, we consider, at NLO+PS-accuracy, SM W production and decay process

$$pp \rightarrow W \rightarrow \mu \nu_\mu. \tag{5.3}$$

In figure 9a, we present normalized distributions for the ratio of the missing transverse energy (both in the inclusive and exclusive cases) to the transverse momentum of the final-state neutrino. For both $R = 0.4$ and $R = 1.0$ jets, the inclusive \cancel{E}_T definition of eq. (5.1) describes the light neutrino p_T very well, as one would expect, with more than 99% of the distribution being contained within $0.9 < \cancel{E}_T/p_T^\nu < 1.1$. On the other hand, requiring jets to satisfy $p_T^j > 5\text{-}20$ GeV reduces this fraction to 50-80%, the rest of the distribution being smeared evenly around the origin. The broadening is alleviated for larger jet radii, due to their inherently more inclusive nature. However, the change is marginal for larger jet p_T requirements.

In figure 9b is the distribution of the transverse mass reconstructed from the (μp_T) -system, as defined below in eq. (6.10). For the inclusive case, about 25% of the distribution is contained in the bin spanning $75 \text{ GeV} < M_T < 82.5 \text{ GeV}$. In contrast, for various exclusive \cancel{E}_T definitions, the peaks drop to about consisting only of 16-18% of the distribution. Once again a larger R choice tames the effects due to increase inclusiveness. Consequently, using exclusive \cancel{E}_T definitions can undermine efforts to search for resonant structures when using \cancel{E}_T -based observables.

A second impact of imposing a $p_T^j > p_T^{\text{min}}$ requirement in building \cancel{E}_T is the generation of non-global logarithms (NGLs) of the form $\alpha_s \log [\cancel{E}_T / (\cancel{E}_T - p_T^{\text{min}})]$. NGLs arise when the phase space associated with virtual corrections of an exclusive observable is different from the phase space associated with the real corrections [85]. In the inclusive limit, e.g., $p_T^{\text{min}} \rightarrow 0$, such logarithms vanish. Intuitively, NGLs can be understood by imagining a jet that just marginally satisfies the p_T^{min} threshold. Virtual corrections do not change kinematics and therefore leave the missing energy unchanged. However, there exists a corresponding phase space configuration consisting of a wide-angle emission that brings the initial jet below p_T^{min} . Such objects are ignored by the \cancel{E}_T definition of eq. (5.2) and are therefore responsible for inducing shifts in the reconstructed \cancel{E}_T of order $\Delta \cancel{E}_T \sim \mathcal{O}(p_T^{\text{min}})$. This mismatch of the virtual and real phase space configurations can lead to potentially large logarithms that would otherwise vanish for inclusive observables. Further discussion of resumming such NGLs and the residual scale uncertainty are beyond the scope of this study. Nonetheless, it is clear that the uncertainty associated with the \cancel{E}_T reported by ATLAS in ref. [4] is an underestimation.

For our purposes, we employ the inclusive \cancel{E}_T definition of eq. (5.1).

5.2 Exclusive and inclusive jet veto

As for many new physics searches, the leading SM backgrounds for a $pp \rightarrow W_{\text{SSM}} \rightarrow \ell \nu_\ell$ signal include single and pair production of top quarks shown in eq. (4.6). As such, it is common practice to apply anti- b -tagging and reject any event featuring a reconstructed b -tagged jet. At the LHC, b -tagging methods have typical identification efficiencies of 70-80% associated with mis-tagging rates of about 1.5-10% [86]. It is thus pertinent to determine whether it is valuable to replace exclusive single-flavor jet vetoes by inclusive flavor-agnostic jet vetoes.




Figure 10. Normalized p_T distribution of the hardest jet for W' boson and single top (tX) production in proton-proton collision at a center-of-mass energy of 13 TeV. We adopt a jet radius fixed either to $R = 0.4$ (a,c) or to $R = 1$ (b,d), and we either include a b -jet veto assuming a tagging/mis-tagging efficiency of 100%/0% (c,d) or not (a,b).

	R	$W_{\text{SSM}}(300 \text{ GeV})$	$W_{\text{SSM}}(3 \text{ TeV})$	$t\bar{t}$	t_j	$t_{\mu\nu}$
$\sigma_{\text{Tot.}}[\text{No. of } \mu \geq 1] \text{ [fb]}$	N/A	24×10^3	15	120×10^3	29×10^3	6.1×10^3
b -Jet Veto [fb] $p_T^{\text{Veto}} = 30 \text{ GeV}, \epsilon^b = 100\%$	0.4	$24 \times 10^3 (> 99\%)$	15 (98%)	$6.9 \times 10^3 (5.6\%)$	$5.5 \times 10^3 (19\%)$	990 (16%)
	1.0	$24 \times 10^3 (99\%)$	15 (98%)	$6.1 \times 10^3 (5.0\%)$	$4.2 \times 10^3 (14\%)$	720 (12%)
Inclusive Jet Veto [fb] $p_T^{\text{Veto}} = 30 \text{ GeV}$	0.4	$18 \times 10^3 (74\%)$	7.4 (48%)	570 (0.5%)	$1.3 \times 10^3 (5\%)$	160 (3%)
	1.0	$16 \times 10^3 (68\%)$	6.3 (41%)	140 (0.1%)	670 (2%)	48 (0.8%)
Inclusive Jet Veto [fb] $p_T^{\text{Veto}} = 40 \text{ GeV}$	0.4	$19 \times 10^3 (80\%)$	8.6 (56%)	$2.1 \times 10^3 (2\%)$	$3.5 \times 10^3 (12\%)$	400 (7%)
	1.0	$18 \times 10^3 (75\%)$	7.6 (49%)	610 (0.5%)	$2.1 \times 10^3 (7\%)$	150 (2%)
Inclusive Jet Veto [fb] $p_T^{\text{Veto}} = 50 \text{ GeV}$	0.4	$20 \times 10^3 (84\%)$	9.5 (62%)	$6.2 \times 10^3 (5\%)$	$7.3 \times 10^3 (25\%)$	820 (14%)
	1.0	$20 \times 10^3 (81\%)$	8.6 (56%)	$2.0 \times 10^3 (2\%)$	$4.8 \times 10^3 (16\%)$	350 (6%)

Table 7. Signal and top quark background contributions to the inclusive $pp \rightarrow \mu^\pm X$ cross section [fb] and efficiencies (%) at $\sqrt{s} = 13 \text{ TeV}$ without any jet veto (second line), an exclusive b -jet veto with $p_T^{\text{Veto}} = 30 \text{ GeV}$ and 100% tagging/0% mis-tagging efficiency (third line), and an inclusive jet veto with $p_T^{\text{Veto}} = 30, 40, 50 \text{ GeV}$ (fourth, fifth, sixth line) for representative jet distance measures $R = 0.4, 1$.

To assess this, we consider the following W' and top quark production processes at 13 TeV,

$$\begin{aligned}
 pp \rightarrow W'(300 \text{ GeV}) &\rightarrow \mu \nu_\mu && \text{at NLO+PS,} \\
 pp \rightarrow W'(3 \text{ TeV}) &\rightarrow \mu \nu_\mu && \text{at NLO+PS,} \\
 pp \rightarrow t\bar{t} &\rightarrow \mu^\pm + X && \text{at NLO+PS,} \\
 pp \rightarrow tj &\rightarrow \mu^\pm + X && \text{at NLO+PS,} \\
 pp \rightarrow t\mu^\pm\nu_\mu &\rightarrow \mu^\pm + X && \text{at LO+PS.}
 \end{aligned}
 \tag{5.4}$$

As in section 4.3, the differences in the formal accuracies of each calculation allow us to avoid any possible double counting of diagrams. We cluster final states into jets as prescribed in section 2.2.

In figure 10a, we show the normalized p_T distributions of the hardest jet from the five processes of eq. (5.4), using a jet radius of $R = 0.4$. For the top backgrounds, a wide plateau can be seen at $p_T^j \sim m_t(1 - M_W^2/m_t^2)/2 \sim 65 - 70$ GeV, which indicates that the hardest jet in top production is indeed often a b -jet. For the W' processes, jets are characteristically at a lower p_T value since their production is entirely occurring via initial-state radiation and they are thus preferentially soft or collinear to the beam axis. In figure 10b, a jet radius of $R = 1.0$ is used and all distributions are expectedly shifted to higher p_T values.

Assuming an ideal b -jet tagging efficiency of $\varepsilon^b = 100\%$ and a 0% mis-tag rate of a lighter jet as a b -jet, we present in figure 10c for $R = 0.4$ and figure 10d for $R = 1.0$ the same distributions, but after rejecting all events featuring at least one b -jet with a transverse momentum satisfying $p_T^b > p_T^{\text{Veto}} = 30$ GeV. Even in this ideal scenario jets associated with top quarks are still characteristically more energetic, with $p_T^j \sim M_W/2$, than jets associated with W' production. This is related to the sizable single top tj process which proceeds through a t -channel W -boson exchange.

We summarize our findings in table 7. Here we present the total inclusive production cross section (including decays to at least one μ) for all processes in eq. (5.4) (second line), after applying an exclusive jet veto (third line) with $p_T^{\text{Veto}} = 30$ GeV and 100% tagging/0% mis-tagging efficiency, as well as after alternatively applying an inclusive veto with $p_T^{\text{Veto}} = 30, 40, \text{ and } 50$ GeV (fourth, fifth, sixth lines). We assume jet radii of $R = 0.4$ (above) and $R = 1$ (lower). The corresponding selection efficiencies are shown in parentheses and are evaluated with respect to the total rates. We observe that applying more inclusive jet vetoes, in terms of both jet radius and flavor composition, considerably increases the signal-to-noise ratio. With respect to b -jet vetoes, inclusive vetoes of $p_T^{\text{Veto}} = 30 - 40$ GeV can further suppress top quark production by an additional factor of 2 – 50 at a modest signal rate cost of 15 – 50%.

6 Observability of W_{SSM} with jet vetoes at hadron colliders

We now investigate the impact of employing jet vetoes on the discovery potential of W' bosons in the $pp \rightarrow W' \rightarrow \ell\nu_\ell$ channel at the 13 TeV LHC. Simulation of background and signal samples is described in section 4. Our analysis follows, where possible, the 13 TeV SSM W' search methodology employed by the CMS collaboration [5, 7].

We emulate the detector inefficiencies by smearing the momenta of all stable charged leptons ($\ell = e, \mu$) and jets reconstructed from the stable hadrons. In all cases, the smearing profile is Gaussian [5, 87], but with different scaling profiles: for muons, the p_T deviation ($\sigma_{p_T^\mu}$) is parameterized by

$$\sigma_{p_T^\mu} = a_{p_T^\mu}^\mu p_T^2, \tag{6.1}$$

where $a_{p_T^\mu}^\mu$ is 10% TeV^{-1} and 20% TeV^{-1} for central muons ($|\eta^\mu| < 0.9$) and forward muons ($|\eta^\mu| > 0.9$), respectively [5, 7]. The smearing in p_T is then translated into a change of the energy so that the momentum direction is kept unmodified. Unlike its energy scale, the direction of an infinitely energetic stable lepton can still be measured.

Similarly, electron energy uncertainties are parameterized by [5, 7]

$$\sigma_{E^e} = b_E^e E \quad \text{with} \quad b_E^e = 4\%. \tag{6.2}$$

The difference in parameterizations is due to electron energies being determined via calorimeters whereas muon momenta are derived from curvature measurements in a magnetic field. For jets, we follow the 13 TeV CMS $tt+nj$ analysis [87] which exploits dedicated energy calibration and p_T resolution measurements [88]. Jet energies and p_T are smeared independently according to

$$\sigma_{\hat{O}^j} = b_{\hat{O}^j}^j \times \hat{O} \quad \text{for} \quad \hat{O} \in \{E, p_T\}, \tag{6.3}$$

where the forward (central) coefficient, associated with jet pseudorapidities satisfying $|\eta| > 3$ ($|\eta| < 3$) are fixed to $b_E^j = 3\%$ (5%) and $b_{p_T}^j = 10\%$ (20%) [88]. The change in the jet momentum is translated into a shift in the jet mass, leaving the jet direction unmodified.

6.1 Signal definition and event selection

To test the production of generic W' bosons at colliders, we focus on the process

$$pp \rightarrow W' \rightarrow \mu \nu_\mu \rightarrow \mu + \cancel{E}_T. \tag{6.4}$$

The jet veto is agnostic to the lepton flavor; we therefore restrict ourselves to the study of the muon channel for simplicity. As discussed in section 3.1, the proposed methodology holds generally for any color-singlet process in hadron collisions, including multi-boson and Higgs production. Moreover, the $e + \cancel{E}_T$ mode consists of a multi-jet background [7] and hence is further enhanced by a jet veto but is otherwise identical to the above channel. Applying jet vetoes to the $\tau + \cancel{E}_T$ final state is debatable due to τ leptons preferential decays to hadrons.

We identify stable leptons ℓ^\pm as hadronically isolated objects for which the sum of the total hadronic E_T within a distance of $\Delta R_{\ell X} < 0.3$ centered on the the lepton candidate is less than 10% of its E_T , i.e.,

$$\sum_{X \in \{\text{jets}\}} E_T^X / E_T^\ell < 0.1 \quad \text{for} \quad \Delta R_{\ell X} < 0.3. \tag{6.5}$$

We select events containing a single muon candidate meeting the following kinematic, fiducial, and leptonic isolation requirements [5, 7]:

$$p_T^\mu > 53 \text{ GeV}, \quad |\eta^\mu| < 2.4, \quad \Delta R_{\mu\ell} > 0.3. \quad (6.6)$$

We reject events with additional isolated electrons and muons satisfying

$$\begin{aligned} p_T^e > 25 \text{ GeV} & \quad \text{with} \quad |\eta^e| < 1.444 \quad \text{or} \quad 1.566 < |\eta^e| < 2.5, \\ p_T^\mu > 35 \text{ GeV} & \quad \text{with} \quad |\eta^\mu| < 2.4. \end{aligned} \quad (6.7)$$

Following the results of sections 4 and 5, we cluster stable hadrons into jets according to the anti- k_T algorithm [65] with a separation scale of $R = 1$. We base our jet veto on the efficiencies of table 7 and reject any event with one or more jets whose properties satisfy

$$p_T^j > p_T^{\text{Veto}} = 40 \text{ GeV} \quad \text{and} \quad |\eta^j| < 4.7. \quad (6.8)$$

We subscribe to the CMS inclusive \cancel{E}_T definition [83, 84], as given in eq. (5.1), and sum over all charged leptons (including non-isolated objects) as well as all clustered hadronic activity satisfying $p_T^{\text{Had.}} > 0.1 \text{ GeV}$ and $|\eta^{\text{Had.}}| < 4.7$.

The following selection is then performed to enhance the signal-over-noise S/B ratio [5],

$$|\Delta\phi(\vec{p}^\ell, \vec{\cancel{p}})| > 2.5 \quad \text{and} \quad 0.4 < p_T^\ell / \cancel{E}_T < 1.5, \quad (6.9)$$

where we respectively constrain the azimuthal separation between the selected muon and the missing momentum and the ratio of the lepton transverse momentum to the missing transverse energy. As longitudinal momenta of light neutrinos cannot be generically inferred in hadron collisions, the transverse mass (M_T) of the $(\ell\cancel{E}_T)$ -system,

$$M_T = \sqrt{2 p_T^\ell \cancel{E}_T [1 - \cos \Delta\phi(\vec{p}^\ell, \vec{\cancel{p}})]}, \quad (6.10)$$

is eventually used as a discriminating variable. K -factors accounting for QCD corrections beyond NLO are applied according to the prescriptions given in section 4.

6.2 Extended discovery potential and sensitivity at 13 TeV

To quantify the discovery potential of a positive $W' \rightarrow \mu\nu$ signal at the LHC, we use Gaussian statistics to define the significance of a would-be discovery as

$$\sigma^{\text{Discovery}} = \frac{n^s}{\sqrt{n^s + (1 + \delta_b)n^b}} \quad \text{where} \quad n^{s,b} = \mathcal{L} \times \sigma^{s,b}. \quad (6.11)$$

Here, $n^{s,b}$ represents the expected number of signal and background events given an integrated luminosity \mathcal{L} and a (fiducial) signal and background cross section $\sigma^{s,b}$. Conservatively, we have introduced a δ_b parameter to account for the potential systematic uncertainties, which we chose to be $\delta_b = 20\%$. While for the discovery potential we require $\sigma^{\text{Discovery}} > 5$, we impose $\sigma^{\text{Discovery}} < 2$ for approximately evaluating the 95% confidence level exclusion range.




Figure 11. Discovery potential for W' boson searches via the $W' \rightarrow \mu\nu$ channel. The results are presented in terms of the requisite luminosity for a 5σ statistical significance (a) of the signal over the SM background at the LHC (operating at a center-of-mass energy of 13 TeV) with (dash-dot) and without (dash) the use of a jet veto of $p_T^{\text{Veto}} = 40$ GeV in the analysis, assuming the W' coupling normalization of eq. (2.15), as well as in terms of a 95% confidence level upper limit on the signal strength (b) when assuming an integrated luminosity of 100 and 3000 fb^{-1} .

$M_{W'}$ [GeV]	30	50	300	500	1000	3000	5000
$\delta(S/B)$	9.8%	6.3%	23%	250%	-4.0%	-5.5%	-2.4%

Table 8. Change in signal-to-noise ratio [%] for $pp \rightarrow W' \rightarrow \mu + \cancel{E}_T$ searches at the 13 TeV LHC after applying a jet veto of $p_T^{\text{Veto}} = 40$ GeV.

In figure 11a, as a function of $M_{W'}$, we show the requisite integrated luminosity for obtaining a 5σ statistical significance (or equivalently to claim discovery of a signal) with (dash-dot) and without (dash) including a jet veto of $p_T^{\text{Veto}} = 40$ GeV in the analysis. As a benchmark, we assume SSM W' coupling to fermions as given in eq. (2.15), so that $\kappa_L^{q,\ell} = g'/g_{\text{SM}} = 2.6 \times 10^{-2}$. For light and moderate W' boson masses of $M_{W'} \in [30, 900]$ GeV, we observe a systematic, albeit marginally for the lighter cases, improvement in the discovery potential. For $M_{W'} \lesssim 100$ GeV, the signal-to-noise ratio slightly increases by about 6-10% when a jet veto is employed, whereas the improvement reaches approximately 20% and 200% for heavier W' boson masses of about 300 GeV or 500 GeV, respectively. This translates to requiring 5 – 90% less data to achieve the same 5σ statistical sensitivity as without a jet veto for this particular mass regime. The large variation in the utility of a jet veto is due to the relative contribution of the top quark and DY processes in the SM background. For small $M_{W'}$ values, non-colored-singlet backgrounds make up only $\mathcal{O}(10\%)$ of the total background, a number that grows dramatically for W' scenarios above the top quark mass threshold. For situations in which the W' boson mass is heavier than 700-800 GeV, the veto ceases to be useful as the SM background essentially vanishes.

Subsequently, the veto only acts to decrease the rates leaving the signal-to-noise ratio unchanged, which therefore worsens the sensitivity. Alternatively, we show in table 8 the corresponding changes in the signal-to-noise ratio when a jet veto is applied. While improvements are at the 10% level for light W' bosons, they drastically increase from 20% to 250% for moderate W' boson masses ranging from 300 GeV to 800 GeV, before worsening the search for the very massive W' bosons.

In figure 11b, we translate this discovery potential into a sensitivity on signal strength μ defined as

$$\mu = \sigma^{\text{Expected}} / \sigma^{pp \rightarrow W_{\text{SSM}} \rightarrow \mu + \nu}, \quad (6.12)$$

where σ^{Expected} is the expected fiducial signal cross section for any W' boson scenario that one may consider and $\sigma^{pp \rightarrow W_{\text{SSM}} \rightarrow \mu + \nu}$ is the analogous SSM boson cross section obtained when using the couplings of eq. (2.15). We obtain similar results to (a) and observe that jet vetoes can potentially improve the sensitivity by 5 – 70% for moderate W' boson lying in the 300 – 900 GeV mass window.

7 Summary and conclusion

The origin of tiny nonzero neutrinos masses, the particle nature of dark matter and the weakness of gravity are longstanding issues, among others, that can potentially be understood and studied at collider experiments via the probes for the existence of new W' and Z' gauge bosons. Due to their color-singlet nature, the QCD radiation pattern of W' and Z' boson production at hadron colliders is intrinsically softer than the W'/Z' mass scale and more collinear with respect to the beam axis than the pattern associated with the leading color non-singlet background processes. As a consequence, the sensitivity to color-singlet new physics searches can be improved with the usage of jet vetoes provided the QCD processes are a non-negligible fraction of the background.

As a proof of principle, we have studied at the 13 TeV LHC, the muonic signature of a generic W' signal,

$$pp \rightarrow W' \rightarrow \mu\nu, \quad (7.1)$$

focusing on the increased discovery potential gained by employing jet vetoes. We have systematically considered both signal and background processes at NLO+PS accuracy, and included, for color-singlet signal and background channels, the resummation of jet veto logarithms up to the NNLL accuracy with matching to NLO fixed-order results. This has necessitated the development of a new FEYNRULES model in which we have implemented in a generic fashion new W' and Z' gauge bosons with model-independent chiral couplings. See section 2 for more details. Associated model files are public available from the FEYNRULES model database [48].

We have investigated the impact of several classes of uncertainties that are attached to jet veto resummation calculations. We have probed the dependence on the choice of jet definition, which suggests larger jet radii ($R \sim 1$) lead to smaller uncertainties than smaller radii. See section 4 for additional details.

In section 5, we studied the dependence of our collider analysis on missing transverse energy definitions as well as the flavor-dependence of the jet vetoes. We have described how exclusive missing energy definitions, such as the one used in 13 TeV ATLAS analyses, broaden all missing-energy-based observables and subsequently leads to a decrease in experimental sensitivity. This choice of \cancel{E}_T additionally leads to the rise of a new class of non-global logarithms that are responsible for a potentially large theoretical uncertainties that have not been previously taken into account. On different lines, we have found that with respect to a b -jet veto of $p_T^{\text{Veto}} = 30$ GeV, flavor-agnostic jet vetoes of $p_T^{\text{Veto}} = 30 - 40$ GeV can further reduce single top and top-antitop quark production by a factor of 2 – 50 at a mild cost of the signal rate.

We have applied our finding to the specific case of a W_{SSM} boson, and observed that for a new physics coupling strength taken as large as allowed by the current constraints, $\kappa_L^{q,\ell} = g'/g_{\text{SM}} = 2.6 \times 10^{-2}$. The usage of jet vetoes can increase the signal-to-noise ratios by roughly 10% for very light bosons masses of 30 – 50 GeV and 25%-250% for moderately heavy bosons of 300 – 800 GeV. Beyond this, vetoes lose their usefulness as they decrease the signal rates by a large amount, leaving the almost vanishing SM backgrounds almost unaffected. Conversely, W_{SSM} -bosons could be discovered by using 2-10 times for moderate $M_{W'}$, the moderate mass range, in contrast to any other mass scale where the change is milder.

Acknowledgments

Thomas Becher, Lydia Brenner, Fabrizio Caola, Mihoko Nojiri, Cedric Weiland are thanked for valuable discussions. The work of RR was funded in part by the U.K. Science and Technology Facilities Council (STFC), the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 690575, and Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture (MEXT), Japan (Nos. 16H06492 and 16H03991 for M. M. Nojiri). The authors appreciate the hospitality of IHEP where this work was initiated. RR acknowledges the generous hospitality KEK and the European Union.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] G. Altarelli, B. Mele and M. Ruiz-Altaba, *Searching for new heavy vector bosons in $p\bar{p}$ colliders*, *Z. Phys.* **C 45** (1989) 109 [*Erratum ibid.* **C 47** (1990) 676] [[INSPIRE](#)].
- [2] ATLAS collaboration, *Search for new high-mass resonances in the dilepton final state using proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [ATLAS-CONF-2016-045](#), CERN, Geneva Switzerland, (2016).

- [3] CMS collaboration, *Search for a high-mass resonance decaying into a dilepton final state in 13 fb^{-1} of pp collisions at $\sqrt{s} = 13\text{ TeV}$* , [CMS-PAS-EXO-16-031](#), CERN, Geneva Switzerland, (2016).
- [4] ATLAS collaboration, *Search for new resonances decaying to a charged lepton and a neutrino in pp collisions at $\sqrt{s} = 13\text{ TeV}$ with the ATLAS detector*, [ATLAS-CONF-2016-061](#), CERN, Geneva Switzerland, (2016)
- [5] CMS collaboration, *Search for $SSM W'$ production, in the lepton+MET final state at a center-of-mass energy of 13 TeV* , [CMS-PAS-EXO-15-006](#), CERN, Geneva Switzerland, (2015).
- [6] CMS collaboration, *Search for W' decaying to tau lepton and neutrino in proton-proton collisions at $\sqrt{s} = 13\text{ TeV}$* , [CMS-PAS-EXO-16-006](#), CERN, Geneva Switzerland, (2016).
- [7] CMS collaboration, *Search for heavy gauge W' boson in events with an energetic lepton and large missing transverse momentum at $\sqrt{s} = 13\text{ TeV}$* , [arXiv:1612.09274](#) [INSPIRE].
- [8] D. Binosi and L. Theussl, *JaxoDraw: a graphical user interface for drawing Feynman diagrams*, *Comput. Phys. Commun.* **161** (2004) 76 [[hep-ph/0309015](#)] [INSPIRE].
- [9] V.D. Barger, K.-M. Cheung, T. Han and D. Zeppenfeld, *Single forward jet tagging and central jet vetoing to identify the leptonic WW decay mode of a heavy Higgs boson*, *Phys. Rev. D* **44** (1991) 2701 [*Erratum ibid.* **D 48** (1993) 5444] [INSPIRE].
- [10] T. Becher and M. Neubert, *Factorization and NNLL resummation for Higgs production with a jet veto*, *JHEP* **07** (2012) 108 [[arXiv:1205.3806](#)] [INSPIRE].
- [11] A. Banfi, G.P. Salam and G. Zanderighi, *NLL+NNLO predictions for jet-veto efficiencies in Higgs-boson and Drell-Yan production*, *JHEP* **06** (2012) 159 [[arXiv:1203.5773](#)] [INSPIRE].
- [12] A. Banfi, P.F. Monni, G.P. Salam and G. Zanderighi, *Higgs and Z-boson production with a jet veto*, *Phys. Rev. Lett.* **109** (2012) 202001 [[arXiv:1206.4998](#)] [INSPIRE].
- [13] S. Dawson, T. Han, W.K. Lai, A.K. Leibovich and I. Lewis, *Resummation effects in vector-boson and Higgs associated production*, *Phys. Rev. D* **86** (2012) 074007 [[arXiv:1207.4207](#)] [INSPIRE].
- [14] T. Becher, M. Neubert and L. Rothen, *Factorization and N^3LL_p +NNLO predictions for the Higgs cross section with a jet veto*, *JHEP* **10** (2013) 125 [[arXiv:1307.0025](#)] [INSPIRE].
- [15] P. Meade, H. Ramani and M. Zeng, *Transverse momentum resummation effects in W^+W^- measurements*, *Phys. Rev. D* **90** (2014) 114006 [[arXiv:1407.4481](#)] [INSPIRE].
- [16] P. Jaiswal and T. Okui, *Explanation of the WW excess at the LHC by jet-veto resummation*, *Phys. Rev. D* **90** (2014) 073009 [[arXiv:1407.4537](#)] [INSPIRE].
- [17] P.F. Monni and G. Zanderighi, *On the excess in the inclusive $W^+W^- \rightarrow \ell^+\ell^-\nu\bar{\nu}$ cross section*, *JHEP* **05** (2015) 013 [[arXiv:1410.4745](#)] [INSPIRE].
- [18] T. Becher, R. Frederix, M. Neubert and L. Rothen, *Automated NNLL+NLO resummation for jet-veto cross sections*, *Eur. Phys. J. C* **75** (2015) 154 [[arXiv:1412.8408](#)] [INSPIRE].
- [19] P. Jaiswal, P. Meade and H. Ramani, *Precision diboson measurements and the interplay of p_T and jet-veto resummations*, *Phys. Rev. D* **93** (2016) 093007 [[arXiv:1509.07118](#)] [INSPIRE].
- [20] A. Banfi et al., *Jet-vetoed Higgs cross section in gluon fusion at N^3LO +NNLL with small- R resummation*, *JHEP* **04** (2016) 049 [[arXiv:1511.02886](#)] [INSPIRE].

- [21] S. Dawson, P. Jaiswal, Y. Li, H. Ramani and M. Zeng, *Resummation of jet veto logarithms at $N^3LL_a + NNLO$ for W^+W^- production at the LHC*, *Phys. Rev. D* **94** (2016) 114014 [[arXiv:1606.01034](#)] [[INSPIRE](#)].
- [22] I.W. Stewart, F.J. Tackmann and W.J. Waalewijn, *Factorization at the LHC: from PDFs to initial state jets*, *Phys. Rev. D* **81** (2010) 094035 [[arXiv:0910.0467](#)] [[INSPIRE](#)].
- [23] I.W. Stewart, F.J. Tackmann and W.J. Waalewijn, *The beam thrust cross section for Drell-Yan at NNLL order*, *Phys. Rev. Lett.* **106** (2011) 032001 [[arXiv:1005.4060](#)] [[INSPIRE](#)].
- [24] C.F. Berger, C. Marcantonini, I.W. Stewart, F.J. Tackmann and W.J. Waalewijn, *Higgs production with a central jet veto at NNLL+NNLO*, *JHEP* **04** (2011) 092 [[arXiv:1012.4480](#)] [[INSPIRE](#)].
- [25] I.W. Stewart, F.J. Tackmann, J.R. Walsh and S. Zuberi, *Jet p_T resummation in Higgs production at NNLL'+NNLO*, *Phys. Rev. D* **89** (2014) 054001 [[arXiv:1307.1808](#)] [[INSPIRE](#)].
- [26] F.J. Tackmann, J.R. Walsh and S. Zuberi, *Resummation properties of jet vetoes at the LHC*, *Phys. Rev. D* **86** (2012) 053011 [[arXiv:1206.4312](#)] [[INSPIRE](#)].
- [27] M.A. Ebert et al., *Exploiting jet binning to identify the initial state of high-mass resonances*, *Phys. Rev. D* **94** (2016) 051901 [[arXiv:1605.06114](#)] [[INSPIRE](#)].
- [28] ATLAS collaboration, *Search for dark matter candidates and large extra dimensions in events with a photon and missing transverse momentum in pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector*, *Phys. Rev. Lett.* **110** (2013) 011802 [[arXiv:1209.4625](#)] [[INSPIRE](#)].
- [29] ATLAS collaboration, *Search for dark matter in events with a Z boson and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys. Rev. D* **90** (2014) 012004 [[arXiv:1404.0051](#)] [[INSPIRE](#)].
- [30] F.J. Tackmann, W.J. Waalewijn and L. Zeune, *Impact of jet veto resummation on slepton searches*, *JHEP* **07** (2016) 119 [[arXiv:1603.03052](#)] [[INSPIRE](#)].
- [31] ATLAS collaboration, *Measurement of W^+W^- production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector and limits on anomalous WWZ and WW γ couplings*, *Phys. Rev. D* **87** (2013) 112001 [*Erratum ibid.* **88** (2013) 079906] [[arXiv:1210.2979](#)] [[INSPIRE](#)].
- [32] CMS collaboration, *Measurement of the W^+W^- cross section in pp collisions at $\sqrt{s} = 7$ TeV and limits on anomalous WW γ and WWZ couplings*, *Eur. Phys. J. C* **73** (2013) 2610 [[arXiv:1306.1126](#)] [[INSPIRE](#)].
- [33] CMS collaboration, *Measurement of W^+W^- and ZZ production cross sections in pp collisions at $\sqrt{s} = 8$ TeV*, *Phys. Lett. B* **721** (2013) 190 [[arXiv:1301.4698](#)] [[INSPIRE](#)].
- [34] CMS collaboration, *Measurement of the W^+W^- cross section in pp collisions at $\sqrt{s} = 8$ TeV and limits on anomalous gauge couplings*, *Eur. Phys. J. C* **76** (2016) 401 [[arXiv:1507.03268](#)] [[INSPIRE](#)].
- [35] J.C. Collins, D.E. Soper and G.F. Sterman, *Factorization for short distance hadron-hadron scattering*, *Nucl. Phys. B* **261** (1985) 104 [[INSPIRE](#)].
- [36] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].

- [37] C.W. Bauer, S. Fleming, D. Pirjol and I.W. Stewart, *An effective field theory for collinear and soft gluons: heavy to light decays*, *Phys. Rev. D* **63** (2001) 114020 [[hep-ph/0011336](#)] [[INSPIRE](#)].
- [38] C.W. Bauer, D. Pirjol and I.W. Stewart, *Soft collinear factorization in effective field theory*, *Phys. Rev. D* **65** (2002) 054022 [[hep-ph/0109045](#)] [[INSPIRE](#)].
- [39] M. Beneke, A.P. Chapovsky, M. Diehl and T. Feldmann, *Soft collinear effective theory and heavy to light currents beyond leading power*, *Nucl. Phys. B* **643** (2002) 431 [[hep-ph/0206152](#)] [[INSPIRE](#)].
- [40] T. Golling et al., *Physics at a 100 TeV pp collider: beyond the Standard Model phenomena*, submitted to *Phys. Rept.* (2016) [[arXiv:1606.00947](#)] [[INSPIRE](#)].
- [41] N. Arkani-Hamed, T. Han, M. Mangano and L.-T. Wang, *Physics opportunities of a 100 TeV proton-proton collider*, *Phys. Rept.* **652** (2016) 1 [[arXiv:1511.06495](#)] [[INSPIRE](#)].
- [42] *Frontier Hadron Collider (FHC) webpage*, <https://fcc.web.cern.ch/Pages/Hadron-Collider.aspx>.
- [43] *International Workshop on Future High Energy Circular Colliders webpage*, <http://indico.ihep.ac.cn/event/3813>, (2013).
- [44] A. Alloul, N.D. Christensen, C. Degrande, C. Duhr and B. Fuks, *FeynRules 2.0 — a complete toolbox for tree-level phenomenology*, *Comput. Phys. Commun.* **185** (2014) 2250 [[arXiv:1310.1921](#)] [[INSPIRE](#)].
- [45] C. Degrande, *Automatic evaluation of UV and R_2 terms for beyond the Standard Model Lagrangians: a proof-of-principle*, *Comput. Phys. Commun.* **197** (2015) 239 [[arXiv:1406.3030](#)] [[INSPIRE](#)].
- [46] T. Hahn, *Generating Feynman diagrams and amplitudes with FeynArts 3*, *Comput. Phys. Commun.* **140** (2001) 418 [[hep-ph/0012260](#)] [[INSPIRE](#)].
- [47] C. Degrande, C. Duhr, B. Fuks, D. Grellscheid, O. Mattelaer and T. Reiter, *UFO — the Universal FeynRules Output*, *Comput. Phys. Commun.* **183** (2012) 1201 [[arXiv:1108.2040](#)] [[INSPIRE](#)].
- [48] B. Fuks and R. Ruiz, *W' and Z' bosons at NLO in QCD*, <http://feynrules.irmp.ucl.ac.be/wiki/WZPrimeAtNLO>.
- [49] S. Gopalakrishna, T. Han, I. Lewis, Z.-G. Si and Y.-F. Zhou, *Chiral couplings of W' and top quark polarization at the LHC*, *Phys. Rev. D* **82** (2010) 115020 [[arXiv:1008.3508](#)] [[INSPIRE](#)].
- [50] T. Han, I. Lewis, R. Ruiz and Z.-G. Si, *Lepton number violation and W' chiral couplings at the LHC*, *Phys. Rev. D* **87** (2013) 035011 [*Erratum ibid.* **D 87** (2013) 039906] [[arXiv:1211.6447](#)] [[INSPIRE](#)].
- [51] O. Mattelaer, M. Mitra and R. Ruiz, *Automated neutrino jet and top jet predictions at next-to-leading-order with parton shower matching in effective left-right symmetric models*, [arXiv:1610.08985](#) [[INSPIRE](#)].
- [52] M. Mitra, R. Ruiz, D.J. Scott and M. Spannowsky, *Neutrino jets from high-mass W_R gauge bosons in TeV-scale left-right symmetric models*, *Phys. Rev. D* **94** (2016) 095016 [[arXiv:1607.03504](#)] [[INSPIRE](#)].
- [53] Z. Sullivan, *Fully differential W' production and decay at next-to-leading order in QCD*, *Phys. Rev. D* **66** (2002) 075011 [[hep-ph/0207290](#)] [[INSPIRE](#)].

- [54] J. Alwall, C. Duhr, B. Fuks, O. Mattelaer, D.G. Öztürk and C.-H. Shen, *Computing decay rates for new physics theories with FeynRules and MadGraph 5_aMC@NLO*, *Comput. Phys. Commun.* **197** (2015) 312 [[arXiv:1402.1178](#)] [[INSPIRE](#)].
- [55] ATLAS collaboration, *Search for new phenomena in dijet mass and angular distributions from pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **754** (2016) 302 [[arXiv:1512.01530](#)] [[INSPIRE](#)].
- [56] CMS collaboration, *Search for narrow resonances decaying to dijets in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **116** (2016) 071801 [[arXiv:1512.01224](#)] [[INSPIRE](#)].
- [57] ATLAS collaboration, *Search for new light resonances decaying to jet pairs and produced in association with a photon or a jet in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, ATLAS-CONF-2016-070, CERN, Geneva Switzerland, (2016).
- [58] PARTICLE DATA GROUP collaboration, K.A. Olive et al., *Review of particle physics*, *Chin. Phys. C* **38** (2014) 090001 [[INSPIRE](#)].
- [59] V. Hirschi, R. Frederix, S. Frixione, M.V. Garzelli, F. Maltoni and R. Pittau, *Automation of one-loop QCD corrections*, *JHEP* **05** (2011) 044 [[arXiv:1103.0621](#)] [[INSPIRE](#)].
- [60] S. Frixione, Z. Kunszt and A. Signer, *Three jet cross-sections to next-to-leading order*, *Nucl. Phys. B* **467** (1996) 399 [[hep-ph/9512328](#)] [[INSPIRE](#)].
- [61] R. Frederix, S. Frixione, F. Maltoni and T. Stelzer, *Automation of next-to-leading order computations in QCD: the FKS subtraction*, *JHEP* **10** (2009) 003 [[arXiv:0908.4272](#)] [[INSPIRE](#)].
- [62] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [[INSPIRE](#)].
- [63] E. Conte, B. Fuks and G. Serret, *MadAnalysis 5, a user-friendly framework for collider phenomenology*, *Comput. Phys. Commun.* **184** (2013) 222 [[arXiv:1206.1599](#)] [[INSPIRE](#)].
- [64] M. Cacciari, G.P. Salam and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896 [[arXiv:1111.6097](#)] [[INSPIRE](#)].
- [65] M. Cacciari, G.P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063 [[arXiv:0802.1189](#)] [[INSPIRE](#)].
- [66] A. Buckley et al., *LHAPDF6: parton density access in the LHC precision era*, *Eur. Phys. J. C* **75** (2015) 132 [[arXiv:1412.7420](#)] [[INSPIRE](#)].
- [67] NNPDF collaboration, R.D. Ball et al., *Parton distributions for the LHC Run II*, *JHEP* **04** (2015) 040 [[arXiv:1410.8849](#)] [[INSPIRE](#)].
- [68] P. Skands, S. Carrazza and J. Rojo, *Tuning PYTHIA 8.1: the Monash 2013 tune*, *Eur. Phys. J. C* **74** (2014) 3024 [[arXiv:1404.5630](#)] [[INSPIRE](#)].
- [69] T. Becher, A. Broggio and A. Ferroglia, *Introduction to soft-collinear effective theory*, *Lect. Notes Phys.* **896** (2015) pp.1–206 [[arXiv:1410.1892](#)] [[INSPIRE](#)].
- [70] T. Becher and M. Neubert, *Drell-Yan production at small q_T , transverse parton distributions and the collinear anomaly*, *Eur. Phys. J. C* **71** (2011) 1665 [[arXiv:1007.4005](#)] [[INSPIRE](#)].
- [71] C. Muselli, S. Forte and G. Ridolfi, *Combined threshold and transverse momentum resummation for inclusive observables*, *JHEP* **03** (2017) 106 [[arXiv:1701.01464](#)] [[INSPIRE](#)].

- [72] M. Zeng, *Drell-Yan process with jet vetoes: breaking of generalized factorization*, *JHEP* **10** (2015) 189 [[arXiv:1507.01652](#)] [[INSPIRE](#)].
- [73] T. Becher and G. Bell, *Enhanced nonperturbative effects through the collinear anomaly*, *Phys. Rev. Lett.* **112** (2014) 182002 [[arXiv:1312.5327](#)] [[INSPIRE](#)].
- [74] M. Dasgupta, L. Magnea and G.P. Salam, *Non-perturbative QCD effects in jets at hadron colliders*, *JHEP* **02** (2008) 055 [[arXiv:0712.3014](#)] [[INSPIRE](#)].
- [75] M. Dasgupta, F. Dreyer, G.P. Salam and G. Soyez, *Small-radius jets to all orders in QCD*, *JHEP* **04** (2015) 039 [[arXiv:1411.5182](#)] [[INSPIRE](#)].
- [76] R. Ruiz, *QCD corrections to pair production of type III seesaw leptons at hadron colliders*, *JHEP* **12** (2015) 165 [[arXiv:1509.05416](#)] [[INSPIRE](#)].
- [77] T. Ježo, J.M. Lindert, P. Nason, C. Oleari and S. Pozzorini, *An NLO+PS generator for $t\bar{t}$ and Wt production and decay including non-resonant and interference effects*, *Eur. Phys. J. C* **76** (2016) 691 [[arXiv:1607.04538](#)] [[INSPIRE](#)].
- [78] T. Ježo and P. Nason, *On the treatment of resonances in next-to-leading order calculations matched to a parton shower*, *JHEP* **12** (2015) 065 [[arXiv:1509.09071](#)] [[INSPIRE](#)].
- [79] M. Czakon, P. Fiedler and A. Mitov, *Total top-quark pair-production cross section at hadron colliders through $O(\alpha_s^4)$* , *Phys. Rev. Lett.* **110** (2013) 252004 [[arXiv:1303.6254](#)] [[INSPIRE](#)].
- [80] N. Kidonakis, *NNLL resummation for s-channel single top quark production*, *Phys. Rev. D* **81** (2010) 054028 [[arXiv:1001.5034](#)] [[INSPIRE](#)].
- [81] N. Kidonakis, *NNLL threshold resummation for top-pair and single-top production*, *Phys. Part. Nucl.* **45** (2014) 714 [[arXiv:1210.7813](#)] [[INSPIRE](#)].
- [82] CMS collaboration, *Measurement of the differential production cross section for top-quark pairs as a function of jet multiplicity in the lepton+jets final state at $\sqrt{s} = 8$ TeV with the CMS detector*, **CMS-PAS-TOP-15-006**, CERN, Geneva Switzerland, (2015).
- [83] CMS collaboration, *Particle-flow event reconstruction in CMS and performance for jets, taus and MET*, **CMS-PAS-PFT-09-001**, CERN, Geneva Switzerland, (2009).
- [84] CMS collaboration, *Commissioning of the particle-flow event reconstruction with the first LHC collisions recorded in the CMS detector*, **CMS-PAS-PFT-10-001**, CERN, Geneva Switzerland, (2010).
- [85] M. Dasgupta and G.P. Salam, *Resummation of nonglobal QCD observables*, *Phys. Lett. B* **512** (2001) 323 [[hep-ph/0104277](#)] [[INSPIRE](#)].
- [86] CMS collaboration, *Identification of b-quark jets with the CMS experiment*, **2013 JINST 8 P04013** [[arXiv:1211.4462](#)] [[INSPIRE](#)].
- [87] CMS collaboration, *Measurement of $t\bar{t}$ production with additional jet activity, including b-quark jets, in the dilepton decay channel using pp collisions at $\sqrt{s} = 8$ TeV*, *Eur. Phys. J. C* **76** (2016) 379 [[arXiv:1510.03072](#)] [[INSPIRE](#)].
- [88] CMS collaboration, *Determination of jet energy calibration and transverse momentum resolution in CMS*, **2011 JINST 6 P11002** [[arXiv:1107.4277](#)] [[INSPIRE](#)].