

RECEIVED: January 2, 2018

REVISED: June 25, 2018

ACCEPTED: June 28, 2018

PUBLISHED: July 13, 2018

A viable QCD axion in the MeV mass range

Daniele S.M. Alves^{a,b,c} and Neal Weiner^a

^aCenter for Cosmology and Particle Physics, Department of Physics, New York University,
New York, NY 10003, U.S.A.

^bDepartment of Physics, Princeton University,
Princeton, NJ 08544, U.S.A.

^cTheoretical Division, Los Alamos National Laboratory,
Los Alamos, NM 87545, U.S.A.

E-mail: spier@lanl.gov, neal.weiner@nyu.edu

ABSTRACT: The QCD axion is one of the most compelling solutions of the strong CP problem. There are major current efforts into searching for an ultralight, *invisible* axion, which is believed to be the only phenomenologically viable realization of the QCD axion. *Visible* axions with decay constants at or below the electroweak scale are believed to have been long excluded by laboratory searches. Considering the significance of the axion solution to the strong CP problem, we revisit experimental constraints on QCD axions in the $\mathcal{O}(10\text{ MeV})$ mass window. In particular, we find a variant axion model that remains compatible with existing constraints. This model predicts new states at the GeV scale coupled hadronically, and a variety of low-energy axion signatures, such as rare meson decays, nuclear de-excitations via axion emission, and production in e^+e^- annihilation and fixed target experiments. This reopens the possibility of solving the strong CP problem at the GeV scale.

KEYWORDS: Beyond Standard Model, CP violation, Chiral Lagrangians, Kaon Physics

ARXIV EPRINT: [1710.03764](https://arxiv.org/abs/1710.03764)

Contents

1	Introduction	1
2	Constraints on generic MeV axions	2
2.1	Constraints amenable to model-building	3
2.2	Pion-phobia	4
2.3	Non-robust/uncertain constraints	4
3	Axion-meson mixings in χPT @ LO	5
4	Constraints from pion decays	7
4.1	The KTeV anomaly	8
5	Constraints from charged kaon decays	10
5.1	Experimental bounds on $K^+ \rightarrow \pi^+(a \rightarrow e^+e^-)$	10
5.2	Estimation of the amplitude for $K^+ \rightarrow \pi^+a$	12
5.3	Axion-eta mixing	15
6	The physical axion current	17
6.1	Axion-photon coupling	19
6.2	Axion-nucleon couplings and nuclear de-excitations	20
7	Axion-electron coupling	24
7.1	Beam dump constraints	24
7.2	Constraints from $(g-2)_e$	25
7.3	Constraints from searches for dark photons	26
8	Other constraints	29
9	GeV scale completions of MeV axion models	30
10	EW scale completions of MeV axion models	32
11	Discussion	34

1 Introduction

The Peccei-Quinn (PQ) mechanism is arguably the most compelling solution of the strong CP problem. Soon after it was originally proposed [1, 2], it was realized that a light pseudoscalar would emerge in the infrared spectrum as a manifestation of the underlying PQ mechanism - the QCD axion [3, 4]. As the parameters characterizing the QCD axion,

such as mass and decay constant, span several orders of magnitude, its phenomenology changes dramatically, with implications ranging from hadronic physics to astrophysics and cosmology. For the past four decades, this rich phenomenology has been explored over an ever broadening range of decay constants, $\mathcal{O}(\text{GeV}) \lesssim f_a \lesssim \mathcal{O}(M_{\text{Pl}})$ [5–10], and there is still an ongoing and vigorous experimental effort to test the QCD axion.

The existing consensus is that *visible* QCD axions, i.e., those with decay constants at or below the weak scale, have long been excluded by laboratory searches [11, 12], such as beam dump experiments, rare meson decays, and nuclear de-excitations.¹ This has motivated the formulation of *invisible* axion models [17–20], which, combined with further astrophysical bounds from stellar evolution, CMB and BBN, redirected experimental efforts to extremely weakly coupled axions ($f_a \gtrsim 10^9 \text{ GeV}$) [21–32], which are ultra-light ($m_a \lesssim 10^{-3} \text{ eV}$), and could be non-thermal dark matter [33–35].

Given the significance of the QCD axion and the many resources dedicated to probing its existence, it is important to ensure that no caveats have been overlooked, and that no gaps in already probed regions of parameter space were left uncovered. Motivated by this, we revisit constraints on the MeV mass window for the QCD axion and its variants, and discuss a particular realization of the QCD axion that has not yet been definitively excluded. Our results reopen the possibility that the strong CP problem might be solved below the weak scale, and suggest new, hadronically-coupled degrees of freedom at the GeV scale.

In sections 2–5 of this paper, we refute previous, premature conclusions that the MeV mass range for the QCD axion has been completely ruled out. We then discuss a viable axion variant and its couplings to photons, nucleons, and electrons, and the relevant experimental implications (sections 6–8). Finally, we comment on UV completions of such variants and associated phenomenology (sections 9–10). For an outline of this paper, vide table of contents.

2 Constraints on generic MeV axions

Generic constraints on QCD axions in the MeV mass window can be broadly classified into three categories: (i) amenable to model-building, (ii) plagued by large hadronic uncertainties, and (iii) evaded only by *pion-phobia*. The first category of constraints can be evaded by well-established model-building tools. Constraints in the second category are more difficult to avoid with model-building, but suffer from significant uncertainties which preclude them from fully and unambiguously ruling out the MeV mass range for the QCD axion. Finally, the third category encompasses the strongest constraints, which can only be avoided by a special class of axion variants which are *pion-phobic*, i.e., which have suppressed mixing with the neutral pion. While this can be achieved by model-building to some degree, the extreme *pion-phobia* needed to avoid exclusion also depends critically on the light quark mass ratio being close to a ratio of Peccei-Quinn charges (this will become

¹For $f_a \gtrsim 100 \text{ GeV}$, the QCD axion is also constrained by stellar evolution [13–15], and most recently by a combination of the CMB power spectrum and primordial ^4He and D/H abundances [16].

clear in section 4). Indeed the most up-to-date determinations of m_u/m_d indicate that it is very close 1/2, making extreme *pion-phobia* a realistic possibility.

2.1 Constraints amenable to model-building

We start by discussing the main experimental observables that have excluded generic QCD axions in the MeV mass window. Despite being severe, these constraints can be “model-built away” by deviating from generic models.

If the axion couples to heavy quark flavors, such as charm or bottom, it is strongly constrained by radiative decays of quarkonia, such as $J/\psi \rightarrow \gamma a$ and $\Upsilon \rightarrow \gamma a$. Wilczek [36] showed that such decay widths can be related to the leptonic widths via:

$$\Gamma(J/\psi \rightarrow \gamma a) = \Gamma(J/\psi \rightarrow \mu^+ \mu^-) \frac{\lambda_c^2}{2\pi\alpha} C_{J/\psi}, \quad (2.1)$$

$$\Gamma(\Upsilon \rightarrow \gamma a) = \Gamma(\Upsilon \rightarrow \mu^+ \mu^-) \frac{\lambda_b^2}{2\pi\alpha} C_\Upsilon, \quad (2.2)$$

where λ_c and λ_b are the axion couplings to $\bar{c} i\gamma_5 c$ and $\bar{b} i\gamma_5 b$, respectively, and $C_{J/\psi}$, $C_\Upsilon \sim \mathcal{O}(1)$ encode QCD and relativistic corrections [37]. The MeV mass range for the QCD axion corresponds to decay constants f_a in the $\mathcal{O}(1 - 10)$ GeV range, and consequently to large couplings of the axion to charm and/or bottom quarks, namely, $\lambda_c \sim \mathcal{O}(m_c/f_a)$ and/or $\lambda_b \sim \mathcal{O}(m_b/f_a)$. With such large couplings, radiative decays of quarkonia to γa would dominate over leptonic modes, in gross contradiction with observation. In fact, bounds from quarkonia decays alone [38–42] were sufficient to exclude the entire parameter space of the original PQWW axion (see, for instance, [43]). These and other bounds led to much activity during the 1980’s in model building QCD axion *variants*, i.e., variations of the original PQWW axion that could evade existing constraints at the time and remain viable.

A class of visible axion variants that trivially evades quarkonia bounds are those that couple the axion exclusively to first generation quarks.² Nevertheless, they still have to contend with other constraints. For instance, if such axions have suppressed couplings to leptons and decay dominantly to a pair of photons, they are sufficiently long lived and hence robustly excluded by a variety of beam dump experiments [44–47] in the range $100 \text{ keV} \lesssim m_a \lesssim 30 \text{ MeV}$. Limits from beam dumps are substantially degraded if axions couple to electrons with strength $\mathcal{O}(m_e/f_a)$. In this case, axions heavier than 5–10 MeV become very shortly lived, $\tau_a \lesssim 10^{-13} \text{ s}$, and their decay products stop in the earth shielding before reaching the detector, evading this class of bounds. Short-lived axions decaying to $e^+ e^-$ also evade severe constraints from $K^+ \rightarrow \pi^+(a \rightarrow \text{invisible})$, whose branching ratio is bounded to be $\lesssim 4.5 \times 10^{-11}$ [48–50].

While coupling the axion to electrons is desirable, an analogous coupling to muons with strength $\mathcal{O}(m_\mu/f_a)$ would induce contributions to $(g-2)_\mu$ that would violate present bounds unless $f_a \gtrsim v_{\text{EW}}$. Therefore, in this work we shall restrict ourselves to variant axion models that couple exclusively the first generation fermions, namely, u , d , and e .

²At least at tree level — axion couplings to second and third generations, including flavor non-diagonal couplings, will invariably be generated radiatively upon electroweak and PQ symmetry breakings. These couplings however are sufficiently suppressed to avoid present bounds, and will be ignored for the remainder of this paper.

2.2 Pion-phobia

The mixing of the axion with the neutral pion poses a major challenge to the viability MeV axion models. Severe constraints on this mixing have been placed three decades ago, and the idea of avoiding them via *pion-phobia* is just as old [51].

Axion-pion mixing induces the rare decay process $\pi^+ \rightarrow e^+ \nu_e a \rightarrow e^+ \nu_e e^+ e^-$. The width is given by [52]:

$$\Gamma(\pi^+ \rightarrow e^+ \nu_e a) = \frac{\cos^2 \theta_c}{384\pi^3} G_F^2 m_\pi^5 \theta_{a\pi}^2, \quad (2.3)$$

where $\theta_{a\pi}$ is the axion-pion mixing angle, and θ_c is the Cabibbo angle. The SINDRUM collaboration [53] searched for this specific decay, and put bounds on $\text{Br}(\pi^+ \rightarrow e^+ \nu_e (a \rightarrow e^+ e^-))$ ranging from $(0.5 - 1) \times 10^{-10}$ in the mass range $m_a \sim (1 - 20)$ MeV. Using (2.3), this translates into a severe upper limit on $\theta_{a\pi}$:

$$|\theta_{a\pi}| \lesssim (0.5 - 0.7) \times 10^{-4}. \quad (2.4)$$

As we shall see in section 4, typical MeV axion variants are in conflict with (2.4), since they predict $\theta_{a\pi} \sim \mathcal{O}(f_\pi/f_a) \sim (0.5 - 10) \times 10^{-2}$. As a consequence, the experimental upper bound on $\theta_{a\pi}$ pushes viable models into special regions of parameter space where the axion is *pion-phobic*, and therefore has suppressed couplings to isovector currents.

2.3 Non-robust/uncertain constraints

Finally, some constraints are plagued by large hadronic uncertainties, and until those are under better control no robust exclusion claim can be made. That is the case of $a - \eta$ and $a - \eta'$ mixings, which provide the dominant contribution to $K^+ \rightarrow \pi^+ a$.

Unlike $K^+ \rightarrow \pi^+(a \rightarrow \text{invisible})$, the rare decay $K^+ \rightarrow \pi^+(a \rightarrow e^+ e^-)$ is much less constrained, $\text{Br}(K^+ \rightarrow \pi^+(a \rightarrow e^+ e^-)) \lesssim \mathcal{O}(10^{-6})$. In light of the allowed range (2.4) for $\theta_{a\pi}$, the contribution to this decay from $\pi^0 - a$ mixing easily satisfies the experimental bound. However, it has been claimed in the literature [54, 55] that the contribution to this amplitude from $a - \eta$ mixing is octet ($\Delta I = 1/2$) enhanced, and translates into a bound $\theta_{a\eta} \lesssim \mathcal{O}(10^{-4})$. Based on naïve estimates of $\theta_{a\eta}$ from leading order in chiral perturbation theory, it was then concluded that axions in the MeV mass range were hopelessly ruled out by K^+ decay bounds. In section 5, we will revisit these statements and constraints, and argue that due to the uncertainties involved in these estimates, previous claims of exclusion were overstated, and bounds from K^+ decays alone cannot definitively rule out the class of axion variants we consider.

Table 1 summarizes the main experimental constraints relevant for generic MeV axions, and also lists the reasons as to why the variant model we shall introduce avoids all present bounds. This is not a standalone table - we urge the reader to follow our arguments in the main body of the paper to become fully aware of all the assumptions and caveats implicit in the information contained in table 1.

experimental limit on observable	translates into bound on model	expectation for generic MeV axions (LO in χ PT)	expectation for MeV axion variant in (3.1) (NLO in χ PT)	reason
beam dumps (see figure 1)	$\tau_a \lesssim 10^{-13}$ s (see figure 1)	$\tau_a < 10^{-12}$ s if $ Q_e^{\text{PQ}} \sim \mathcal{O}(1)$ $\tau_a \gtrsim 10^{-11}$ s if $Q_e^{\text{PQ}} = 0$	$ Q_e^{\text{PQ}} \sim \mathcal{O}(1)$ $\tau_a \lesssim 10^{-13}$ s for $m_a \gtrsim 5-10$ MeV	MB
$\text{Br}(J/\Psi \rightarrow \gamma a) \lesssim 6 \times 10^{-2}$	$ Q_c^{\text{PQ}} \lesssim 0.25 \left(\frac{f_a}{\text{GeV}}\right)$	$ Q_c^{\text{PQ}} \sim \mathcal{O}(1)$	$Q_c^{\text{PQ}} = 0$	MB
$\text{Br}(T \rightarrow \gamma a) \lesssim 3 \times 10^{-4}$	$ Q_b^{\text{PQ}} \lesssim 0.8 \times 10^{-2} \left(\frac{f_a}{\text{GeV}}\right)$	$ Q_b^{\text{PQ}} \sim \mathcal{O}(1)$	$Q_b^{\text{PQ}} = 0$	MB
$-\Delta(g-2)_\mu < 0.5 \times 10^{-8}$	$ Q_\mu^{\text{PQ}} \lesssim 4 \times 10^{-3} \left(\frac{f_a}{\text{GeV}}\right)$	$ Q_\mu^{\text{PQ}} \sim \mathcal{O}(1)$	$Q_\mu^{\text{PQ}} = 0$	MB
$-\Delta(g-2)_e < 0.5 \times 10^{-11}$	$ Q_e^{\text{PQ}} \lesssim \mathcal{O}(\text{few})$ $g_{a\gamma\gamma} \lesssim \mathcal{O}(0.1) \text{ TeV}^{-1}$	$ Q_e^{\text{PQ}} \sim \mathcal{O}(1)$ $g_{a\gamma\gamma} \sim \mathcal{O}(0.1-1) \text{ TeV}^{-1}$	$ Q_e^{\text{PQ}} \sim \mathcal{O}(1)$ $g_{a\gamma\gamma} \sim \mathcal{O}(0.01-0.1) \text{ TeV}^{-1}$	MB PP, NLO
$\text{Br}(\pi^+ \rightarrow e^+ \nu(a \rightarrow e^+ e^-)) \lesssim 10^{-10}$	$ \theta_{a\pi} \lesssim (0.5-0.7) \times 10^{-4}$	$ \theta_{a\pi} \sim \mathcal{O}(0.01-0.1) \left(\frac{\text{GeV}}{f_a}\right)$	$\theta_{a\pi} \sim (0.2 \pm 3) \times 10^{-3} \left(\frac{\text{GeV}}{f_a}\right)$	PP, NLO
$\Delta \text{Br}(\pi^0 \rightarrow e^+ e^-) \lesssim 2 \times 10^{-8}$	$Q_e^{\text{PQ}} \times \theta_{a\pi} \lesssim 1.6 \times 10^{-4} \left(\frac{f_a}{\text{GeV}}\right)$			
$\text{Br}(K^+ \rightarrow \pi^+(a \rightarrow e^+ e^-)) \lesssim 10^{-5}-10^{-6}$	$ \theta_{a\eta_{ud}} \lesssim 10^{-4}$ if octet enhanced $ \theta_{a\eta_{ud}} \lesssim 10^{-2}$ if not	$ \theta_{a\eta_{ud}} \sim \mathcal{O}(10^{-3}-10^{-2}) \left(\frac{\text{GeV}}{f_a}\right)$	$\theta_{a\eta_{ud}} \sim (-2 \pm 8) \times 10^{-3} \left(\frac{\text{GeV}}{f_a}\right)$	NLO
$\text{Br}(K^+ \rightarrow \pi^+(a \rightarrow \gamma\gamma)) \lesssim 10^{-9}$ $\text{Br}(\Phi \rightarrow \gamma(a \rightarrow e^+ e^-)) \lesssim 5 \times 10^{-5}$	$ \theta_{a\eta_s} \lesssim \mathcal{O}(10^{-1})$	$ \theta_{a\eta_s} \sim \mathcal{O}(10^{-2}) \left(\frac{\text{GeV}}{f_a}\right)$	$\theta_{a\eta_s} \sim (1 \pm 2) \times 10^{-2} \left(\frac{\text{GeV}}{f_a}\right)$	NLO
$\text{Br}(K^+ \rightarrow \pi^+(a \rightarrow \text{inv})) \lesssim 0.5 \times 10^{-10}$	$\text{Br}(a \rightarrow \text{inv}) \lesssim \mathcal{O}(10^{-4})$ (assuming $\tau_a < 10^{-12}$ s)	$\text{Br}(a \rightarrow \nu\bar{\nu}) \sim (Q_\nu^{\text{PQ}} m_\nu / Q_e^{\text{PQ}} m_e)^2$	$\text{Br}(a \rightarrow \chi\chi) \sim (Q_\chi^{\text{PQ}} m_\chi / Q_e^{\text{PQ}} m_e)^2$ $\chi = \nu_{(e,\mu,\tau,s)}$, sub-MeV DM, ...	MB

Table 1. Summary of the most relevant existing bounds on MeV axions, as well as the conditions for experimental viability discussed in this paper. The 3rd column indicates the typical range in generic axion models for the quantity being constrained, estimated at leading order (LO) in chiral perturbation theory (χ PT). The 4th column contrasts it with the corresponding expectation for a short-lived, *pion-phobic* MeV axion which, as we argue in the text, remains compatible with experimental bounds. The 4th column takes into account large uncertainties from next-to-leading order (NLO) in χ PT. The fifth column indicates the reason for the difference in expected properties of generic models (3rd column) and of the viable variant (4th column), where: **MB** indicates that the quantity in the 4th column is achievable via model building; **NLO** indicates that the error in the quantity in the 4th column comes from large corrections at NLO in χ PT; and **PP** indicates that the quantity in the 4th column is suppressed due to accidental cancellations at LO in the axion-pion mixing angle $\theta_{a\pi}$, i.e., *pion-phobia*.

3 Axion-meson mixings in χ PT @ LO

We can trivially evade the constraints discussed in section 2.1 by restricting our investigation to *QCD axion variants coupled only to the first generation of SM fermions*. This assumption is implied for the remainder of this paper.

We are now in a position to introduce more concrete notation and review the extraction of axion-meson mixing angles to leading order in chiral perturbation theory (χ PT).

We define the axion couplings slightly above the QCD scale, where all heavy fermions (c , b and t) have been integrated out. We also choose a specific basis (motivated by UV completions to be discussed in section 9) in which the operators $(a/f_a) G\tilde{G}$ and $\partial_\mu(a/f_a) J_{\text{SM}}^{5\mu}$ are not present, or have been rotated away. With this choice of basis, the variant models we shall consider are unambiguously defined by the following couplings:

$$\mathcal{L}_a^{\text{eff}} = m_u e^i Q_u^{\text{PQ}} a/f_a uu^c + m_d e^i Q_d^{\text{PQ}} a/f_a dd^c + m_e e^i Q_e^{\text{PQ}} a/f_a ee^c + \text{h.c.} \quad (3.1)$$

Above, Q_f^{PQ} are Peccei-Quinn (PQ) charges that determine the transformation of

fermions $f = u, d, e$ under a PQ rotation. Below the QCD scale, (3.1) can be mapped onto the effective chiral Lagrangian by treating the axion as a complex phase of the quark masses. To leading order in χ PT, the axion couplings to the $U(3)_\chi$ meson nonet can be extracted from:

$$\mathcal{L}_\chi^{(0)} = \frac{f_\pi^2}{4} \text{Tr} [2BM_q(a)U + \text{h.c.}] - \frac{1}{2}M_0^2\eta_0^2, \quad (3.2)$$

where $f_\pi = 92 \text{ MeV}$, B has mass dimensions and is set by the QCD quark condensate, $M_0 \sim \mathcal{O}(\text{GeV})$ parameterizes the strong anomaly contribution to the mass of the $SU(3)_\chi$ singlet η_0 , $M_q(a)$ is the light quark mass matrix,³

$$M_q(a) \equiv \begin{pmatrix} m_u e^{iQ_u a/f_a} & & \\ & m_d e^{iQ_d a/f_a} & \\ & & m_s \end{pmatrix}, \quad (3.3)$$

and U is the non-linear representation of the meson nonet under chiral $U(3)_\chi$ symmetry

$$U \equiv \text{Exp} i \frac{\sqrt{2}}{f_\pi} \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta_8}{\sqrt{6}} + \frac{\eta_0}{\sqrt{3}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta_8}{\sqrt{6}} + \frac{\eta_0}{\sqrt{3}} & K^0 \\ K^- & \bar{K}^0 & -\frac{\eta_8}{\sqrt{3/2}} + \frac{\eta_0}{\sqrt{3}} \end{pmatrix}. \quad (3.4)$$

Expanding the leading order chiral Lagrangian (3.2) to quadratic terms in the fields and diagonalizing the mass matrix, we obtain:

$$\begin{aligned} m_a^2 &= (Q_u + Q_d)^2 \frac{B m_u m_d m_s}{(m_u m_d + m_u m_s + m_d m_s + \frac{6Bm_s}{M_0^2})} \frac{f_\pi^2}{f_a^2} \\ &= \frac{(Q_u + Q_d)^2}{(1 + \epsilon_{\eta\eta'})} \frac{m_u m_d}{(m_u + m_d)^2} \frac{m_\pi^2 f_\pi^2}{f_a^2}, \end{aligned} \quad (3.5)$$

with

$$\begin{aligned} \epsilon_{\eta\eta'} &\equiv \frac{m_u m_d}{m_s(m_u + m_d)} \left(1 + 6 \frac{B m_s}{M_0^2} \right) \\ &\approx \frac{m_u m_d}{(m_u + m_d)^2} \frac{m_\pi^2}{m_K^2} \left(1 + 6 \frac{m_K^2}{m_{\eta'}^2} \right) \simeq 0.04, \end{aligned} \quad (3.6)$$

and

$$\theta_{a\pi}^{(0)} = -\frac{1}{(1 + \epsilon_{\eta\eta'})} \left(\frac{(Q_u m_u - Q_d m_d)}{(m_u + m_d)} + \epsilon_{\eta\eta'} \frac{(Q_u - Q_d)}{2} \right) \frac{f_\pi}{f_a}, \quad (3.7)$$

$$\theta_{a\eta_8}^{(0)} = -\sqrt{\frac{3}{2}} \frac{\epsilon_{\eta\eta'}}{(1 + \epsilon_{\eta\eta'})} (Q_u + Q_d) \frac{f_\pi}{f_a}, \quad (3.8)$$

$$\theta_{a\eta_0}^{(0)} = -\sqrt{6} (Q_u + Q_d)^{-1} \frac{f_a}{f_\pi} \frac{m_a^2}{M_0^2}, \quad (3.9)$$

³From this point forward we will suppress the superscript in the PQ charges, $Q_f^{\text{PQ}} \rightarrow Q_f$, to lighten the notation.

where the superscript ⁽⁰⁾ on the mixing angles indicates that these expressions stem from leading order in χ PT. Note, in particular, that when any of the quark masses is taken to zero, both the axion mass, m_a , and its mixing with the $SU(3)_\chi$ singlet, $\theta_{a\eta_0}$, vanish. This is consistent with the fact that, in the limit of a massless quark, the axion becomes a *bona fide* Goldstone boson associated with a non-anomalous conserved current.

Previous studies in the literature have assumed that these leading order estimations of the axion-meson mixing angles were good enough approximations. Based on those, they went on to extract rates for processes involving the axion, and infer bounds that widely excluded all axion variants with $m_a \gtrsim \mathcal{O}(\text{MeV})$. The core argument of this paper is that these assumptions, while correct for a wide class of axion variants, fail under special circumstances and invalidate broad exclusion claims. In these special circumstances, the mixing angles in (3.7)–(3.9) receive $\mathcal{O}(1)$ corrections from the next order in the chiral expansion, introducing large uncertainties to the axion couplings and dramatically affecting the inference of bounds that depend on these mixing angles.

4 Constraints from pion decays

As discussed in section 2.2, bounds from $\pi^+ \rightarrow e^+ \nu_e a$ severely restrict viable MeV axions to the extreme *pion-phobic* region. In this section we will discuss under what conditions *pion-phobia* can be achieved, as well as implications for another relevant pion decay, namely, $\pi^0 \rightarrow e^+ e^-$.

The leading order expression for the axion mixing with the neutral pion in (3.7) indicates that, for generic $\mathcal{O}(1)$ values of $Q_{u,d}$, $\theta_{a\pi} \sim \mathcal{O}(f_\pi/f_a)$. For this generic situation, the axion-pion mixing is about one to two orders of magnitude larger than the other axion-meson mixings, and therefore dominates the axion phenomenology involving hadronic couplings.

However, notice that $|\theta_{a\pi}^{(0)}|$ in (3.7) is not bounded from below. In particular, for

$$\frac{Q_u}{Q_d} = 2 \Rightarrow \theta_{a\pi}^{(0)} \approx \frac{4Q_d}{3} \frac{f_\pi}{f_a} \left(\frac{1}{2} - \frac{m_u}{m_d} \right) \approx 0. \tag{4.1}$$

Indeed, plugging in the PDG’s 2017 weighted average for the light quark mass ratio [10], $m_u/m_d = 0.483 \pm 0.027$, and setting $Q_d = 1$ without loss of generality, (3.7) gives:

$$\theta_{a\pi}^{(0)} \approx \frac{(0.2 \pm 3) \times 10^{-3}}{f_a/\text{GeV}}, \tag{4.2}$$

which contains the *pion-phobic* range (2.4) still allowed by $\pi^+ \rightarrow e^+ \nu_e a$.

Therefore, extreme *pion-phobia* might be achievable in variant models with $Q_d/Q_u = 1/2 \approx m_u/m_d$. Unfortunately, under these circumstances, the leading order expression (3.7) is not a reliable estimate of $\theta_{a\pi}$. The reason is that the second order expansion in χ PT can give comparable contributions to $\theta_{a\pi}$, which should be included for a reliable estimate of this quantity. However, since the coefficients of many $\mathcal{O}(p^4)$ χ PT operators that contribute to $\theta_{a\pi}$ are poorly known, a precise determination of $\theta_{a\pi}$ is not possible. Moreover, as we

will discuss in section 9, additional GeV states from the UV completion of such models also give model dependent contributions to $\theta_{a\pi}$.

In summary, we have identified a particular class of axion variants, defined in (3.1) with $Q_u/Q_d = 2$, which is compatible with the condition of extreme *pion-phobia*, and hence remains a viable possibility. Due to present errors in the determination of m_u/m_d , uncertainties in the second order expansion in χ PT, and model dependence of associated UV completions, the estimated range for the axion-pion mixing (4.2) is much broader than the experimentally allowed range in (2.4), but it is nonetheless consistent with those bounds given errors.

4.1 The KTeV anomaly

Were such *pion-phobic* axion to exist, $\theta_{a\pi}$ would ideally be determined from experiment. Another very sensitive probe of this mixing angle is the π^0 decay width to e^+e^- , which is highly suppressed in the Standard Model. In fact, there appears to be a discrepancy in the observed π^0 decay width to e^+e^- at the $\sim 3\sigma$ level. The most precise measurement of this branching ratio, from the KTeV-E778 collaboration [56], is:

$$\text{Br}(\pi^0 \rightarrow e^+e^-)|_{\text{KTeV}} = (7.48 \pm 0.29 \pm 0.25) \times 10^{-8}, \quad (4.3)$$

where the first and second errors are statistical and systematic, respectively. Theoretical estimates of this branching ratio [57–59] in the SM predict:

$$\text{Br}(\pi^0 \rightarrow e^+e^-)|_{\text{SM}} = (6.23 \pm 0.09) \times 10^{-8}, \quad (4.4)$$

corresponding to a 3.2σ discrepancy between theory and experiment.⁴

In the SM, this decay is loop induced via the $\pi^0\gamma\gamma$ coupling. The imaginary (absorptive) part of this amplitude is well understood and gives an irreducible contribution, the so called “unitarity bound”, to this branching ratio [62]:

$$\text{Br}(\pi^0 \rightarrow e^+e^-)|_{\text{unit.}} = \text{Br}(\pi^0 \rightarrow \gamma\gamma) \frac{\alpha^2}{2\beta} \log^2\left(\frac{1+\beta}{1-\beta}\right) \frac{m_e^2}{m_\pi^2} \approx 4.7 \times 10^{-8}, \quad (4.5)$$

where $\beta \equiv \sqrt{1 - 4m_e^2/m_\pi^2}$. By noting that the width for $\pi^0 \rightarrow \gamma\gamma$ is given by:

$$\Gamma(\pi^0 \rightarrow \gamma\gamma) = \left(\frac{\alpha}{4\pi f_\pi}\right)^2 \frac{m_\pi^3}{4\pi}, \quad (4.6)$$

we can model the imaginary part of the SM amplitude $\text{Im}\mathcal{M}(\pi^0 \rightarrow e^+e^-)_{\text{SM}}$ via an effective $\pi^0 e^+e^-$ vertex given by [63]:

$$\mathcal{L}_{\pi ee} \supset i y_{\pi ee}^{\text{SM}} \pi^0 \bar{e} \gamma^5 e, \quad y_{\pi ee}^{\text{SM}} \equiv \frac{1}{2\beta} \log\left(\frac{1+\beta}{1-\beta}\right) \frac{\alpha^2}{2\pi} \frac{m_e}{f_\pi}. \quad (4.7)$$

⁴The KTeV-E778 collaboration extrapolated its measurement in the exclusive region $m_{ee}^2/m_\pi^2 > 0.95$ to obtain the inclusive result with final state radiation removed [56]. Refs. [60, 61] revisited the QED radiative corrections used in this extrapolation, and claimed that the discrepancy between theory and experiment could be reduced to the 2σ level.

Parameterizing BSM corrections to this coupling as $y_{\pi ee}^{\text{BSM}}$, we can finally write

$$\mathcal{L}_{\pi ee} = i (y_{\pi ee}^{\text{SM}} + y_{\pi ee}^{\text{BSM}}) \pi^0 \bar{e} \gamma^5 e, \quad (4.8)$$

and predict the contribution to the width from the imaginary part of the amplitude:

$$\Gamma(\pi^0 \rightarrow e^+ e^-) \Big|_{\text{Im}\mathcal{M}} = \beta \frac{m_\pi}{8\pi} |y_{\pi ee}^{\text{SM}} + y_{\pi ee}^{\text{BSM}}|^2. \quad (4.9)$$

As a conservative bound on the BSM contribution to $\Gamma(\pi^0 \rightarrow e^+ e^-)$, we can demand that (4.9) does not exceed the observed value (4.3) by more than two standard deviations, leading to:

$$-6.0 \times 10^{-7} \lesssim y_{\pi ee}^{\text{BSM}} \lesssim 0.83 \times 10^{-7}. \quad (4.10)$$

Note that when $y_{\pi ee}^{\text{BSM}} < 0$, the BSM amplitude destructively interferes with the SM one, and in the range $y_{\pi ee}^{\text{BSM}} \in [-5.2, 0] \times 10^{-7}$ the contribution (4.9) to the width from the imaginary part of the amplitude is smaller than the SM unitarity bound (4.5). In that case, the real (dispersive) part of the amplitude would have to be unexpectedly large to account for the observed width.

If instead we attempt to explain the excess,⁵ i.e., fit the BSM contribution to account for the discrepancy between (4.3) and (4.4), we obtain the following 2σ range:

$$y_{\pi ee}^{\text{BSM}} \Big|_{\text{KTeV}} = \begin{cases} (0.3 \pm 0.2) \times 10^{-7} & \text{or} \\ (-5.5 \pm 0.2) \times 10^{-7}. \end{cases} \quad (4.11)$$

So far we have kept this discussion general, since for some UV completions of the axion variants we are considering, not only the axion, but also other GeV states, can contribute to $y_{\pi ee}^{\text{BSM}}$. Nevertheless, we can consider the limit of these models in which the only important contribution to $y_{\pi ee}^{\text{BSM}}$ comes from the QCD axion. In this case,

$$y_{\pi ee}^{\text{BSM}} = \theta_{a\pi} \times \frac{Q_e m_e}{f_a}. \quad (4.12)$$

The 2σ bound (4.10) in this instance then becomes:

$$\theta_{a\pi} \lesssim \frac{1.6 \times 10^{-4}}{Q_e (\text{GeV}/f_a)}, \quad (4.13)$$

and the range needed to fit the KTeV anomaly within 2 standard deviations is:

$$\theta_{a\pi} \Big|_{\text{KTeV}} \approx \frac{(0.6 \pm 0.4) \times 10^{-4}}{Q_e (\text{GeV}/f_a)}. \quad (4.14)$$

Remarkably, the $\theta_{a\pi}$ fit to the KTeV anomaly, (4.14), is compatible with the *pion-phobic* range imposed by $\pi^+ \rightarrow e^+ \nu_e a$.

⁵See also [64] for an alternative BSM explanation of the KTeV anomaly.

5 Constraints from charged kaon decays

It was widely claimed in the literature [54, 55] that bounds from $K^+ \rightarrow \pi^+ a$ ruled out all QCD axion variants in the MeV mass range. In this section we critically re-examine the arguments which have led to this claim. We find that there are large uncertainties involved in obtaining $K^+ \rightarrow \pi^+ a$, from the interpretation of experimental analyses, to assumptions in estimating the $K^+ \rightarrow \pi^+(\eta^* \rightarrow a)$ amplitude, and finally to the derivation of the $\eta - a$ mixing angle. We conclude that these uncertainties are significant enough to preclude a definitive exclusion of MeV axions from existing bounds on rare K^+ decays.

5.1 Experimental bounds on $K^+ \rightarrow \pi^+(a \rightarrow e^+e^-)$

Measurements of rare K^+ decays such as $K^+ \rightarrow \pi^+\nu\bar{\nu}$, $K^+ \rightarrow \pi^+\gamma\gamma$, and $K^+ \rightarrow \pi^+e^+e^-$ have been improving over time for the past five decades or so. However, for the last 30 years, the low $m_{e^+e^-}$ region of $K^+ \rightarrow \pi^+e^+e^-$ has been neglected from experimental scrutiny for BSM physics. This is due in part to the very large backgrounds from π^0 Dalitz decays, namely, $K^+ \rightarrow \pi^+(\pi^0 \rightarrow e^+e^-\gamma)$, which make the study of the low $m_{e^+e^-}$ region very difficult. To the best of our knowledge, the last time dedicated searches were performed in this region was in the 1980's, by two different experiments, one at KEK [65] and the other at BNL [66]. We shall now discuss these searches in some detail.

The KEK experiment E89 [65] was published in 1984. It consisted of a high resolution spectrograph that measured the momentum of charged pions from the decay of charged kaons at rest. They searched for a peak in the π^+ momentum distribution, which would be evidence of a two-body decay of the charged kaon, $K^+ \rightarrow \pi^+X^0$, to a new pseudoscalar X^0 . Since they did not impose any vetoes nor requirements on the remaining decay products of K^+ , this search was sensitive to $X^0 \rightarrow$ anything. Unfortunately, we believe there are a couple of issues with this study.

First, it is not clear what was the lowest value of m_{X^0} to which this search was sensitive. The abstract states that “*bounds are presented for the mass range of X^0 from 10 to 300 MeV/c².*” The concluding paragraph, on the other hand, quotes 5 – 300 MeV as the range of exclusion for m_{X^0} . The acceptance curve for the pion momentum, shown in figure 1 of [65], has a lower range of $P_{\pi^+} \sim 224$ MeV, corresponding to a lower range of 50 MeV for m_{X^0} . Finally, the exclusion curve presented in figure 2, of $\text{Br}(K^+ \rightarrow \pi^+X^0)$ versus m_{X^0} , has a lower range of $m_{X^0} = 10$ MeV.

Moreover, there are issues with figure 2. It misrepresents limits from previous analysis, namely, Asano et al. [67] and Abrams et al. [68]. The limits from Asano et al. on $\text{Br}(K^+ \rightarrow \pi^+(X^0 \rightarrow \gamma\gamma))$ are depicted in figure 2 as a factor of ~ 5 weaker than in [67], and shown only in the range $m_{X^0} \in (50 - 90 \text{ MeV})$, when in fact Asano et al.'s limits apply from $(0 - 100 \text{ MeV})$ as long as $\tau_{X^0} < 10^{-9} \text{ s}$. It is unclear which limit from Abrams et al. [68] is being depicted in figure 2, but assuming it is the limit on $\text{Br}(K^+ \rightarrow \pi^+\gamma\gamma)$ in the region $K_{\pi^+} < 70 \text{ MeV}$, corresponding to $P_{\pi^+} > 156 \text{ MeV}$ ($M_{\gamma\gamma} > 237 \text{ MeV}$), then figure 2 of [65] misrepresents it by a factor of ~ 5 as well.

An earlier conference note [69] presenting preliminary results of the KEK E89 analysis might shed light on the origin of these discrepancies.⁶ In figure 6 of conference note [69], limits were presented in terms of the relative branching ratio $\text{Br}(K^+ \rightarrow \pi^+ X^0)/\text{Br}(K^+ \rightarrow \pi^+ \pi^0) \approx 5 \times \text{Br}(K^+ \rightarrow \pi^+ X^0)$, and represented more accurately the results from Asano et al. [67] and Abrams et al. [68]. This suggests that the factor of ~ 5 discrepancy in the final publication [65] was a typo in recasting the plot from relative branching ratio to absolute branching ratio. Adding to the ambiguity on the lowest X^0 mass to which KEK E89 was sensitive, the limits presented on figure 6 of the conference note [69] did not extend below $m_{X^0} \sim 50$ MeV.

The second issue with the KEK E89 analysis was their statistical inference of exclusion limits in the low statistics region $m_{X^0} \lesssim 80$ MeV. The formula used for obtaining limits is appropriate in the case of a “bump-hunt” on top of a very large background, where the errors are gaussian and scale as $\sqrt{N_{\text{background}}}$. It fails, however, in the Poisson limit where the number of expected events from signal plus background is $\mathcal{O}(1)$, where it can overestimate the exclusion power by a factor of several.

All things considered, we conservatively choose to ignore the limits from KEK E89 analysis [65] in the region $m_{X^0} \lesssim 50$ MeV.

That leaves us with the other dedicated search for the rare decay $K^+ \rightarrow \pi^+(a \rightarrow e^+e^-)$ in the low $m_{e^+e^-}$ region, published in 1987 by Baker et al. [66]. Their apparatus, located at BNL’s AGS, was optimized for the decay of K^+ to three charged tracks, with efficient discrimination between electrons, pions and muons. They collected a large sample of approximately 2.8×10^4 $K^+ \rightarrow \pi^+e^+e^-$ events, mostly Dalitz decays, $K^+ \rightarrow \pi^+(\pi^0 \rightarrow e^+e^-\gamma)$, and applied optimized cuts to remove this background while remaining sensitive to the $K^+ \rightarrow \pi^+a$ signal. While they claim that their results exclude branching ratios $\text{Br}(K^+ \rightarrow \pi^+a) \gtrsim 0.8 \times 10^{-7}$ for any mass in the range $m_a \in (1.8 - 100$ MeV) at 90% confidence level, we believe their exclusion is too aggressive for two reasons.

Firstly, in their final signal region, they fit the observed $m_{e^+e^-}$ distribution in the range 1.8 – 100 MeV as a constant background not properly modeled by their Monte Carlo (MC), and then subtract it before extracting limits. We believe that a more conservative approach when deriving limits would be to treat any residual events in the signal region as potential signal. We can estimate what those more conservative limits would be based on the data shown in figure 2(d) of [66], together with the assumption that the signal efficiency as a function of m_a does not vary substantially from the one quoted at $m_a = 1.8$ MeV, and trusting their Monte Carlo signal yield of 5 expected events for $\text{Br}(K^+ \rightarrow \pi^+a) = 10^{-6}$. Under these assumptions, we infer the limit $\text{Br}(K^+ \rightarrow \pi^+a) \lesssim 2.4 \times 10^{-6}$ in the range $m_a \sim (10 - 25)$ MeV.

Our second concern regards this analysis’ sole reliance on Monte Carlo to estimate all the acceptances of both background and signal. As figure 2(d) of [66] shows, their Monte Carlo grossly misestimates the Dalitz background contamination in the signal region by about one order of magnitude. They do not address this mismodeling and simply re-scale

⁶We also attempted to contact some of the authors of [65], but since three decades have elapsed since the time of publication of this analysis, we understandably did not get a response to our queries.

their expected background to match the observed rate. More importantly, they do not address whether their Monte Carlo estimation of the *signal acceptance* could possibly be mismodeled as well. Were their signal acceptance off by a similar order of magnitude, their limits could be weakened to as much as $\text{Br}(K^+ \rightarrow \pi^+ a) \lesssim 10^{-5}$ in the low $m_{e^+e^-}$ region.

We therefore choose to remain agnostic about the precise experimental limit on MeV axions from kaon decays, and for the remainder of this paper we discuss implications from K^+ decays to axion-meson mixing parameters assuming the following range:

$$\text{Br}(K^+ \rightarrow \pi^+ a \rightarrow \pi^+ e^+ e^-) \lesssim 10^{-6} - 10^{-5}. \quad (5.1)$$

We stress that new experimental studies of the low m_{ee} kinematic region in $K^+ \rightarrow \pi^+ e^+ e^-$ are relevant and warranted, considering the sensitivity of this final state to rare new physics processes, and the fact that measurements in this region could be greatly improved by modern experiments.

5.2 Estimation of the amplitude for $K^+ \rightarrow \pi^+ a$

Next, we discuss how to relate the amplitude for $K^+ \rightarrow \pi^+ a$ to the SM amplitudes for $K^+ \rightarrow \pi^+ \pi^0$, $K^+ \rightarrow \pi^+ \eta_8$, and $K^+ \rightarrow \pi^+ \eta_0$, assuming that the axion-meson mixing angles $\theta_{a\pi}$, $\theta_{a\eta_8}$, and $\theta_{a\eta_0}$ are known.

Since the π^0 is on-shell in the SM process $K^+ \rightarrow \pi^+ \pi^0$, it is trivial to relate the amplitude for this decay to the amplitude for $K^+ \rightarrow \pi^+ a$ via axion-pion mixing:

$$\mathcal{M}(K^+ \rightarrow \pi^+ a)|_{a-\pi \text{ mixing}} = \theta_{a\pi} \mathcal{M}(K^+ \rightarrow \pi^+ \pi^0). \quad (5.2)$$

If (5.2) were the dominant contribution to $\mathcal{M}(K^+ \rightarrow \pi^+ a)$, we could plug in the upper bound (2.4) on $\theta_{a\pi}$ to obtain:

$$\text{Br}(K^+ \rightarrow \pi^+ a)|_{a-\pi \text{ mixing}} \lesssim (0.5 \times 10^{-8}) \times \text{Br}(K^+ \rightarrow \pi^+ \pi^0) \approx 10^{-9}, \quad (5.3)$$

which is at least three orders of magnitude below the existing bounds discussed in section 5.1.

Antoniadis et al. [54] and Bardeen et al. [55] pointed out that the axion mixing with η and η' also contribute to $\mathcal{M}(K^+ \rightarrow \pi^+ a)$:

$$\mathcal{M}(K^+ \rightarrow \pi^+ a)|_{a-\eta_{8,0} \text{ mixing}} = \theta_{a\eta_8} \mathcal{M}(K^+ \rightarrow \pi^+ \eta_8) + \theta_{a\eta_0} \mathcal{M}(K^+ \rightarrow \pi^+ \eta_0), \quad (5.4)$$

and that, due to $\Delta I = 1/2$ *enhancement* (a.k.a. *octet enhancement*) of these amplitudes, these contributions dominate the branching ratio for $K^+ \rightarrow \pi^+ a$ and lead to tension with experimental constraints.

Let us briefly review the arguments of [55]. In order to obtain the amplitudes $\mathcal{M}(K^+ \rightarrow \pi^+ \eta_{8,0})$, [55] considers the leading order $\Delta S = 1$ chiral Lagrangian describing non-leptonic kaon decays:

$$\mathcal{L}_\chi^{\Delta S=1} \Big|_{\mathcal{O}(p^2)} = g_8 f_\pi^2 \text{Tr}(\lambda_{ds} \partial_\mu U \partial^\mu U^\dagger) + g_{27} f_\pi^2 C_{ab} \text{Tr}(\lambda_a \partial_\mu U U^\dagger \lambda_b \partial^\mu U U^\dagger) + \text{h.c.}, \quad (5.5)$$

where λ_a are the Gell-Mann matrices, $\lambda_{ds} \equiv (\lambda_6 + i\lambda_7)/2$, and the Clebsch-Gordan coefficients C_{ab} are given for instance in [70]. The first operator transforms as an octet under chiral $SU(3)_\chi$, while the second transforms as a 27-plet. In particular, while $\mathcal{M}(K^0 \rightarrow \pi\pi)$ receives contributions from both operators, only the 27-plet operator contributes to $\mathcal{M}(K^+ \rightarrow \pi^+\pi^0)$:

$$\mathcal{M}(K^0 \rightarrow \pi^+\pi^-) = \frac{\sqrt{2}}{f_\pi}(m_K^2 - m_\pi^2) \left(g_8 e^{i\delta_0} + g_{27} e^{i\delta_2} \right), \quad (5.6)$$

$$\mathcal{M}(K^0 \rightarrow \pi^0\pi^0) = \frac{\sqrt{2}}{f_\pi}(m_K^2 - m_\pi^2) \left(g_8 e^{i\delta_0} - 2g_{27} e^{i\delta_2} \right), \quad (5.7)$$

$$\mathcal{M}(K^+ \rightarrow \pi^+\pi^0) = \frac{3}{f_\pi}(m_K^2 - m_\pi^2) g_{27} e^{i\delta_2}. \quad (5.8)$$

Above, δ_0, δ_2 are strong interaction S-wave $\pi\pi$ phase shifts. Within the effective framework of χ PT, the coefficients g_8 and g_{27} cannot be obtained from first principles. However, under the standard assumption that (5.6), (5.7), (5.8) provide the leading contributions non-leptonic kaon decays, g_8 and g_{27} can be fit to match the observed $K \rightarrow \pi\pi$ amplitudes [70] (see also [71]):

$$g_8 \simeq 7.8 \times 10^{-8}, \quad g_{27} \simeq 0.25 \times 10^{-8}. \quad (5.9)$$

Whereas naïvely one would expect $g_8 \sim g_{27}$, (5.9) shows a large enhancement of the octet coefficient relative to the 27-plet's, $g_8/g_{27} \simeq 31.2$, tied to the fact that the hadronic width of K_S^0 is much broader than that of K^\pm , e.g.,

$$\frac{\Gamma(K_S^0 \rightarrow \pi\pi)}{\Gamma(K^\pm \rightarrow \pi^+\pi^0)} \approx 668. \quad (5.10)$$

This is known as the *octet enhancement* in non-leptonic kaon decays.

Ref. [55] noted that the octet operator in (5.5) also contributes to the (off-shell) amplitudes:

$$\mathcal{M}(K^+ \rightarrow \pi^+\eta_8) = \frac{1}{\sqrt{3}f_\pi}(2m_K^2 + m_\pi^2 - 3p_{\eta_8}^2) g_8, \quad (5.11)$$

$$\mathcal{M}(K^+ \rightarrow \pi^+\eta_0) = \frac{2\sqrt{2}}{\sqrt{3}f_\pi}(m_K^2 - m_\pi^2) g_8. \quad (5.12)$$

Defining the quark flavor basis:

$$\eta_{ud} \equiv \frac{\eta_8}{\sqrt{3}} + \frac{\eta_0}{\sqrt{3/2}}, \quad \eta_s \equiv -\frac{\eta_8}{\sqrt{3/2}} + \frac{\eta_0}{\sqrt{3}}, \quad (5.13)$$

and neglecting terms $\mathcal{O}(g_{27}/g_8)$ and $\mathcal{O}(m_\pi^2/m_K^2)$, we can then use (5.11), (5.12), (5.4) and (5.6) to obtain the following relation:⁷

$$\mathcal{M}(K^+ \rightarrow \pi^+a) \Big|_{\text{octet enh.}} \approx \theta_{a\eta_{ud}} \sqrt{2} \mathcal{M}(K^0 \rightarrow \pi^+\pi^-). \quad (5.17)$$

⁷To be precise, the definition of axion mixing angles with states that are not mass eigenstates, such as

Ref. [54] noted that expression (5.17) does not take into account the absence of strong final-state interactions between π^+ and a . Following [54], we correct this by introducing a fudge factor $D_{\pi\pi} \sim 1/\sqrt{3}$ multiplying the r.h.s. of (5.17). We then finally obtain:

$$\begin{aligned} \text{Br}(K^+ \rightarrow \pi^+ a) \Big|_{\text{octet enh.}} &\approx \theta_{a\eta_{ud}}^2 2 \frac{\Gamma_{K_s^0}}{\Gamma_{K^+}} \text{Br}(K_s^0 \rightarrow \pi^+ \pi^-) \frac{|\vec{p}_a|}{|\vec{p}_\pi|} D_{\pi\pi}^2 \\ &\approx 64 \theta_{a\eta_{ud}}^2. \end{aligned} \quad (5.18)$$

Finally, using the experimental upper bound (5.1), we infer the following constraint:

$$|\theta_{a\eta_{ud}}| \Big|_{\text{octet enh.}} \lesssim (1-4) \times 10^{-4}. \quad (5.19)$$

We would like to stress that the arguments above, leading to (5.18) and (5.19), hinge on the assumption that the $\mathcal{O}(p^2)$ octet operator, shown in (5.5), is enhanced. While this assumption is not considered controversial, it is nonetheless possible that this is not the correct description of octet enhancement in non-leptonic kaon decays [72–74]. In particular, an equally good description is obtained by transferring the enhancement to the $\mathcal{O}(p^4)$ expansion of $\mathcal{L}_\chi^{\Delta S=1}$. Consider, for instance,

$$\mathcal{L}_\chi^{\Delta S=1} \Big|_{\mathcal{O}(p^4)} \supset g'_8 \frac{f_\pi^2}{\Lambda^2} B \text{Tr}(\lambda_{ds} M_q U) \text{Tr}(\partial_\mu U \partial^\mu U^\dagger) + \text{h.c.}, \quad (5.20)$$

where $\Lambda \sim 4\pi f_\pi$ is a natural cut-off. If one assumes that the $\mathcal{O}(p^2)$ coefficients g_8 and g_{27} are comparable, and instead the $\mathcal{O}(p^4)$ coefficient g'_8 is enhanced,

$$\frac{g_8}{g_{27}} \sim \mathcal{O}(1), \quad \frac{g'_8}{g_{27}} \sim \mathcal{O}(100), \quad g'_8 \simeq \frac{\Lambda^2}{2m_K^2} g_8 \Big|_{(5.9)} \approx 1.6 \times 10^{-7}, \quad (5.21)$$

one obtains an equally good phenomenological fit of $K \rightarrow \pi\pi$ and $K \rightarrow \pi\pi\pi$ data compared to the standard fit in (5.9).⁸

However, unlike the octet operator in (5.5), (5.20) *does not contribute* to $K^+ \rightarrow \pi^+ \eta_{8,0}$. Therefore, the implication of the alternative fit in (5.21) would be that the amplitude for $K^+ \rightarrow \pi^+ a$, which is proportional to g_8 , but unaffected by g'_8 , would not be octet enhanced. In that case, one would have to properly redo the fit to kaon data to obtain the precise

η_{ud} and η_s , goes as follows. Consider the interactions in the (canonically normalized) quark flavor basis:

$$V \supset \frac{\mu_q^2}{2} \eta_{ud}^2 + \frac{\mu_s^2}{2} \eta_s^2 + \mu_{qs}^2 \eta_{ud} \eta_s + \mu_{aq}^2 a \eta_{ud} + \mu_{as}^2 a \eta_s + \mathcal{J}_{ud} \eta_{ud} + \mathcal{J}_s \eta_s, \quad (5.14)$$

where \mathcal{J}_{ud} and \mathcal{J}_s are external sources. Integrating η_{ud} and η_s out, one obtains:

$$V \supset \theta_{a\eta_{ud}} a \mathcal{J}_{ud} + \theta_{a\eta_s} a \mathcal{J}_s, \quad (5.15)$$

with

$$\theta_{a\eta_{ud}} = -\frac{\mu_{aq}^2 \mu_s^2 - \mu_{qs}^2 \mu_{as}^2}{\mu_q^2 \mu_s^2 - \mu_{qs}^4}, \quad \theta_{a\eta_s} = -\frac{\mu_{as}^2 \mu_q^2 - \mu_{qs}^2 \mu_{aq}^2}{\mu_q^2 \mu_s^2 - \mu_{qs}^4}. \quad (5.16)$$

⁸Specifically, there is a degeneracy in the $\{g_8, g'_8\}$ dependence of the amplitudes for $K \rightarrow \pi\pi$ and $K \rightarrow \pi\pi\pi$, which can only be broken with additional assumptions. See [75, 76].

value of g_8 , but as a rough estimate, taking $g_8 \sim \mathcal{O}(g_{27})$, the constraint on $a - \eta$ mixing would be weakened by a factor of ~ 30 ,

$$|\theta_{a\eta_{ud}}| \Big|_{\text{not octet enh.}} \lesssim (0.4 - 1) \times 10^{-2}. \tag{5.22}$$

In short, the enhancement of the amplitude $\mathcal{M}(K^+ \rightarrow \pi^+ a)$ depends on how octet enhancement is realized in χ PT. If the amplitudes $\mathcal{M}(K^+ \rightarrow \pi^+ \eta_{8,0})$ are not octet enhanced, the resulting bound (5.22) on $\theta_{a\eta_{ud}}$ is relatively weak and does not presently pose a threat to the viability of MeV axions. Under the more conventional assumption of octet enhancement at $\mathcal{O}(p^2)$, however, it becomes important to establish how compatible is $\theta_{a\eta_{ud}}$ with the bound in (5.19), which we shall now address.

5.3 Axion-eta mixing

The last sources of uncertainty in predicting the rare decay $K^+ \rightarrow \pi^+ a$, namely, the mixing angles $\theta_{a\eta_8}$ and $\theta_{a\eta_0}$, have been the least critically examined in the axion literature.

It is well known that the leading order expansion in χ PT does not adequately describe the η and η' masses and mixing angles [77, 78]. Indeed, the second order expansion of the Chiral Lagrangian provides important corrections to masses, decay constants and mixing angles of singlet and octet mesons, which typically scale as:

$$\frac{32 m_K^2}{f_\pi^2} L_i \sim \mathcal{O}(10^3) L_i. \tag{5.23}$$

Above, L_i are the dimensionless coefficients of the $\mathcal{O}(p^4)$ operators in the chiral expansion, and are commonly known as *Low Energy Constants* (LECs) [79, 80]. Many LECs are reasonably well-determined from experimental and/or lattice data, their typical size being $L_i \sim \mathcal{O}(10^{-3})$. From (5.23) it is then evident that these encode $\mathcal{O}(1)$ effects in η - η' mixing, and may very well have comparable importance in describing a - η and a - η' mixing.

In order to illustrate the uncertainties involved in obtaining $\theta_{a\eta,\eta'}$, we consider Leutwyler's study of η - η' mixing in [81], which, based on $1/N_c$ -expansion counting rules, retained only the following operators:⁹

$$\begin{aligned} \mathcal{L}_\chi^{\eta-\eta'} = & \frac{F^2}{4} \text{Tr} \left[\partial_\mu U \partial^\mu U^\dagger \right] + \frac{F^2}{4} \text{Tr} \left[(2BM_q) U + \text{h.c.} \right] - \frac{1}{2} M_0^2 \eta_0^2 \\ & + L_5 \text{Tr} \left[\partial^\mu (2BM_q U) \partial_\mu U^\dagger U + \text{h.c.} \right] + L_8 \text{Tr} \left[(2BM_q) U (2BM_q) U + \text{h.c.} \right] \\ & + \text{OZI violating terms.} \end{aligned} \tag{5.24}$$

Loop corrections do not contribute at this order, and [81] obtains $F = 90.6 \text{ MeV}$, $L_5 = 2.2 \times 10^{-3}$, $L_8 = 1.0 \times 10^{-3}$, and $M_0 \simeq 1030 \text{ MeV}$.

⁹For simplicity, in this exercise we omit the OZI violating terms in (5.24). Their numerical values obtained in the fit of [81] change our results by $\mathcal{O}(10\%)$.

Remembering that the axion is formally a phase of the light quark mass matrix M_q (see (3.3)), and setting $Q_d = Q_u/2 = 1$, we can expand (5.24) to obtain:

$$\begin{aligned} \mathcal{L}_\chi^{\eta-\eta'} \supset & \frac{1}{2} \partial_\mu \eta_{ud} \partial^\mu \eta_{ud} + \frac{1}{2} \left(1 + \frac{\hat{L}_5}{2}\right) \partial_\mu \eta_s \partial^\mu \eta_s + \frac{1}{2} \partial_\mu a \partial^\mu a \\ & + \frac{1}{2} 2(1 + \hat{L}_8) m_K^2 \eta_s^2 + \frac{1}{2} M_0^2 \left(\frac{\eta_s}{\sqrt{3}} + \frac{\eta_{ud}}{\sqrt{3/2}}\right)^2 + \frac{1}{2} m_a^2 a^2 \\ & + \frac{\hat{L}_5}{6} \frac{m_\pi^2}{m_K^2} \frac{F}{f_a} \partial_\mu \eta_{ud} \partial^\mu a + \frac{4 m_\pi^2}{3} \frac{F}{f_a} \eta_{ud} a. \end{aligned}$$

Above, we have expanded the coefficients in powers of $\frac{m_\pi^2}{m_K^2}$ and kept only the leading terms; we have also omitted all π and K dependent terms, since they are not relevant for the present discussion. Furthermore, we have simplified the notation by defining:

$$\hat{L}_i \equiv \frac{32 m_K^2}{F^2} L_i. \quad (5.25)$$

Canonically normalizing the kinetic terms and integrating out η_{ud} and η_s , we obtain:

$$\theta_{a\eta_{ud}} = - \left(\frac{1 - \hat{L}_5(1 + \hat{L}_8)/2}{3(1 + \hat{L}_8)} + 2 \frac{m_K^2}{M_0^2} \right) \frac{m_\pi^2}{m_K^2} \frac{F}{f_a}. \quad (5.26)$$

Plugging in the numerical fit of the LECs obtained by [81] yields:

$$\theta_{a\eta_{ud}} \approx \frac{-1.9 \times 10^{-3}}{f_a/\text{GeV}} \approx 0.2 \times \theta_{a\eta_{ud}}^{(0)}. \quad (5.27)$$

Rather than claiming that we have calculated $\theta_{a\eta_{ud}}$ more precisely, our purpose with this exercise is to make the point that contributions from the second order chiral expansion change the leading order estimate, $\theta_{a\eta_{ud}}^{(0)}$, by $\mathcal{O}(1)$. While we have restricted this exercise to a couple of LECs at tree level, inclusion of the full $\mathcal{O}(p^4)$ expansion and loop corrections will most likely change the answer from (5.27). Had we derived such expression, even then we would not be able to evaluate it numerically because many of the LECs contributing to $\theta_{a\eta_{ud}}$ are still undetermined. For instance, operators such as [82]

$$\begin{aligned} \mathcal{L}_\chi^{\mathcal{O}(p^4)} \supset & L_7 \text{Tr} \left[(2BM_q) U - U^\dagger (2BM_q)^\dagger \right]^2 + \lambda_2 F^2 \frac{\eta_0}{F} \text{Tr} \left[(2BM_q) U - U^\dagger (2BM_q)^\dagger \right] \\ & + L_{18} \text{Tr} \left[U^\dagger \partial^\mu U \right] \text{Tr} \left[\partial_\mu (U^\dagger (2BM_q)^\dagger - (2BM_q) U) \right] \\ & + i L_{25} \text{Tr} \left[U^\dagger (2BM_q)^\dagger U^\dagger (2BM_q)^\dagger - (2BM_q) U (2BM_q) U \right] \\ & + i L_{26} \left(\text{Tr} \left[U^\dagger (2BM_q)^\dagger \right]^2 - \text{Tr} \left[(2BM_q) U \right]^2 \right) + \dots \end{aligned} \quad (5.28)$$

also give important contributions $\theta_{a\eta_{ud}}$ and $\theta_{a\eta_s}$. While such operators also affect $\eta-\eta'$ mixing, and hence are subject to constraints, there are too many independent LECs to be fit by observations, and without further assumptions, the parameterization of the $\eta-\eta'$ phenomenology in χ PT is underconstrained.

We conclude, therefore, that currently it is impossible to reliably estimate $\theta_{a\eta_{ud}}$ and $\theta_{a\eta_s}$, given that these mixing angles can receive $\mathcal{O}(1)$ corrections from the second order chiral expansion whose numerical values are undetermined by existing data. Therefore, one cannot claim with confidence that $\theta_{a\eta_{ud}}$ violates the $K^+ \rightarrow \pi^+ a$ bounds in (5.19) or (5.22), and therefore K^+ decay bounds by themselves do not provide a *definitive* exclusion of the QCD axion variant we are considering. Most likely, a reliable determination of the mixing angles $\theta_{a\eta_{ud}}$ and $\theta_{a\eta_s}$ would have to come from lattice calculations, or, were such axion to be realized in nature, from direct measurements.

6 The physical axion current

Because of much confusion in the literature, in this section we review the differences between the PQ current, the Bardeen-Tye current, and the current associated with the axion mass eigenstate. Identifying the latter, in particular, is a critical part of working out the proper phenomenology of axions, since it is the physical axion current that determines the coupling of the axion to the chiral $U(3)_\chi$ currents that mediate meson decays, nuclear de-excitations, as well as the electromagnetic anomaly that induces the axion coupling to two photons.

In the UV, the PQ current for the class of axion variants defined in (3.1) is given by:¹⁰

$$J_\mu^{\text{PQ}} = f_a \partial_\mu a - \frac{Q_u}{2} \bar{u} \gamma_\mu \gamma_5 u - \frac{Q_d}{2} \bar{d} \gamma_\mu \gamma_5 d. \quad (6.1)$$

Bardeen and Tye [83] pointed out that the PQ current cannot be associated with the physical axion, i.e., the light mass eigenstate, since the divergence of this current receives a large contribution from the strong anomaly:

$$\partial^\mu J_\mu^{\text{PQ}} = -\frac{\alpha_s}{8\pi} (Q_u + Q_d) G_{\mu\nu} \tilde{G}^{\mu\nu}, \quad (6.2)$$

which does not vanish in the limit $m_q \rightarrow 0$. The physical axion, on the other hand, becomes a *bona fide* Goldstone boson in the limit of a massless quark, and should therefore be associated with a current that is conserved as $m_q \rightarrow 0$.

In order to identify the physical current, [83] subtracted the strong anomaly from J^{PQ} , obtaining the *Bardeen-Tye* current:

$$J_\mu^{\text{BT}} \equiv J_\mu^{\text{PQ}} + \frac{(Q_u + Q_d)/2}{(m_u^{-1} + m_d^{-1} + m_s^{-1})} (m_u^{-1} \bar{u} \gamma_\mu \gamma_5 u + m_d^{-1} \bar{d} \gamma_\mu \gamma_5 d + m_s^{-1} \bar{s} \gamma_\mu \gamma_5 s), \quad (6.3)$$

which is conserved in the limit $m_q \rightarrow 0$:

$$\partial^\mu J_\mu^{\text{BT}} = (Q_u + Q_d) \frac{m_u m_d m_s}{m_d m_s + m_u m_s + m_u m_d} (\bar{u} i \gamma_5 u + \bar{d} i \gamma_5 d + \bar{s} i \gamma_5 s). \quad (6.4)$$

¹⁰Since this discussion is concerned with the hadronic properties of the axion, we ignore the leptonic current and electromagnetic anomaly for the remainder of this subsection.

We can rewrite J^{BT} in order to show its explicit dependence on the chiral $U(3)_\chi$ currents. First we define:

$$J_{5\mu}^{(3)} = \frac{\bar{u}\gamma_\mu\gamma_5 u - \bar{d}\gamma_\mu\gamma_5 d}{2} \equiv f_\pi \partial_\mu \pi_3, \quad (6.5a)$$

$$J_{5\mu}^{(8)} = \frac{\bar{u}\gamma_\mu\gamma_5 u + \bar{d}\gamma_\mu\gamma_5 d - 2\bar{s}\gamma_\mu\gamma_5 s}{2\sqrt{3}} \equiv f_\pi \partial_\mu \eta_8, \quad (6.5b)$$

$$J_{5\mu}^{(0)} = \frac{\bar{u}\gamma_\mu\gamma_5 u + \bar{d}\gamma_\mu\gamma_5 d + \bar{s}\gamma_\mu\gamma_5 s}{\sqrt{6}} \equiv f_\pi \partial_\mu \eta_0. \quad (6.5c)$$

Above, π_3 , η_8 and η_0 are *not* mass eigenstates.

The Bardeen-Tye current in (6.3) can then be written as:

$$J_\mu^{\text{BT}} = f_a \partial_\mu a - \left(\frac{(Q_u m_u - Q_d m_d)}{(1 + \epsilon_\eta)(m_u + m_d)} + \epsilon_\eta \frac{(Q_u - Q_d)}{2(1 + \epsilon_\eta)} \right) J_{5\mu}^{(3)} - \left(\frac{\sqrt{3}}{2} \frac{\epsilon_\eta}{(1 + \epsilon_\eta)} (Q_u + Q_d) \right) J_{5\mu}^{(8)}, \quad (6.6)$$

$$\epsilon_\eta \equiv \frac{m_u m_d}{m_s(m_u + m_d)} \approx \frac{m_u m_d}{(m_u + m_d)^2} \frac{m_\pi^2}{m_K^2}. \quad (6.7)$$

The Bardeen-Tye current has been identified in numerous studies as the physical axion current. Under this assumption, it is straightforward to read the mixing angles $\theta_{a\pi/\eta}$ from (6.5) and (6.6):

$$\theta_{a\pi}^{\text{BT}} = -\frac{1}{(1 + \epsilon_\eta)} \left(\frac{(Q_u m_u - Q_d m_d)}{(m_u + m_d)} + \epsilon_\eta \frac{(Q_u - Q_d)}{2} \right) \frac{f_\pi}{f_a}, \quad (6.8)$$

$$\theta_{a\eta_8}^{\text{BT}} = -\sqrt{\frac{3}{2}} \frac{\epsilon_\eta}{(1 + \epsilon_\eta)} (Q_u + Q_d) \frac{f_\pi}{f_a}, \quad (6.9)$$

$$\theta_{a\eta_0}^{\text{BT}} = 0. \quad (6.10)$$

Note, however, that this assumption is only strictly correct in the limit where the η' is decoupled, i.e., in the limit $M_0 \rightarrow \infty$ (see (3.2)) and $\eta' \rightarrow \eta_0$. Indeed, the Bardeen-Tye prescription to obtain the mixing angles $\theta_{a\pi/\eta}$ is *equivalent* to the 1st order χ PT prescription with η_0 decoupled. This is corroborated by the fact that (3.7), (3.8), (3.9) and (6.8), (6.9), (6.10) agree in the limit $M_0 \rightarrow \infty$.

We point out as a side remark that the condition that $\partial^\mu J_\mu^{\text{BT}}$ vanishes in the limit $m_q \rightarrow 0$ is not sufficient to uniquely determine the Bardeen-Tye current. In particular, one could define:

$$J_\mu^{\prime\text{BT}} = J_\mu^{\text{BT}} + \frac{m_u m_d m_s}{(m_u + m_d)(m_u + m_s)(m_d + m_s)} (c_3 J_{5\mu}^{(3)} + c_8 J_{5\mu}^{(8)} + c_0 J_{5\mu}^{(0)}), \quad (6.11)$$

which also satisfies the aforementioned condition with arbitrary finite coefficients $c_{3,8,0}$. Again, in this context, this ambiguity can only be removed in the limit of decoupled η' . In this limit, c_0 in (6.11) must necessarily vanish, since $\theta_{a\eta_0} \propto c_0 \rightarrow 0$ as $m_{\eta'} \rightarrow \infty$.¹¹

¹¹The vanishing of $\theta_{a\eta_0}$ as $M_0 \rightarrow \infty$ only holds for bases where the coupling $(a/f_a)G\tilde{G}$ has been rotated away, which is the case of (3.1). However, the statement that the Bardeen-Tye prescription is equivalent to taking the limit $M_0 \rightarrow \infty$ in 1st order χ PT is basis-independent.

Moreover, in this limit the mass eigenstates a_{phys} , π and η are entirely determined by the non-anomalous currents J'^{BT} , $J^{(3)}$ and $J^{(8)}$. Imposing then that J'^{BT} commutes with J^π and J^η , one can estimate c_3 and c_8 via current algebra methods, and conclude that these coefficients provide only small corrections to (6.8) and (6.9) of order m_a^2/m_π^2 and m_a^2/m_η^2 , respectively.

Unfortunately, once the η' is included in the spectrum, the Bardeen-Tye prescription is no longer sufficient to determine the physical axion current. In particular, while π_3 in (6.5a) is a good approximation to the mass eigenstate π^0 , the physical states η and η' have substantial components of both η_8 and η_0 , and hence are not amenable to current algebra methods, which are not applicable to anomalous currents.

Therefore, generically the physical axion current has components from all three neutral chiral $U(3)_\chi$ currents:

$$J_\mu^{a_{\text{phys}}} \equiv f_a \partial_\mu a_{\text{phys}} = f_a \partial_\mu a + \xi_{a3} J_{5\mu}^{(3)} + \xi_{a8} J_{5\mu}^{(8)} + \xi_{a0} J_{5\mu}^{(0)}. \quad (6.12)$$

The coefficients ξ_{a3} , ξ_{a8} and ξ_{a0} in (6.12) are related to the axion-meson mixing angles by:

$$\theta_{a\pi} = \xi_{a3} \frac{f_\pi}{f_a}, \quad \theta_{a\eta_8} = \xi_{a8} \frac{f_\pi}{f_a}, \quad \theta_{a\eta_0} = \xi_{a0} \frac{f_\pi}{f_a}. \quad (6.13)$$

This discussion provides yet another perspective on the difficulties in estimating the hadronic mixing angles of the axion. This propagates into uncertainties in the axion's phenomenology, such as rates of rare meson decays, discussed in the previous two sections, as well as the axion couplings to photons and nucleons, which we will now consider.

6.1 Axion-photon coupling

As light quarks confine below the QCD scale, an effective operator coupling the axion to the electromagnetic dual field strength is generated. In this section we derive this operator, estimate the axion decay branching ratio to a pair of photons, and comment on bounds from $K^+ \rightarrow \pi^+(a \rightarrow \gamma\gamma)$.

We start by rewriting the physical axion current (6.12) as:

$$J_\mu^{a_{\text{phys}}} = f_a \partial_\mu a + \xi_u \bar{u} \gamma_\mu \gamma_5 u + \xi_d \bar{d} \gamma_\mu \gamma_5 d + \xi_s \bar{s} \gamma_\mu \gamma_5 s, \quad (6.14)$$

where

$$\xi_u = \frac{f_a}{f_\pi} \left(\frac{\theta_{a\pi}}{2} + \frac{\theta_{a\eta_8}}{2\sqrt{3}} + \frac{\theta_{a\eta_0}}{\sqrt{6}} \right) = \frac{f_a}{f_\pi} \left(\frac{\theta_{a\pi}}{2} + \frac{\theta_{a\eta_{ud}}}{2} \right), \quad (6.15)$$

$$\xi_d = \frac{f_a}{f_\pi} \left(-\frac{\theta_{a\pi}}{2} + \frac{\theta_{a\eta_8}}{2\sqrt{3}} + \frac{\theta_{a\eta_0}}{\sqrt{6}} \right) = \frac{f_a}{f_\pi} \left(-\frac{\theta_{a\pi}}{2} + \frac{\theta_{a\eta_{ud}}}{2} \right) \quad (6.16)$$

$$\xi_s = \frac{f_a}{f_\pi} \left(-\frac{\theta_{a\eta_8}}{\sqrt{3}} + \frac{\theta_{a\eta_0}}{\sqrt{6}} \right) = \frac{f_a}{f_\pi} \left(\frac{\theta_{a\eta_s}}{\sqrt{2}} \right). \quad (6.17)$$

With this notation, we can derive the electromagnetic anomaly of $J_\mu^{a\text{phys}}$ and the axion coupling to photons straightforwardly:

$$\begin{aligned}
 \mathcal{L}_a &\supset \frac{\alpha}{2\pi f_a} 3 \left(\xi_u \left(\frac{2}{3}\right)^2 + \xi_d \left(-\frac{1}{3}\right)^2 + \xi_s \left(-\frac{1}{3}\right)^2 \right) a F_{\mu\nu} \tilde{F}^{\mu\nu} \\
 &= \frac{\alpha}{4\pi f_\pi} \left(\theta_{a\pi} + \frac{\theta_{a\eta_8}}{\sqrt{3}} + 2 \frac{\theta_{a\eta_0}}{\sqrt{3/2}} \right) a F_{\mu\nu} \tilde{F}^{\mu\nu} \\
 &= \frac{\alpha}{4\pi f_\pi} \left(\theta_{a\pi} + \frac{5}{3} \theta_{a\eta_{ud}} + \frac{\sqrt{2}}{3} \theta_{a\eta_s} \right) a F_{\mu\nu} \tilde{F}^{\mu\nu}, \tag{6.18}
 \end{aligned}$$

where we assume that $m_a > 2m_e$, so that the coupling of the axion to electrons does not contribute to this effective operator.

The axion decay width to $\gamma\gamma$ is then given by:

$$\Gamma(a \rightarrow \gamma\gamma) = \left(\theta_{a\pi} + \frac{5}{3} \theta_{a\eta_{ud}} + \frac{\sqrt{2}}{3} \theta_{a\eta_s} \right)^2 \left(\frac{\alpha}{4\pi f_\pi} \right)^2 \frac{m_a^3}{4\pi}. \tag{6.19}$$

This expression neglects the loop induced contribution coming from the axion coupling to electrons, which becomes comparable to the hadronic contribution above if the linear combination of mixing angles in (6.19) is $\mathcal{O}(10^{-4})$.

The relative $\gamma\gamma$ to e^+e^- branching ratio is then:

$$\frac{\text{Br}(a \rightarrow \gamma\gamma)}{\text{Br}(a \rightarrow e^+e^-)} = \frac{2}{\beta_e} \frac{m_u m_d}{(m_u + m_d)^2} \left(\frac{Q_u + Q_d}{Q_e} \right)^2 \left(\frac{\alpha}{4\pi} \right)^2 \left(\frac{m_\pi}{m_e} \right)^2 \left(\theta_{a\pi} + \frac{5}{3} \theta_{a\eta_{ud}} + \frac{\sqrt{2}}{3} \theta_{a\eta_s} \right)^2 \tag{6.20}$$

where $\beta_e = \sqrt{1 - 4m_e^2/m_a^2}$. Note the very weak dependence of this branching ratio on the axion mass when $m_a \gg m_e$.

From (6.20) we can estimate the branching ratio for the rare decay $K^+ \rightarrow \pi^+(a \rightarrow \gamma\gamma)$ in terms $\text{Br}(K^+ \rightarrow \pi^+(a \rightarrow e^+e^-))$:

$$\text{Br}(K^+ \rightarrow \pi^+(a \rightarrow \gamma\gamma)) \approx 10^{-11} \left(\frac{\theta_{a\pi} + \frac{5}{3} \theta_{a\eta_{ud}} + \frac{\sqrt{2}}{3} \theta_{a\eta_s}}{10^{-2}} \right)^2 \left(\frac{\text{Br}(K^+ \rightarrow \pi^+(a \rightarrow e^+e^-))}{10^{-6}} \right), \tag{6.21}$$

which is still below current experimental sensitivity by two or three orders of magnitude [84].

6.2 Axion-nucleon couplings and nuclear de-excitations

Another classic signature of visible axions is axion emission in nuclear de-excitations, first studied in [85–87]. We shall briefly review here the standard results from the literature.

From (6.12) or (6.14) we can infer the axion coupling to protons and neutrons:

$$\mathcal{L}_{aNN} = \bar{N} i\gamma_5 g_{aNN} a N, \tag{6.22}$$

$$g_{aNN} = \frac{2m_N}{f_\pi} \left(\frac{\theta_{a\pi}}{2} g_A^{(3)} \tau^3 + \frac{\theta_{a\eta_8}}{2\sqrt{3}} g_A^{(8)} + \frac{\theta_{a\eta_0}}{\sqrt{6}} g_A^{(0)} \right), \tag{6.23}$$

where $N = \begin{pmatrix} p \\ n \end{pmatrix}$ is the nucleon isospin doublet, m_N is the nucleon mass, and $\tau^3 = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}$ is a Pauli matrix. The axial coupling constants, $g_A^{(i)}$, are the axial-vector form factors of $\langle N | J_5^i | N \rangle$ at $q^2 = 0$,

$$g_A^{(3)} = \Delta u - \Delta d, \tag{6.24}$$

$$g_A^{(8)} = \Delta u + \Delta d - 2 \Delta s, \tag{6.25}$$

$$g_A^{(0)} = \Delta u + \Delta d + \Delta s, \tag{6.26}$$

where Δq is defined as:

$$2 s_\mu \Delta q = \langle N | \bar{q} \gamma_\mu \gamma_5 q | N \rangle \tag{6.27}$$

with s_μ being the nucleon spin-vector. In terms of Δq , we can alternatively recast g_{aNN} as:

$$g_{aNN} = \frac{m_N}{f_\pi} \left(\theta_{a\pi} (\Delta u - \Delta d) \tau^3 + \theta_{a\eta_{ud}} (\Delta u + \Delta d) + \sqrt{2} \theta_{a\eta_s} \Delta s \right). \tag{6.28}$$

The pion-nucleon coupling, set by $g_A^{(3)}$, is well determined from nuclear β -decay,

$$g_A^{(3)} = \Delta u - \Delta d \simeq 1.27. \tag{6.29}$$

The other axial coupling constants, $g_A^{(8)}$ and $g_A^{(0)}$, are much harder to extract from experiment, and various estimates based on data from semi-leptonic hyperon decays, proton deep inelastic scattering, and lattice calculations yield the following ranges [88–100]:

$$0.09 \lesssim \Delta u + \Delta d \lesssim 0.62 \quad \text{and} \quad -0.35 \lesssim \Delta s \lesssim 0. \tag{6.30}$$

These uncertainties, compounded by the difficulties in extracting the axion hadronic mixing angles $\theta_{a\pi/\eta}$, are obstacles to making firm predictions for nuclear de-excitation rates via axion emission.¹²

Nevertheless, the parametric dependence of such rates on nuclear and axion properties is relatively well understood. Donnelly et al. [86] showed that the axion acts as a “magnetic photon” in nuclear transitions, and using standard multipole techniques in the

¹²Ref. [101] also considered properties of the axion in great detail, extracting the axion mass and coupling to photons and nucleons by combining results from Lattice QCD (LQCD) and χ PT at NLO. We believe that the authors of [101] were overly optimistic regarding the smallness of the errors in some of the LECs used, as well as the errors in LQCD extractions of the nucleon spin content Δq . But more importantly, their approach differs substantially from ours. For instance, they associate the axion with the Bardeen-Tye current in the two-flavor effective theory (i.e., chiral EFT with the strange quark integrated out). While the inclusion of NLO contributions in χ PT should in principle correct for the use of the (non physical) Bardeen-Tye current, it is unclear to us whether the authors of [101] did it in a consistent manner such that the resulting axion couplings to photons and nucleons would indeed correspond to that of the mass eigenstate. For instance, applying their procedure to our model to extract the couplings of the axion to nucleons, one would be led to conclude that our axion variant does not couple to nucleons, which is incorrect. (In particular, in their notation, our model would correspond to: $c_u^0 = Q_u/3$, $c_d^0 = Q_d/3$, $c_{s,c,b,t}^0 = 0$, $f_a \mapsto (Q_u + Q_d)/f_a$, $E/N = 2$ in eq. (1) of [101]; this would translate into $c_u = c_d = c_{s,c,b,t} = 0$ in eq. (46) of [101]). A thorough comparison of our treatment with that of [101] is a non-trivial task which is beyond the scope of this paper.

long-wavelength limit, derived the ratio of axion to photon de-excitation rates of a nuclear state. For an isoscalar ($\Delta T = 0$), p -wave magnetic (M1) transition, [86] derived:

$$\frac{\Gamma_a}{\Gamma_\gamma} \Big|_{\Delta T=0} = \frac{1}{2\pi\alpha} \left(\frac{g_{aNN}^{(0)}}{\mu^{(0)} - \eta^{(0)}} \right)^2 \left(\frac{|\vec{p}_a|}{|\vec{p}_\gamma|} \right)^3 \quad (6.31)$$

$$\approx \frac{1}{2\pi\alpha} \frac{m_N^2}{f_\pi^2} \left(\frac{\theta_{a\eta_{ud}}(\Delta u + \Delta d) + \sqrt{2}\theta_{a\eta_s}\Delta s}{\mu^{(0)} - \eta^{(0)}} \right)^2 \left(1 - \frac{m_a^2}{\Delta E^2} \right)^{3/2}. \quad (6.32)$$

Above, ΔE is the energy splitting between the two nuclear levels. The parameters $\mu^{(0)} \approx 0.88$ and $\eta^{(0)} \approx 1/2$ are related to the nuclear magnetic moment and the ratio of convection to magnetization currents, respectively.

For an isovector ($\Delta T = 1$) M1 transition, the analogous expression is [86]:

$$\frac{\Gamma_a}{\Gamma_\gamma} \Big|_{\Delta T=1} = \frac{1}{2\pi\alpha} \left(\frac{g_{aNN}^{(1)}}{\mu^{(1)} - \eta^{(1)}} \right)^2 \left(\frac{|\vec{p}_a|}{|\vec{p}_\gamma|} \right)^3 \quad (6.33)$$

$$\approx \frac{1}{2\pi\alpha} \left(\frac{m_N g_A^{(3)}}{f_\pi} \right)^2 \left(\frac{\theta_{a\pi}}{\mu^{(1)}} \right)^2 \left(1 - \frac{m_a^2}{\Delta E^2} \right)^{3/2}, \quad (6.34)$$

where $(\mu^{(1)} - \eta^{(1)}) \approx \mu^{(1)} \approx 4.7$. For mixed isospin transitions, the generalization is [102]:

$$\frac{\Gamma_a}{\Gamma_\gamma} = \frac{1}{2\pi\alpha} \left(\frac{c_0 g_{aNN}^{(0)} + c_1 g_{aNN}^{(1)}}{c_0 (\mu^{(0)} - \eta^{(0)}) + c_1 (\mu^{(1)} - \eta^{(1)})} \right)^2 \left(\frac{|\vec{p}_a|}{|\vec{p}_\gamma|} \right)^3, \quad (6.35)$$

where $c_{0,1}$ are the probability amplitudes for the different isospin components.

During the 80's, several experiments have searched for variants of the visible QCD axion in nuclear de-excitations [102–110]. Since most nuclear levels have splittings of a few MeV, the results of these experiments were only relevant for a range of m_a already ruled-out by other measurements, in particular beam dump experiments (see section 7). In 1990, de Boer et al. [111] studied transitions of the ^8Be nucleus which were energetic enough to probe axion masses of up to ≈ 15 MeV. A few years later, however, the results of this experiment were revisited by the authors in [112, 113] and the claimed limits were weakened substantially. We shall therefore not consider the results of [111–113] here.

Most recently, the ATOMKI collaboration [114] has measured several ^8Be nuclear transitions via emission of e^+e^- pairs. The two relevant transitions for our discussion are the M1 de-excitations of the $J^P = 1^+$ isospin doublet states, namely, $^8\text{Be}^*(17.64)$ and $^8\text{Be}^*(18.15)$, to the $J^P = 0^+$ isospin singlet ground state, $^8\text{Be}(0)$:

$$^8\text{Be}^*(17.64) \rightarrow ^8\text{Be}(0), \quad \Delta E = 17.64 \text{ MeV}, \quad \Delta T \approx 1, \quad (6.36)$$

$$^8\text{Be}^*(18.15) \rightarrow ^8\text{Be}(0), \quad \Delta E = 18.15 \text{ MeV}, \quad \Delta T \approx 0. \quad (6.37)$$

The ATOMKI collaboration claimed to have observed a deviation in the e^+e^- spectrum of the $^8\text{Be}^*(18.15) \rightarrow ^8\text{Be}(0)$ transition relative to the SM prediction of internal pair conversion ($\gamma^* \rightarrow e^+e^-$). According to [114], this deviation was consistent with the on-shell emission of a narrow resonance X of mass $m_X \approx (16.6 \pm 0.9)$ MeV promptly decaying

to e^+e^- . The best fit for the relative de-excitation rate was $\Gamma_X/\Gamma_\gamma \approx 5.8 \times 10^{-6}$, with a statistical significance of 6.8σ . Moreover, in the original publication [114], no excess was observed in the e^+e^- spectrum of the ${}^8\text{Be}^*(17.64) \rightarrow {}^8\text{Be}(0)$ transition. No error bars were quoted for either measurement, neither were upper bounds on emission rates of generic new particles, such as light vectors or pseudoscalars. Subsequent studies have attempted to understand these results via nuclear physics models [115], or via emission of a new light resonance [116–123].

The ATOMKI collaboration is revisiting these measurements with an improved experimental set-up [124, 125]. At the time of writing of this paper, preliminary results by Krasznahorkay et al. have been released [125] indicating a possible excess in the ${}^8\text{Be}^*(17.64) \rightarrow {}^8\text{Be}(0)$ transition as well, which was fit by a similar hypothetical particle X with mass $m_X = (17.0 \pm 0.2) \text{ MeV}$, at a rate of $\Gamma_X/\Gamma_\gamma \approx 4.0 \times 10^{-6}$.

At present, we believe the claims from the ATOMKI collaboration are inconclusive, and an independent measurement is warranted. Nonetheless, we can make order of magnitude predictions for axion emission in ${}^8\text{Be}$ M1 transitions using (6.35). While the excited states ${}^8\text{Be}^*(17.64)$ and ${}^8\text{Be}^*(18.15)$ are predominantly $T = 1$ and $T = 0$, respectively, their widths strongly indicate that they are isospin-mixed:

$$|{}^8\text{Be}^*(17.64)\rangle = \sin\theta_{1+} |T=0\rangle + \cos\theta_{1+} |T=1\rangle, \quad (6.38)$$

$$|{}^8\text{Be}^*(18.15)\rangle = \cos\theta_{1+} |T=0\rangle - \sin\theta_{1+} |T=1\rangle. \quad (6.39)$$

Estimates on the level of isospin mixing vary between $0.18 \lesssim \sin\theta_{1+} \lesssim 0.43$ [118, 122, 126]. Taking the value of $\sin\theta_{1+} = 0.35$ suggested in [122], we obtain the following range for the ${}^8\text{Be}^*(18.15)$ axion de-excitation rate:

$$\begin{aligned} \left. \frac{\Gamma_a}{\Gamma_\gamma} \right|_{{}^8\text{Be}^*(18.15)} &= \frac{1}{2\pi\alpha} \left(\frac{\cos\theta_{1+} g_{aNN}^{(0)} - \sin\theta_{1+} g_{aNN}^{(1)}}{\cos\theta_{1+} (\mu^{(0)} - \eta^{(0)}) - \sin\theta_{1+} (\mu^{(1)} - \eta^{(1)})} \right)^2 \left(1 - \frac{m_a^2}{(18.15 \text{ MeV})^2} \right)^{3/2} \\ &\approx (1 - 11) \times 10^{-6} \quad \text{for} \quad \frac{|\theta_{a\eta_{ud}}(\Delta u + \Delta d) + \sqrt{2}\theta_{a\eta_s}\Delta s|}{10^{-4}} \sim (1.4 - 4). \end{aligned} \quad (6.40)$$

Above, we also assume $m_a = 16.6 \text{ MeV}$ and $\theta_{a\pi} = 0.5 \times 10^{-4}$. Reiterating our point, while uncertainties in nuclear and axion parameters preclude us from making a precise prediction for the rate of axion emission in the de-excitation of ${}^8\text{Be}^*(18.15)$, the range in (6.40) is compatible with the excess observed by the ATOMKI collaboration.

Analogously, for the ${}^8\text{Be}^*(17.64)$ de-excitation we obtain:

$$\begin{aligned} \left. \frac{\Gamma_a}{\Gamma_\gamma} \right|_{{}^8\text{Be}^*(17.64)} &= \frac{1}{2\pi\alpha} \left(\frac{\sin\theta_{1+} g_{aNN}^{(0)} + \cos\theta_{1+} g_{aNN}^{(1)}}{\sin\theta_{1+} (\mu^{(0)} - \eta^{(0)}) + \cos\theta_{1+} (\mu^{(1)} - \eta^{(1)})} \right)^2 \left(1 - \frac{m_a^2}{(17.64 \text{ MeV})^2} \right)^{3/2} \\ &\approx (0.5 - 1.7) \times 10^{-7} \quad \text{for} \quad \frac{|\theta_{a\eta_{ud}}(\Delta u + \Delta d) + \sqrt{2}\theta_{a\eta_s}\Delta s|}{10^{-4}} \sim (1.4 - 4). \end{aligned} \quad (6.41)$$

Here, despite uncertainties in (6.41), we are able to make the more firm prediction that, should an anomaly persist in the ${}^8\text{Be}^*(17.64)$ de-excitation rate at the level of $\Gamma_X/\Gamma_\gamma \approx \mathcal{O}(10^{-6})$, this would preclude our QCD axion variant from offering as a viable explanation of this excess.

7 Axion-electron coupling

Unlike its hadronic couplings, the axion coupling to electrons bears no consequence to the solution of the strong CP problem, and therefore it is a model dependent parameter that can be adjusted according to phenomenological constraints. Moreover, it is much less susceptible to calculational uncertainties, and can be probed in a variety of existing and upcoming experiments. In fact, it is conceivable that this MeV axion variant could be definitively excluded via its electron couplings before any substantial progress is made regarding its hadronic phenomenology. In this section, we will review existing bounds on the axion-electron coupling, and briefly discuss on-going experimental efforts to search for hidden photons that could test the MeV axion scenario as well.

For specificity, we reiterate the axion coupling to electrons:

$$\mathcal{L}_a^{\text{eff}} \supset \frac{Q_e}{f_a} m_e a \bar{e} i \gamma_5 e, \quad (7.1)$$

as well as the relation between the axion's mass and decay constant:

$$\begin{aligned} m_a &= |Q_u + Q_d| \frac{\sqrt{m_u m_d}}{m_u + m_d} \frac{m_\pi f_\pi}{f_a} \\ &\approx \frac{\sqrt{2} m_\pi f_\pi}{f_a}, \end{aligned} \quad (7.2)$$

and the axion lifetime, assuming that it is dominated by $a \rightarrow e^+ e^-$:

$$\tau_a^{-1} = \frac{1}{8\pi} \left(\frac{Q_e}{f_a} m_e \right)^2 m_a. \quad (7.3)$$

The constraints discussed in this section are summarized in figure 1.

7.1 Beam dump constraints

In the 80's, several beam dump experiments have specifically targeted the QCD axion. Since the results of these searches require no re-interpretation, we refer the reader to the original papers for details on production and detection mechanisms. In this subsection, we compile the most significant constraints in the region of parameter space of interest.

Constraints from beam dumps can be avoided if axions are sufficiently short-lived so as to decay in the earth shielding, before reaching the detectors. Moreover, in order to remain experimentally viable, invisible decay modes of the axion must be subdominant by at least $\mathcal{O}(10^{-4})$ in order to avoid stringent constraints from $K^+ \rightarrow \pi^+(a \rightarrow \text{invisible})$. In order to fulfill these requirements, the axion must be heavier than a few MeV and couple significantly to electrons.

In figure 1, we show contours of the axion lifetime as a function of the axion mass and coupling to electrons, Q_e/f_a . We also show the most relevant beam dump constraints as shaded gray regions, namely, FNAL E774 [127], SLAC E141 [128], and Orsay [129]. Other overlapping but less powerful constraints, such as Bechis et al. [132], SLAC E56 [133], FNAL E605/772 [134, 135], and KEK [136] are omitted in order to make the plot more readable. Experiments with longer earth shieldings, such as SLAC E137 [45] and CHARM [44] lie outside the plot.




Figure 1. Existing constraints on the axion-electron coupling for our specific axion variant. The red curves illustrate the m_a vs. $|Q_e|/f_a$ relation for different values of $|Q_e|$. The black dashed lines are contours of the axion’s lifetime. The shaded gray regions are excluded by beam dump experiments [127–129], and the shaded orange and green regions are excluded by BaBar [130] and KLOE [131] searches for dark photons, respectively. Constraints from the electron’s anomalous magnetic moment are model dependent and therefore not shown in this plot (see section 7.2 for details).

7.2 Constraints from $(g - 2)_e$

Next, we discuss the MeV axion contribution to the electron’s anomalous magnetic moment. Constraints coming from $(g - 2)_e$ are difficult to pin down due to uncertainties in the axion’s two-loop contribution, as well as the model dependence associated with UV completions of MeV axions. As we shall discuss in section 9, generic UV completions of such models contain additional particles that can also contribute substantially to $(g - 2)_e$.

State-of-the-art calculations of $a_e \equiv (g - 2)_e/2$ in the SM predict [137]:

$$a_e^{\text{SM}} = 1\,159\,652\,181.643(25)(23)(16)(763) \times 10^{-12}, \tag{7.4}$$

where the first three uncertainties are theoretical, and the last one stems from the error in the measurement of the fine structure constant.

The most precise measurement of a_e to date [138, 139], on the other hand, gives:

$$a_e^{\text{exp}} = 1\,159\,652\,180.73(0.28) \times 10^{-12}, \tag{7.5}$$

and differs from the theoretical expectation in the SM by:

$$a_e^{\text{exp}} - a_e^{\text{SM}} = -(0.91 \pm 0.77 \pm 0.28) \times 10^{-12}. \tag{7.6}$$

The dominant contribution to this quantity from the MeV axion comes at one-loop and is given by:

$$\begin{aligned} \Delta a_e^{1\text{-loop}} &= -\frac{2}{(4\pi)^2} \left(\frac{Q_e}{f_a} m_e\right)^2 \left(\frac{m_e}{m_a}\right)^2 \int_0^1 dx \frac{x^3}{(1-x) + \left(\frac{m_e}{m_a}\right)^2 x^2} \\ &\approx -\frac{Q_e^2}{(4\pi)^2} \left(\frac{m_e^2}{m_\pi f_\pi}\right)^2 \left(\log\left[\frac{m_a^2}{m_e^2}\right] - \frac{11}{6}\right), \end{aligned} \tag{7.7}$$

where we have used (7.2) in the second equality.

The two-loop contribution (which is a Barr-Zee type diagram) suffers from large uncertainties due to the poor determination of $g_{a\gamma\gamma}$, as well as the precise value of the cut-off scale Λ above which $g_{a\gamma\gamma}$ can no longer be treated as a point-like coupling [140]. Setting $\Lambda \sim 4\pi f_\pi$, we have [141–143]:

$$\Delta a_e^{2\text{-loop}} \sim \left(\frac{Q_e}{f_a} m_e\right) \frac{m_e}{\pi^2} g_{a\gamma\gamma} f_{PS} \left[\frac{4\pi f_\pi}{m_a}\right], \tag{7.8}$$

where, from (6.18),

$$g_{a\gamma\gamma} = \frac{\alpha}{4\pi f_\pi} \left(\theta_{a\pi} + \frac{5}{3}\theta_{a\eta_{ud}} + \frac{\sqrt{2}}{3}\theta_{a\eta_s}\right), \tag{7.9}$$

and

$$\begin{aligned} f_{PS}[z] &= \int_0^1 dx \frac{z^2/2}{x(1-x) - z^2} \log\left(\frac{x(1-x)}{z^2}\right) \\ &\approx \log z + 1 \quad \text{for } z \gg 1. \end{aligned} \tag{7.10}$$

A few observations are in order: the one-loop contribution is negative and exceeds the allowed range in (7.6) by more than two standard deviations if $|Q_e| \gtrsim 1/2$. However, the two-loop contribution has the opposite sign if the product $Q_e g_{a\gamma\gamma}$ is positive, and could in principle partially cancel the one-loop contribution to a substantial degree. In particular, if the linear combination of hadronic mixing angles in (7.9) is $\sim 2 \times 10^{-2}$, then $Q_e \lesssim 1.3$ is still consistent with (7.6) at 2σ . Note that this range of mixing angles, and therefore of $g_{a\gamma\gamma}$, is consistent with other experimental constraints, mainly because of the very poor determination of $\theta_{a\eta_s}$. The rare decay $K^+ \rightarrow \pi^+(a \rightarrow \gamma\gamma)$ is not yet sensitive to $\theta_{a\eta_s} \lesssim \mathcal{O}(10^{-1})$ (see section 6.1), and the contribution of $\theta_{a\eta_s}$ to the ${}^8\text{Be}$ nuclear transitions is mild if $\Delta s \sim -0.02$, which is suggested by recent lattice results [94–100].

Given all these considerations, existing measurements of the electron anomalous magnetic moment cannot provide a robust and model independent exclusion of the MeV axion parameter space, and therefore we do not include it in figure 1.

7.3 Constraints from searches for dark photons

Over the past decade, there has been a spur in phenomenological studies of light, weakly coupled vectors (“dark photons”), mostly motivated by dark matter phenomenology [144–147].

As a result, many experimental proposals have been put forth to search for dark photons in the near future [148–151]. As it turns out, dark photon production and detection mechanisms are very similar to those of light pseudoscalars such as the axion, and several constraints on dark photons decaying visibly can be easily recast as limits on the MeV axion.

In this section, we briefly comment on two experimental strategies that are relevant for MeV axions, namely, production in e^+e^- collisions, and at fixed-target experiments. We also translate existing bounds as well as future projections of proposed experiments in the MeV axion parameter space.

The axion can be produced in association with a photon in e^+e^- annihilation via the $a\bar{e}i\gamma_5e$ vertex. In the limit that the center-of-mass energy $\sqrt{s} \gg m_a$, the cross-section computed at leading order is:

$$\frac{d\sigma(e^+e^- \rightarrow \gamma a)}{d\cos\theta_\gamma} = \left(\frac{Q_e}{f_a} m_e\right)^2 \frac{\alpha}{2s \sin^2\theta_\gamma + \frac{m_e^2}{2}}, \quad (7.11)$$

where θ_γ is the angle of the final state photon with respect to the beam axis, and we show the explicit dependence on m_e that regulates the collinear divergence when $|\sin\theta_\gamma| \rightarrow 0$.

Recent results from BaBar [130] and KLOE [131] looking for $e^+e^- \rightarrow \gamma (A' \rightarrow e^+e^-)$ can be easily recast in the axion parameter space using (7.11). We display the corresponding limits in figure 1.

The axion can also be produced in fixed target experiments via “axion-bremsstrahlung” when an energetic electron scatters off of a heavy nuclear target. The differential cross-section as a function of the axion emission angle with respect to the beam, θ_a , and the fraction x_a of the incoming electron energy carried by the radiated axion, $x_a \equiv E_a/E_0$, has been worked out in [152]. Below, we just quote the result:

$$\frac{d\sigma(eN \rightarrow eNa)}{dx_a d\cos\theta_a} = \frac{\alpha^2}{\pi} \left(\frac{Q_e}{f_a} m_e\right)^2 \beta_a E_0^2 \frac{x_a}{U^2} \left(\frac{x_a^2}{2} - \frac{m_a^2}{U^2} x_a(1-x_a)(E_0^2 \theta_a^2 x_a)\right) \chi, \quad (7.12)$$

where

$$U(x_a, \theta_a) \equiv E_0^2 \theta_a^2 x_a + m_a^2 \frac{1-x_a}{x_a} + m_e^2 x_a \quad , \quad \beta_a \equiv \sqrt{1 - m_a^2/E_0^2}, \quad (7.13)$$

and χ encodes nuclear form factors (in the notation of [145], $\chi = Z^2 \mathcal{L}og$). For further details, see [152–154].

Much like the case of dark photon bremsstrahlung, axion emission is almost collinear, and its characteristic emission angle is parametrically smaller than the angle of its decay products with respect to the incident beam [145]. Integrating (7.12) over emission angles, we obtain

$$\frac{d\sigma(eN \rightarrow eNa)}{dx_a} = \frac{\alpha^2}{3\pi m_a^2} \left(\frac{Q_e}{f_a} m_e\right)^2 \beta_a \chi \frac{x_a^3}{(1-x_a) + \left(\frac{m_e}{m_a}\right)^2 x_a^2}, \quad (7.14)$$

which is more peaked towards $x_a = 1$ than the corresponding $x_{A'}$ dependence of dark photon emission.




Figure 2. Projected reach of several planned dark photon experiments into the MeV axion parameter space, based on [151]. Gray curves and gray shaded areas are from figure 1.

Finally, we quote the total production cross-section integrated over angle and energy:

$$\begin{aligned} \sigma(eN \rightarrow eNa) &= \frac{\alpha^2}{3\pi m_a^2} \left(\frac{Q_e m_e}{f_a} \right)^2 \chi \left(\log \left[\frac{1}{(1-x_a)_c} \right] - \frac{11}{6} + \mathcal{O} \left(\frac{m_a^2}{E_0^2} \right) \right) \\ &\approx \frac{\alpha^2 Q_e^2}{6\pi} \left(\frac{m_e}{m_\pi f_\pi} \right)^2 \chi \left(\log \left[\frac{1}{(1-x_a)_c} \right] - \frac{11}{6} + \mathcal{O} \left(\frac{m_a^2}{E_0^2} \right) \right), \end{aligned} \quad (7.15)$$

where

$$(1-x_a)_c \equiv \max \left(\frac{m_e^2}{m_a^2}, \frac{m_a^2}{E_0^2} \right), \quad (7.16)$$

and we have used (7.2) in the second equality. We see, therefore, that the total cross-section for axion emission increases logarithmically with m_a while $m_a \lesssim \sqrt{m_e E_0}$, whereas it decreases logarithmically with m_a in the regime $\sqrt{m_e E_0} \lesssim m_a \ll E_0$.

In figure 2, we translate the projected reach of several planned dark photon experiments into the MeV axion parameter space, using the following rule of thumb relating the dark photon kinetic mixing parameter $\epsilon_{A'}$ and the axion-electron coupling that would yield comparable signal strengths:

$$\epsilon_{A'}^2 \alpha \sim \frac{1}{4\pi} \left(\frac{Q_e}{f_a} m_e \right)^2. \quad (7.17)$$

We emphasize that (7.17) is only approximate, since the angular dependence of pseudoscalar cross-sections differs from that of vectors. Figure 2 includes ongoing and proposed dark photon searches via e^+e^- annihilation (VEPP-3 at BINP [155, 156], PADME in Frascati [157], MMAPS at Cornell [151] and Belle-II at KEK [158]), as well as dark photon

bremsstrahlung from electrons scattering off of heavy fixed targets (HPS [159] and Dark-Light [160, 161] at JLab, MAGIX [162] at Mainz). These projections were based on the *Dark Sectors 2016 Workshop* community report [151]. We refer the reader to this report for further details.

8 Other constraints

Finally, we comment on other potentially constraining observables that could probe the MeV axion parameter space. We discuss them in lesser detail because the resulting limits are either weaker than the ones previously discussed, suffer from similar hadronic uncertainties, or lack experimental information specific to the axion related signature.

Bounds from the hyperfine splitting between the 1^3S_1 and 1^1S_0 positronium levels [163, 164], for instance, are currently not competitive with beam dumps constraints shown in figure 1. Neither are bounds from neutron-nucleus and neutron-electron scattering [165, 166], since the axion mediated contribution is spin-dependent [167, 168].

Competitive constraints on axion hadronic mixing angles could be potentially obtained from dedicated analyses of existing data on other rare meson decays, such as $K^+ \rightarrow \ell^+ \nu_\ell a$, $K_L^0 \rightarrow \pi\pi a$, $K_S^0 \rightarrow \pi^0 a$, and $\eta, \eta' \rightarrow \pi\pi a$. The relevant final states for these observables have been measured in high statistics samples [169–181], but, to the best of our knowledge, no dedicated searches for e^+e^- resonances in the few MeV invariant mass range have been performed.

Other less sensitive rare decays also exist, which could become competitive if the branching ratio sensitivity were substantially improved. Examples are B meson decays,¹³ $B \rightarrow Ka$ [182, 183]; radiative quarkonia decays, $J/\psi, \Upsilon, \Phi \rightarrow \gamma a$ [38, 184–188]; hyperon decays, such as $\Sigma^+ \rightarrow p^+ a$ [189]; and the not-yet-observed decays $\eta, \eta' \rightarrow e^+e^-$ [190, 191], which are highly suppressed in the SM.

We emphasize that axion production via its coupling to photons¹⁴ should be subdominant to the processes discussed in section 7.3 induced by the axion’s electronic couplings. This can be easily seen by noting that the effective axion-photon coupling, $g_{a\gamma\gamma} a F\tilde{F}$, is suppressed by an effective scale of $\mathcal{O}(10 - 100)$ TeV:

$$g_{a\gamma\gamma} = \frac{\alpha}{4\pi f_\pi} \left(\theta_{a\pi} + \frac{\theta_{a\eta_8}}{\sqrt{3}} + 2 \frac{\theta_{a\eta_0}}{\sqrt{3/2}} \right) \sim \frac{\mathcal{O}(0.1 - 1)}{10 \text{ TeV}}. \quad (8.1)$$

A comparable effective scale suppresses the couplings of the axion to $Z\gamma$, thereby suppressing the rate for the decay $Z \rightarrow \gamma a$ below current experimental bounds. This rare Z decay is typically a sensitive probe of generic “axion-like particles” (ALPs) with low decay constants (see recent studies in [192–194]). A generic ALP ϕ with decay constant f_ϕ couples to γZ with typical strength $g_{\phi\gamma Z} \sim \frac{\alpha}{4\pi f_\phi}$. This parametric dependence does not apply for an axion with $f_a \sim \mathcal{O}(\text{GeV})$ for the following reason: in the case of a generic ALP, the ϕ - γ - Z coupling is generated by integrating out electroweakly charged fermions

¹³Note that EW penguins with charm or top quarks in the loop do not contribute in this case, because the axion does not couple to heavy quarks.

¹⁴See [192] for a recent study of ALP photo-production.

heavier than M_Z . An MeV axion, on the other hand, is typically precluded from coupling to such heavy fermions by the fact that the PQ symmetry must be preserved down to the scale of $f_a \sim \mathcal{O}(1 - 10)$ GeV. In other words, the MeV axion can only have perturbative couplings to fermions with mass $m_f = y_f \mathcal{O}(f_a) \lesssim \mathcal{O}(f_a) \ll M_Z$. Loops of such light fermions induce much smaller $g_{a\gamma Z}$ couplings, $g_{a\gamma Z} \sim \frac{m_f}{M_Z} \frac{\alpha}{4\pi f_a}$.

So far we have assumed that the MeV axion does not couple to neutrinos. This can be naturally realized if neither the lepton doublets nor right-handed Dirac and/or sterile neutrinos carry PQ charge. If we relax this assumption, the typical axion coupling to neutrinos would scale as $m_\nu/f_a \sim \mathcal{O}(10^{-11} - 10^{-10})$, which is likely too small to be phenomenologically interesting. On the other hand, PQ-charged *sterile* neutrinos could couple to the axion much more strongly, namely, with strength $\mathcal{O}(m_{\nu_s}/f_a)$. If m_{ν_s} is heavy enough, this could lead to observable signatures. A particularly well motivated mass range for sterile neutrinos is $m_{\nu_s} \sim \mathcal{O}(1 - 10)$ keV, where sterile neutrinos can constitute warm dark matter and potentially address structure formation problems such as the core vs cusp and missing satellites problems (see [195] and references therein). In this mass range, the branching ratio for axion decay into sterile neutrinos would be:

$$\text{Br}(a \rightarrow \nu_s \nu_s) \sim \mathcal{O}\left(\frac{m_{\nu_s}^2}{m_e^2}\right) \sim \mathcal{O}(10^{-4} - 10^{-6}). \quad (8.2)$$

An interesting consequence of (8.2) would be a contribution to the rare kaon decay $K^+ \rightarrow \pi^+ a \rightarrow \pi^+ \nu_s \nu_s$ at the level of:

$$\text{Br}(K^+ \rightarrow \pi^+ \nu_s \nu_s) \sim \mathcal{O}(10^{-10} - 10^{-12}) \left(\frac{\text{Br}(K^+ \rightarrow \pi^+ a)}{10^{-6}} \right), \quad (8.3)$$

which is compatible with the strongest bounds set by BNL's E787/E949 experiments [48–50], namely, $\text{Br}(K^+ \rightarrow \pi^+ (X^0 \rightarrow \text{invisible})) \lesssim 0.45 \times 10^{-10}$ for $m_{X^0} \lesssim 70$ MeV. CERN's NA62 experiment will soon supersede these bounds by at least one order of magnitude, since it is expected to have sensitivity to the SM prediction of $\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})|_{\text{SM}} = (8.4 \pm 1.0) \times 10^{-11}$ [196] with better than 10% accuracy [197].

Other interesting possibilities for neutrino phenomenology could come from off-diagonal couplings of the axion to active (ν_ℓ) and sterile neutrinos (ν_s), such as $\lambda a \nu_\ell \nu_s$, especially if the sterile neutrinos are light enough to be produced in laboratory experiments and/or astrophysical processes. Effects such as $\nu_s \rightarrow \nu_\ell a \rightarrow \nu_\ell e^+ e^-$ (if $m_{\nu_s} > m_a$), non-standard neutrino interactions, MSW-type resonance effects in neutrino propagation in matter, and neutrino transport in core-collapse supernova are interesting directions to explore, but are beyond the scope of this paper.

9 GeV scale completions of MeV axion models

The Peccei-Quinn breaking scale f_a suggests that there is new dynamics around 1–10 GeV. While there are many constraints on what type of new particles may be associated with this new dynamics, there is still a large degree of model dependence in UV completions of the MeV axion. A thorough exploration of the viable phenomenology is beyond the scope of

this paper, and will be deferred to a future publication [198]. In this section, we briefly illustrate a few possibilities with a simple toy-model.

Consider introducing two new complex scalar degrees of freedom, Φ_u and Φ_d , with PQ charges $Q_{\Phi_u}^{\text{PQ}} = Q_u = 2$ and $Q_{\Phi_d}^{\text{PQ}} = Q_d = 1$, respectively. We can then UV complete the PQ mechanism at the GeV scale by writing:

$$\mathcal{L}_{\text{PQ}}^{\Lambda=\text{GeV}} \supset -(y_u \Phi_u uu^c + y_d \Phi_d dd^c + \text{h.c.}) - V(\Phi_u, \Phi_d). \quad (9.1)$$

The scalar potential $V(\Phi_u, \Phi_d)$ enforces the PQ symmetry, and induces vacuum expectation values for Φ_u and Φ_d , hence breaking the PQ symmetry:

$$\langle \Phi_u \rangle = \frac{f_u}{\sqrt{2}} = \frac{m_u}{y_u}, \quad (9.2)$$

$$\langle \Phi_d \rangle = \frac{f_d}{\sqrt{2}} = \frac{m_d}{y_d}. \quad (9.3)$$

We can then decompose Φ_u and Φ_d into real scalar and pseudoscalar components:

$$\Phi_u = \left(\frac{f_u}{\sqrt{2}} + \frac{\varphi_u}{\sqrt{2}} \right) \text{Exp } i \left(Q_u \frac{a}{f_a} + \frac{Q_d}{\tan \beta_{\text{PQ}}} \frac{\eta_{\text{PQ}}}{f_a} \right), \quad (9.4)$$

$$\Phi_d = \left(\frac{f_d}{\sqrt{2}} + \frac{\varphi_d}{\sqrt{2}} \right) \text{Exp } i \left(Q_d \frac{a}{f_a} - Q_u \tan \beta_{\text{PQ}} \frac{\eta_{\text{PQ}}}{f_a} \right). \quad (9.5)$$

Above, $\tan \beta_{\text{PQ}} \equiv f_u/f_d$, the pseudoscalar a is the MeV axion, and the axion decay constant, f_a , is given by $f_a^2 \equiv Q_u^2 f_u^2 + Q_d^2 f_d^2 = 4f_u^2 + f_d^2$.

In this toy model, we have introduced 3 extra degrees of freedom besides the MeV axion: two real scalars, φ_u and φ_d , and one pseudoscalar, η_{PQ} . The natural scale for their masses is set by f_a . Such light states must, therefore, be electroweak singlets in order to be phenomenologically viable. As a consequence, the couplings of Φ_u and Φ_d to SM fermions in (9.1) descend from higher dimensional operators and must be generated after electroweak symmetry breaking. We shall return to this point in the next section.

Because of their couplings to light quarks, the new states φ_u , φ_d , and η_{PQ} are produced in hadronic interactions, and decay dominantly to hadrons - though with smaller cross-sections and narrower widths than typical QCD hadronic resonances. Unfortunately, estimating their widths and mixings with QCD resonances is challenging, since they lie in a regime where neither perturbative QCD nor chiral perturbation methods are reliable. It is conceivable, at least in principle, that these states are not excluded. The scalar resonances φ_u and φ_d , for instance, might not have been identified if lying in the murky mass range below 2 GeV [199], where many broad scalar resonances overlap and might constitute a large and complicated background to disentangle. The same argument is less likely to apply for the pseudoscalar resonance η_{PQ} , unless it lies in a mass range where the hadronic pseudoscalar spectrum is poorly understood. For instance, the 1300-1500 MeV range contains three 0^{-+} states, namely, $\eta(1295)$, $\eta(1405)$ and $\eta(1475)$. The exact nature of these states is still subject to debate [200], with interpretations ranging from those being radial excitations of lighter pseudoscalars [201–203] to pseudoscalar glueballs [204–212]. Some authors

dispute the existence of the $\eta(1295)$ state and claim that there is a single pseudoscalar meson in this mass range, the $\eta(1440)$ state, which would be the first radial excitation of the η [213, 214]. There are also claims that the $\eta(1405)$ and $\eta(1475)$ structures might originate from a single pole [215], the splitting being due to nodes in the decay amplitudes [213], or amplitude mixing via a triangular singularity [216, 217]. Whatever the nature of $\eta(1295)$, $\eta(1405)$ and $\eta(1475)$ may be, a priori they might present a challenging background for η_{PQ} , and possibly mix substantially with it. A less speculative and more careful investigation of these possibilities will be done in [198].

We conclude by commenting on the coupling of these states to electrons. One possibility is that either Φ_u or Φ_d have electron Yukawa couplings, in which case all real degrees of freedom (namely, φ_u , φ_d , η_{PQ} and a) will couple to electrons (either directly or through mixing), with typical strength $\mathcal{O}(m_e/f_a)$. Alternatively, we may introduce a third PQ-charged complex scalar, $\Phi_e = \varphi_e e^{i\eta_e/f_e}$, which will be responsible for generating the electron mass and the axion-electron coupling. Mixings between the Φ_e and $\Phi_{u,d}$ degrees of freedom will likewise result in couplings of $\varphi_{u,d,e}$ and $\eta_{\text{PQ},e}$ to electrons, although in this case some of the couplings might be suppressed or enhanced relative to m_e/f_a . Depending on the details of a specific model, these states might give important contributions to $(g-2)_e$, $\Gamma(\pi^0 \rightarrow e^+e^-)$, and might be searched for in e^+e^- annihilation and fixed target experiments as well. A phenomenological study will be considered in [198].

10 EW scale completions of MeV axion models

As discussed in the previous section, the dynamics that breaks the PQ symmetry at the GeV scale generically requires new light degrees of freedom coupling to $\text{SU}(2)_W$ -charged quark bilinears. However, these new light particles themselves must not be $\text{SU}(2)_W$ -charged, otherwise they would be excluded, for instance, by measurements of the Z^0 width. Therefore, the Yukawa couplings of the PQ sector to SM fermions violate $\text{SU}(2)_W \times \text{U}(1)_Y$ and can only arise from higher dimensional operators after EW symmetry breaking. These operators can be generated by integrating out new degrees of freedom at the EW scale, such as heavy PQ-charged scalar doublets, or heavy vector-like quarks, leading to interesting and distinct LHC signatures. Although a thorough exploration of all possibilities is beyond the scope of this paper, we will briefly consider a simple EW completion of the toy-model discussed in section 9, and comment on the associated phenomenology.

For each EW singlet PQ scalar Φ_f , we introduce a new $\text{SU}(2)_W$ doublet H_f with the same corresponding PQ charge, $Q_{H_f}^{\text{PQ}} = Q_{\Phi_f}^{\text{PQ}}$. We can then write EW preserving Yukawa couplings, $H_f f_L f^c$, and tri-scalar “ A -terms”:

$$\begin{aligned} \mathcal{L}_{\text{PQ}}^{\Lambda=\text{EW}} \supset & - (Y_u H_u Q_1 u^c + Y_d H_d Q_1 d^c + Y_e H_e L_1 e^c + \text{h.c.}) \\ & - (A_u \Phi_u^\dagger H_u H_{\text{SM}}^\dagger + A_d \Phi_d^\dagger H_d H_{\text{SM}} + A_e \Phi_e^\dagger H_e H_{\text{SM}} + \text{h.c.}). \end{aligned} \quad (10.1)$$

Above, the A_f coefficients have dimensions of mass, H_{SM} is the doublet responsible for EW symmetry breaking and contains the 125 GeV Higgs, and H_e , Φ_e may be independent fields, or may be identified with H_d , Φ_d or H_u^\dagger , Φ_u^\dagger .

Giving large masses $m_{H_f} \gtrsim \mathcal{O}(100 \text{ GeV})$ for the new PQ doublets H_f , we can integrate them out and obtain the effective interactions below the EW symmetry breaking scale:

$$\mathcal{L}_{\text{PQ}}^{\Lambda=\text{GeV}} \supset - (y_u \Phi_u uu^c + y_d \Phi_d dd^c + y_e \Phi_e ee^c + \text{h.c.}) \quad (10.2)$$

where

$$y_f \equiv Y_f \frac{A_f \langle H_{\text{SM}} \rangle}{m_{H_f}^2}, \quad (10.3)$$

with $\langle H_{\text{SM}} \rangle = 174 \text{ GeV}$.

A parameter of particular relevance for phenomenology is the mixing angle between the light singlet Φ_f and the neutral component of the heavy doublet H_f^0 :

$$\theta_{\Phi H}^f \equiv \frac{A_f \langle H_{\text{SM}} \rangle}{m_{H_f}^2}. \quad (10.4)$$

Firstly, this parameter quantifies the degree of tree-level tuning required to maintain the hierarchy between the singlet states (at the GeV scale) and the doublet states (at the EW scale). A simple measure of this fine-tuning is given by:

$$\text{F.T.} \sim \frac{1}{\theta_{\Phi H}^f} \frac{m_{\Phi_f}^2}{m_{H_f}^2} \sim \frac{\mathcal{O}(10^{-4})}{\theta_{\Phi H}^f}. \quad (10.5)$$

Imposing a tolerance of at most 10% tuning, (10.5) requires $\theta_{\Phi H}^f \lesssim 0.03$.

Secondly, $\theta_{\Phi H}^f$ indirectly determines the coupling of H_f to quarks. Considering the up quark for concreteness, we have from (10.3) and (9.2),

$$Y_u = \sqrt{2} \frac{m_u}{f_u} \frac{1}{\theta_{\Phi H}^f} \approx 0.1 \times \left(\frac{0.03}{\theta_{\Phi H}^f} \right) \left(\frac{\text{GeV}}{f_u} \right). \quad (10.6)$$

Such sizable Yukawa coupling would lead to a large production of H_u at the LHC, with 13 TeV cross-sections ranging from $\mathcal{O}(1 - 10^3)$ pb for $\theta_{\Phi H}^f \sim (0.01 - 0.03)$ and $m_{H_u} \sim (100 - 500) \text{ GeV}$. Existing LHC searches for leptophobic vectors (Z') [218] can already place non-trivial upper bounds on processes such as $pp \rightarrow H_u(\rightarrow jj) + j$ [219–221] and $pp \rightarrow H_u(\rightarrow jj) + \gamma$ [221, 222], ranging from $Y_u \lesssim (0.1 - 0.4)$ depending on m_{H_u} . Similar considerations hold for H_d production.

Finally, $\theta_{\Phi H}^f$ controls a variety of rare decays of the 125 GeV Higgs h , Z^0 and H_f to final states with PQ scalars a , η_{PQ} , φ_u , φ_d . We compile some of these possibilities in table 2. The reach of existing measurements and future search strategies for these rare decays will depend on how the boosted PQ scalars will be tagged by LHC detection algorithms. Typically, the GeV states φ_u , φ_d , η_{PQ} will decay promptly to collimated final states of 2π , $\eta\pi$, $K\bar{K}$, 3π , $\eta\pi\pi$, etc, which may be tagged as jets, hadronic taus, photon-jets [223–225] (for final states of 2 or 3 π^0 's), or even single photons (if the detector's granularity is poor enough, which might be the case for LEP detectors). The much lighter and longer lived MeV axion a , with $c\tau_a$ ranging from $0.1 \mu\text{m}$ to 0.1 mm , will produce a

Rare Decay	Potential Signature	Signal Strength
$pp \rightarrow H_{u,d}^0 \rightarrow Z^0 a$	$Z + \gamma^*, Z + LJ$	$\sigma \sim \mathcal{O}(10^{-3} - 1)$ pb
$pp \rightarrow H_{u,d}^\pm \rightarrow W^\pm a$	$W + \gamma^*, W + LJ$	$\sigma \sim \mathcal{O}(10^{-3} - 1)$ pb
$pp \rightarrow H_{u,d}^0 \rightarrow h \varphi_{u,d}/\eta_{PQ}$	$h + j, h + PJ, h + \tau_h$	$\sigma \sim \mathcal{O}(0.1 - 1)$ pb
$pp \rightarrow H_{u,d}^0 \rightarrow Z^0 \varphi_{u,d}/\eta_{PQ}$	$Z + PJ, Z + \tau_h$	$\sigma \sim \mathcal{O}(10^{-3} - 1)$ pb
$pp \rightarrow H_{u,d}^\pm \rightarrow W^\pm \varphi_{u,d}/\eta_{PQ}$	$W + PJ, W + \tau_h$	$\sigma \sim \mathcal{O}(10^{-3} - 1)$ pb
$Z^0 \rightarrow a \phi_{u,d}$	$\Gamma_Z, Z \rightarrow \gamma^* j, Z \rightarrow \gamma^* PJ, Z \rightarrow LJ PJ$	$\text{Br} \sim \mathcal{O}(10^{-10} - 10^{-8})$
$Z^0 \rightarrow \eta_{PQ} \phi_{u,d}$	$R_Z, Z \rightarrow j PJ, Z \rightarrow PJ PJ$	$\text{Br} \sim \mathcal{O}(10^{-10} - 10^{-8})$
$h \rightarrow \phi_{u,d} \phi_{u,d}, \eta_{PQ} \eta_{PQ}$	$h \rightarrow j PJ, h \rightarrow PJ PJ, h \rightarrow \tau_h \tau_h$	$\text{Br} \sim \mathcal{O}(10^{-6} - 10^{-3})$
$h \rightarrow H_{u,d}^0 \varphi_{u,d}/\eta_{PQ} (m_{H_{u,d}} < m_h)$	$h \rightarrow (jj) j, h \rightarrow (jj) PJ, h \rightarrow (jj) \tau_h$	$\text{Br} \sim \mathcal{O}(10^{-3} - 10^{-2})$

Table 2. Compilation of rare decays that could potentially probe the model in (10.1). The middle column shows potential signatures, *assuming* that the MeV axion a would be tagged either as a converted photon (γ^*) or a lepton-jet (LJ); and that $\varphi_{u,d}, \eta_{PQ}$ would be tagged either as a jet (j), a photon-jet (PJ), or a hadronic tau (τ_h). The third column shows typical ranges for signal cross-sections/branching ratios assuming $\theta_{\Phi H}^f \sim (0.01 - 0.03)$ and $m_{H_u} \sim (100 - 300)$ GeV.

highly collimated e^+e^- pair, with a decay vertex displaced by a few mm to several cm. Here, too, the sensitivity of any particular analysis will depend on whether a is tagged as a converted photon, a prompt or displaced lepton-jet [226–237], or whether it will fail quality criteria for standard objects and simply be vetoed. A thorough study of these possibilities is deferred to [198].¹⁵ It suffices to say, nonetheless, that for $\theta_{\Phi H}^f \sim \mathcal{O}(10^{-2})$ all existing bounds are satisfied for the processes listed in table 2, regardless of assumptions on boosted-PQ-object-tagging.

If H_e in (10.1) is independent of $H_{u,d}$, additional leptonic signatures may arise at LEP (depending on the mass of H_e) and/or at the LHC (see [198]).

Finally, note that in order to write the flavor diagonal couplings of $H_{u,d}$ to first generation quarks, we implicitly assumed an MFV-type mechanism which generates the CKM flavor structure of the weak interactions of quarks without spoiling the flavor alignment of Yukawa couplings. Concrete realizations of such mechanism are possible, but are beyond the scope of this study.

11 Discussion

A short-lived, *pion-phobic* QCD axion with mass of several MeV might still offer a viable solution to the strong CP problem. Constraints that have excluded generic MeV axions can be evaded by coupling the axion exclusively to the first generation of SM fermions. Bounds from $K^+ \rightarrow \pi^+ a$, previously believed to be severe, in fact suffer from large hadronic uncertainties and are currently sufficiently ambiguous to experimentally allow portions of the axion parameter space.

¹⁵See also [238, 239] for exotic Z and Higgs decays to light ALPs.

The extreme *pion-phobia* needed to avoid exclusion is a realistic possibility in models with a special relation between the light quark masses and PQ charges, namely,

$$\frac{m_u}{m_d} \simeq \frac{Q_d^{\text{PQ}}}{Q_u^{\text{PQ}}}.$$

In this study we have imposed this relation *ad hoc*, but it is easy to envision how it might arise dynamically. For instance, in supersymmetric models of flavor, quartics are often proportional to charges squared, and flavon VEVs (and thus fermions Yukawas) would then naturally be inversely proportional to the charges.

The associated phenomenology of these variants is rich and testable. Several axion signatures are similar to the those of visibly decaying dark photons, and can be searched for by ongoing and near-future dark photon experiments. They also offer alternative explanations to a few discrepancies in data usually attributed to dark photons, such as the Beryllium-8 and KTeV ($\pi^0 \rightarrow e^+e^-$) anomalies. At the GeV scale, these models predict new states coupled to light hadrons awaiting to be uncovered. Those, along with the MeV axion, may appear in rare decays of the SM Higgs, Z^0 and other BSM states, yielding exotic signatures with thin jets (i.e., “ τ_h -like”), prompt or displaced lepton jets, and photon jets.

Acknowledgments

We thank these colleagues for helpful discussions on various aspects of this study: Wolfgang Altmannshofer, Vincenzo Cirigliano, Kyle Cranmer, Lance Dixon, Bertrand Echenart, Daniel Egaña, Glennys Farrar, Alex Friedland, Greg Gabadadze, Michael Graesser, Andy Haas, Eva Halkiadakis, Yonit Hochberg, Anson Hook, Yoni Kahn, Attila Krasnahorkey, Eric Kuflik, Anthony Palladino, Duccio Pappadopulo, Maxim Pospelov, Josh Ruderman, Philip Schuster, Tim Tait, Flip Tanedo, Natalia Toro, Chris Tully, Ken Van Tilberg, and Graziano Venanzoni.

N.W. and D.S.M.A. were supported by the NSF under grants PHY-0947827 and PHY-1316753. At earlier stages of this work, D.S.M.A. was also supported supported by NSF-PHY-0969510 (the LHC Theory Initiative). D.S.M.A. is currently supported by LANL’s Early Career LDRD program. D.S.M.A. thanks the Aspen Center for Physics (NSF Grant No. PHYS-1066293) for hospitality while progress was made on this study.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] R.D. Peccei and H.R. Quinn, *CP Conservation in the Presence of Instantons*, *Phys. Rev. Lett.* **38** (1977) 1440 [[INSPIRE](#)].
- [2] R.D. Peccei and H.R. Quinn, *Constraints Imposed by CP Conservation in the Presence of Instantons*, *Phys. Rev.* **D 16** (1977) 1791 [[INSPIRE](#)].
- [3] S. Weinberg, *A New Light Boson?*, *Phys. Rev. Lett.* **40** (1978) 223 [[INSPIRE](#)].

- [4] F. Wilczek, *Problem of Strong p and t Invariance in the Presence of Instantons*, *Phys. Rev. Lett.* **40** (1978) 279 [INSPIRE].
- [5] J.E. Kim, *Light Pseudoscalars, Particle Physics and Cosmology*, *Phys. Rept.* **150** (1987) 1 [INSPIRE].
- [6] H.-Y. Cheng, *The Strong CP Problem Revisited*, *Phys. Rept.* **158** (1988) 1 [INSPIRE].
- [7] J.E. Kim and G. Carosi, *Axions and the Strong CP Problem*, *Rev. Mod. Phys.* **82** (2010) 557 [arXiv:0807.3125] [INSPIRE].
- [8] C. Hagmann, H. Murayama, G.G. Raffelt, L.J. Rosenberg and K. van Bibber, *Axions*, *Phys. Lett. B* **667** (2008) 459.
- [9] D.J.E. Marsh, *Axion Cosmology*, *Phys. Rept.* **643** (2016) 1 [arXiv:1510.07633] [INSPIRE].
- [10] PARTICLE DATA GROUP collaboration, C. Patrignani et al., *Review of Particle Physics*, *Chin. Phys. C* **40** (2016) 100001.
- [11] R.D. Peccei, *The Strong CP Problem*, *Adv. Ser. Direct. High Energy Phys.* **3** (1989) 503.
- [12] M.S. Turner, *Windows on the Axion*, *Phys. Rept.* **197** (1990) 67 [INSPIRE].
- [13] G.G. Raffelt, *Astrophysical methods to constrain axions and other novel particle phenomena*, *Phys. Rept.* **198** (1990) 1 [INSPIRE].
- [14] J. Engel, D. Seckel and A.C. Hayes, *Emission and detectability of hadronic axions from SN1987A*, *Phys. Rev. Lett.* **65** (1990) 960 [INSPIRE].
- [15] G.G. Raffelt, *Astrophysical axion bounds*, *Lect. Notes Phys.* **741** (2008) 51 [hep-ph/0611350] [INSPIRE].
- [16] M. Millea, L. Knox and B. Fields, *New Bounds for Axions and Axion-Like Particles with keV-GeV Masses*, *Phys. Rev. D* **92** (2015) 023010 [arXiv:1501.04097] [INSPIRE].
- [17] J.E. Kim, *Weak Interaction Singlet and Strong CP Invariance*, *Phys. Rev. Lett.* **43** (1979) 103 [INSPIRE].
- [18] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Can Confinement Ensure Natural CP Invariance of Strong Interactions?*, *Nucl. Phys. B* **166** (1980) 493 [INSPIRE].
- [19] M. Dine, W. Fischler and M. Srednicki, *A Simple Solution to the Strong CP Problem with a Harmless Axion*, *Phys. Lett.* **104B** (1981) 199 [INSPIRE].
- [20] A.R. Zhitnitsky, *On Possible Suppression of the Axion Hadron Interactions. (In Russian)*, *Sov. J. Nucl. Phys.* **31** (1980) 260 [INSPIRE].
- [21] P. Sikivie, *Detection Rates for 'Invisible' Axion Searches*, *Phys. Rev. D* **32** (1985) 2988 [Erratum *ibid.* **D 36** (1987) 974] [INSPIRE].
- [22] H. Peng et al., *Cryogenic cavity detector for a large scale cold dark-matter axion search*, *Nucl. Instrum. Meth. A* **444** (2000) 569 [INSPIRE].
- [23] G. Mueller, P. Sikivie, D.B. Tanner and K. van Bibber, *Detailed design of a resonantly-enhanced axion-photon regeneration experiment*, *Phys. Rev. D* **80** (2009) 072004 [arXiv:0907.5387] [INSPIRE].
- [24] ADMX collaboration, S.J. Asztalos et al., *A SQUID-based microwave cavity search for dark-matter axions*, *Phys. Rev. Lett.* **104** (2010) 041301 [arXiv:0910.5914] [INSPIRE].

- [25] R. Bähre et al., *Any light particle search II — Technical Design Report, 2013 JINST* **8** T09001 [[arXiv:1302.5647](#)] [[INSPIRE](#)].
- [26] C.E. Aalseth et al., *The CERN Axion Solar Telescope (CAST), Nucl. Phys. Proc. Suppl.* **110** (2002) 85 [[INSPIRE](#)].
- [27] I.G. Irastorza et al., *Towards a new generation axion helioscope, JCAP* **06** (2011) 013 [[arXiv:1103.5334](#)] [[INSPIRE](#)].
- [28] E. Armengaud et al., *Conceptual Design of the International Axion Observatory (IAXO), 2014 JINST* **9** T05002 [[arXiv:1401.3233](#)] [[INSPIRE](#)].
- [29] D. Budker, P.W. Graham, M. Ledbetter, S. Rajendran and A. Sushkov, *Proposal for a Cosmic Axion Spin Precession Experiment (CASPER), Phys. Rev. X* **4** (2014) 021030 [[arXiv:1306.6089](#)] [[INSPIRE](#)].
- [30] Y. Kahn, B.R. Safdi and J. Thaler, *Broadband and Resonant Approaches to Axion Dark Matter Detection, Phys. Rev. Lett.* **117** (2016) 141801 [[arXiv:1602.01086](#)] [[INSPIRE](#)].
- [31] MADMAX WORKING GROUP collaboration, A. Caldwell et al., *Dielectric Haloscopes: A New Way to Detect Axion Dark Matter, Phys. Rev. Lett.* **118** (2017) 091801 [[arXiv:1611.05865](#)] [[INSPIRE](#)].
- [32] P.W. Graham, I.G. Irastorza, S.K. Lamoreaux, A. Lindner and K.A. van Bibber, *Experimental Searches for the Axion and Axion-Like Particles, Ann. Rev. Nucl. Part. Sci.* **65** (2015) 485 [[arXiv:1602.00039](#)] [[INSPIRE](#)].
- [33] J. Preskill, M.B. Wise and F. Wilczek, *Cosmology of the Invisible Axion, Phys. Lett. B* **120** (1983) 127 [[INSPIRE](#)].
- [34] L.F. Abbott and P. Sikivie, *A Cosmological Bound on the Invisible Axion, Phys. Lett. B* **120** (1983) 133 [[INSPIRE](#)].
- [35] M. Dine and W. Fischler, *The Not So Harmless Axion, Phys. Lett. B* **120** (1983) 137 [[INSPIRE](#)].
- [36] F. Wilczek, *Decays of Heavy Vector Mesons Into Higgs Particles, Phys. Rev. Lett.* **39** (1977) 1304 [[INSPIRE](#)].
- [37] P. Nason, *QCD Radiative Corrections to Υ Decay Into Scalar Plus γ and Pseudoscalar Plus γ , Phys. Lett. B* **175** (1986) 223 [[INSPIRE](#)].
- [38] ARGUS collaboration, H. Albrecht et al., *Search for Exotic Decay Modes of the Υ ($1s$), Phys. Lett. B* **179** (1986) 403 [[INSPIRE](#)].
- [39] S.Y. Hsueh and S. Palestini, *The Width of the J/ψ resonance, Phys. Rev. D* **45** (1992) R2181.
- [40] E760 collaboration, T.A. Armstrong et al., *Measurement of the J/ψ and ψ -prime resonance parameters in $\bar{p}p$ annihilation, Phys. Rev. D* **47** (1993) 772 [[INSPIRE](#)].
- [41] BES collaboration, J.Z. Bai et al., *A Measurement of J/ψ decay widths, Phys. Lett. B* **355** (1995) 374 [*Erratum ibid.* **B 363** (1995) 267] [[INSPIRE](#)].
- [42] CLEO collaboration, G.S. Adams et al., *Measurement of $\Gamma(ee)(J/\psi)$, $\Gamma(\text{tot})(J/\psi)$ and $\Gamma(ee)[\psi(2S)]/\Gamma(ee)(J/\psi)$, Phys. Rev. D* **73** (2006) 051103 [[hep-ex/0512046](#)] [[INSPIRE](#)].
- [43] M. Davier, *Searches for New Particles, in Proceedings of 23RD International Conference on High Energy Physics, Berkeley U.S.A. (1986).*

- [44] CHARM collaboration, F. Bergsma et al., *Search for Axion Like Particle Production in 400-GeV Proton-Copper Interactions*, *Phys. Lett.* **157B** (1985) 458 [INSPIRE].
- [45] J.D. Bjorken et al., *Search for Neutral Metastable Penetrating Particles Produced in the SLAC Beam Dump*, *Phys. Rev. D* **38** (1988) 3375 [INSPIRE].
- [46] M. Davier and H. Nguyen Ngoc, *An Unambiguous Search for a Light Higgs Boson*, *Phys. Lett. B* **229** (1989) 150 [INSPIRE].
- [47] J. Blumlein et al., *Limits on neutral light scalar and pseudoscalar particles in a proton beam dump experiment*, *Z. Phys. C* **51** (1991) 341 [INSPIRE].
- [48] E787 collaboration, S.S. Adler et al., *Search for the decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ in the momentum region $P(\pi)$ less than 195-MeV/c*, *Phys. Lett. B* **537** (2002) 211 [hep-ex/0201037] [INSPIRE].
- [49] E787 collaboration, S. Adler et al., *Further search for the decay $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ in the momentum region $P < 195$ -MeV/c*, *Phys. Rev. D* **70** (2004) 037102 [hep-ex/0403034] [INSPIRE].
- [50] E949 collaboration, V.V. Anisimovsky et al., *Improved measurement of the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ branching ratio*, *Phys. Rev. Lett.* **93** (2004) 031801 [hep-ex/0403036] [INSPIRE].
- [51] L.M. Krauss and D.J. Nash, *A Viable weak interaction axion?*, *Phys. Lett. B* **202** (1988) 560 [INSPIRE].
- [52] L.M. Krauss and M.B. Wise, *Constraints on Shortlived Axions From the Decay $\pi^+ \rightarrow e^+ e^- e^+$ Neutrino*, *Phys. Lett. B* **176** (1986) 483 [INSPIRE].
- [53] SINDRUM collaboration, R. Eichler et al., *Limits for Shortlived Neutral Particles Emitted μ^+ or π^+ Decay*, *Phys. Lett. B* **175** (1986) 101 [INSPIRE].
- [54] I. Antoniadis and T.N. Truong, *Lower Bound for Branching Ratio of $K^+ \rightarrow \pi^+$ Axion and Nonexistence of Peccei-Quinn Axion*, *Phys. Lett.* **109B** (1982) 67 [INSPIRE].
- [55] W.A. Bardeen, R.D. Peccei and T. Yanagida, *Constraints on variant axion models*, *Nucl. Phys. B* **279** (1987) 401 [INSPIRE].
- [56] KTeV collaboration, E. Abouzaid et al., *Measurement of the Rare Decay $\pi^0 \rightarrow e^+ e^-$* , *Phys. Rev. D* **75** (2007) 012004 [hep-ex/0610072] [INSPIRE].
- [57] A.E. Dorokhov and M.A. Ivanov, *Rare decay $\pi^0 \rightarrow e^+ e^-$: Theory confronts KTeV data*, *Phys. Rev. D* **75** (2007) 114007 [arXiv:0704.3498] [INSPIRE].
- [58] A.E. Dorokhov, E.A. Kuraev, Yu. M. Bystritskiy and M. Secansky, *QED radiative corrections to the decay $\pi^0 \rightarrow e^+ e^-$* , *Eur. Phys. J. C* **55** (2008) 193 [arXiv:0801.2028] [INSPIRE].
- [59] A.E. Dorokhov, M.A. Ivanov and S.G. Kovalenko, *Complete structure dependent analysis of the decay $P \rightarrow \ell^+ \ell^-$* , *Phys. Lett. B* **677** (2009) 145 [arXiv:0903.4249] [INSPIRE].
- [60] P. Vasko and J. Novotny, *Two-loop QED radiative corrections to the decay $\pi^0 \rightarrow e^+ e^-$: The virtual corrections and soft-photon bremsstrahlung*, *JHEP* **10** (2011) 122 [arXiv:1106.5956] [INSPIRE].
- [61] T. Husek, K. Kampf and J. Novotný, *Rare decay $\pi^0 \rightarrow e^+ e^-$: on corrections beyond the leading order*, *Eur. Phys. J. C* **74** (2014) 3010 [arXiv:1405.6927] [INSPIRE].
- [62] L.G. Landsberg, *Electromagnetic Decays of Light Mesons*, *Phys. Rept.* **128** (1985) 301 [INSPIRE].

- [63] L. Bergstrom, *Radiative corrections to pseudoscalar meson decays*, *Z. Phys. C* **20** (1983) 135 [INSPIRE].
- [64] Y. Kahn, M. Schmitt and T.M.P. Tait, *Enhanced rare pion decays from a model of MeV dark matter*, *Phys. Rev. D* **78** (2008) 115002 [arXiv:0712.0007] [INSPIRE].
- [65] T. Yamazaki et al., *Search for a neutral boson in a two-body decay of $K^+ \rightarrow \pi + X0$* , *Phys. Rev. Lett.* **52** (1984) 1089 [INSPIRE].
- [66] N.J. Baker et al., *Search for Shortlived Neutral Particles Emitted in K^+ Decay*, *Phys. Rev. Lett.* **59** (1987) 2832 [INSPIRE].
- [67] Y. Asano et al., *A New Experimental Limit for the Decay $K^+ \rightarrow \pi^+ \gamma \gamma$* , *Phys. Lett. B* **113** (1982) 195 [INSPIRE].
- [68] R.J. Abrams et al., *New Experimental Upper Limit for the Decay $K^\pm \rightarrow \pi^\pm \gamma \Gamma$* , *Phys. Rev. D* **15** (1977) 22 [INSPIRE].
- [69] T. Yamazaki, *Search for Exotics in Two-Body Decay of K^+ : $K(\mu 2)$ and $K(\pi 2)$ Revisited*, in *Proceedings of the Second LAMPF II Workshop. Vol. II*, Los Alamos U.S.A. (1982), Preprint LA-9572-C, pg. 413.
- [70] J.F. Donoghue, E. Golowich and B.R. Holstein, *Dynamics of the standard model*, *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.* **2** (1992) 1 [Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **35** (2014) 1].
- [71] V. Cirigliano, G. Ecker, H. Neufeld, A. Pich and J. Portoles, *Kaon Decays in the Standard Model*, *Rev. Mod. Phys.* **84** (2012) 399 [arXiv:1107.6001] [INSPIRE].
- [72] J.M. Gerard and J. Weyers, *Trace anomalies and the $\Delta I = 1/2$ rule*, *Phys. Lett. B* **503** (2001) 99 [hep-ph/0011391] [INSPIRE].
- [73] A.J. Buras, J.-M. Gérard and W.A. Bardeen, *Large N Approach to Kaon Decays and Mixing 28 Years Later: $\Delta I = 1/2$ Rule, \hat{B}_K and ΔM_K* , *Eur. Phys. J. C* **74** (2014) 2871 [arXiv:1401.1385] [INSPIRE].
- [74] R.J. Crewther and L.C. Tunstall, *$\Delta I = 1/2$ rule for kaon decays derived from QCD infrared fixed point*, *Phys. Rev. D* **91** (2015) 034016 [arXiv:1312.3319] [INSPIRE].
- [75] J. Kambor, J.H. Missimer and D. Wyler, *$K \rightarrow 2\pi$ and $K \rightarrow 3\pi$ decays in next-to-leading order chiral perturbation theory*, *Phys. Lett. B* **261** (1991) 496 [INSPIRE].
- [76] J. Bijnens, P. Dhonte and F. Borg, *$K \rightarrow 3\pi$ decays in chiral perturbation theory*, *Nucl. Phys. B* **648** (2003) 317 [hep-ph/0205341] [INSPIRE].
- [77] H. Georgi, *A bound on $m(\eta)/m(\eta')$ for large N_c* , *Phys. Rev. D* **49** (1994) 1666 [hep-ph/9310337] [INSPIRE].
- [78] J.M. Gerard and E. Kou, *$\eta - \eta'$ masses and mixing: A Large N_c reappraisal*, *Phys. Lett. B* **616** (2005) 85 [hep-ph/0411292] [INSPIRE].
- [79] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory to One Loop*, *Annals Phys.* **158** (1984) 142 [INSPIRE].
- [80] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory: Expansions in the Mass of the Strange Quark*, *Nucl. Phys. B* **250** (1985) 465 [INSPIRE].
- [81] H. Leutwyler, *On the $1/N$ expansion in chiral perturbation theory*, *Nucl. Phys. Proc. Suppl.* **64** (1998) 223 [hep-ph/9709408] [INSPIRE].

- [82] B. Borasoy and S. Wetzel, *U(3) chiral perturbation theory with infrared regularization*, *Phys. Rev. D* **63** (2001) 074019 [[hep-ph/0105132](#)] [[INSPIRE](#)].
- [83] W.A. Bardeen and S.H.H. Tye, *Current Algebra Applied to Properties of the Light Higgs Boson*, *Phys. Lett.* **74B** (1978) 229 [[INSPIRE](#)].
- [84] E949 collaboration, A.V. Artamonov et al., *Search for the decay $K^+ \rightarrow \pi^+ \gamma \gamma$ in the π^+ momentum region $P > 213 \text{ MeV}/c$* , *Phys. Lett.* **B 623** (2005) 192 [[hep-ex/0505069](#)] [[INSPIRE](#)].
- [85] S.B. Treiman and F. Wilczek, *Axion Emission in Decay of Excited Nuclear States*, *Phys. Lett.* **74B** (1978) 381 [[INSPIRE](#)].
- [86] T.W. Donnelly, S.J. Freedman, R.S. Lytel, R.D. Peccei and M. Schwartz, *Do Axions Exist?*, *Phys. Rev. D* **18** (1978) 1607 [[INSPIRE](#)].
- [87] A. Barroso and N.C. Mukhopadhyay, *Nuclear axion decay*, *Phys. Rev. C* **24** (1981) 2382 [[INSPIRE](#)].
- [88] R.L. Jaffe and A. Manohar, *The G(1) Problem: Fact and Fantasy on the Spin of the Proton*, *Nucl. Phys.* **B 337** (1990) 509 [[INSPIRE](#)].
- [89] M.J. Savage and J. Walden, *SU(3) breaking in neutral current axial matrix elements and the spin content of the nucleon*, *Phys. Rev. D* **55** (1997) 5376 [[hep-ph/9611210](#)] [[INSPIRE](#)].
- [90] M. Karliner and H.J. Lipkin, *Nucleon spin with and without hyperon data: A New tool for analysis*, *Phys. Lett.* **B 461** (1999) 280 [[hep-ph/9906321](#)] [[INSPIRE](#)].
- [91] G.K. Mallot, *The Spin structure of the nucleon*, *Int. J. Mod. Phys. A* **15S1** (2000) 521 [[hep-ex/9912040](#)] [[INSPIRE](#)].
- [92] E. Leader, A.V. Sidorov and D.B. Stamenov, *On the sensitivity of the polarized parton densities to flavor SU(3) symmetry breaking*, *Phys. Lett.* **B 488** (2000) 283 [[hep-ph/0004106](#)] [[INSPIRE](#)].
- [93] H.-Y. Cheng and C.-W. Chiang, *Revisiting Scalar and Pseudoscalar Couplings with Nucleons*, *JHEP* **07** (2012) 009 [[arXiv:1202.1292](#)] [[INSPIRE](#)].
- [94] QCDSF collaboration, G.S. Bali et al., *Strangeness Contribution to the Proton Spin from Lattice QCD*, *Phys. Rev. Lett.* **108** (2012) 222001 [[arXiv:1112.3354](#)] [[INSPIRE](#)].
- [95] M. Engelhardt, *Strange quark contributions to nucleon mass and spin from lattice QCD*, *Phys. Rev. D* **86** (2012) 114510 [[arXiv:1210.0025](#)] [[INSPIRE](#)].
- [96] A. Abdel-Rehim et al., *Disconnected quark loop contributions to nucleon observables in lattice QCD*, *Phys. Rev. D* **89** (2014) 034501 [[arXiv:1310.6339](#)] [[INSPIRE](#)].
- [97] T. Bhattacharya, R. Gupta and B. Yoon, *Disconnected Quark Loop Contributions to Nucleon Structure*, *PoS(LATTICE2014)* 141 [[arXiv:1503.05975](#)] [[INSPIRE](#)].
- [98] A. Abdel-Rehim et al., *Nucleon and pion structure with lattice QCD simulations at physical value of the pion mass*, *Phys. Rev. D* **92** (2015) 114513 [[arXiv:1507.04936](#)] [[INSPIRE](#)].
- [99] A. Abdel-Rehim et al., *Disconnected quark loop contributions to nucleon observables using $N_f = 2$ twisted clover fermions at the physical value of the light quark mass*, *PoS(LATTICE 2015)* 136 [[arXiv:1511.00433](#)] [[INSPIRE](#)].
- [100] J. Green et al., *Up, down and strange nucleon axial form factors from lattice QCD*, *Phys. Rev. D* **95** (2017) 114502 [[arXiv:1703.06703](#)] [[INSPIRE](#)].

- [101] G. Grilli di Cortona, E. Hardy, J. Pardo Vega and G. Villadoro, *The QCD axion, precisely*, *JHEP* **01** (2016) 034 [[arXiv:1511.02867](#)] [[INSPIRE](#)].
- [102] M.J. Savage, B.W. Filippone and L.W. Mitchell, *New Limits on Light Scalar and Pseudoscalar Particles Produced in Nuclear Decay*, *Phys. Rev. D* **37** (1988) 1134 [[INSPIRE](#)].
- [103] F.P. Calaprice et al., *Search for axion emission in the decay of excited states of C-12*, *Phys. Rev. D* **20** (1979) 2708 [[INSPIRE](#)].
- [104] M.J. Savage, R.D. Mckeown, B.W. Filippone and L.W. Mitchell, *Search for a Shortlived Neutral Particle Produced in Nuclear Decay*, *Phys. Rev. Lett.* **57** (1986) 178 [[INSPIRE](#)].
- [105] A.L. Hallin, F.P. Calaprice, R.W. Dunford and A.B. McDonald, *Restrictions on a 1.7-MeV Axion From Nuclear Pair Transitions*, *Phys. Rev. Lett.* **57** (1986) 2105 [[INSPIRE](#)].
- [106] F.W.N. De Boer et al., *Search for Shortlived Axions in a Nuclear Isoscalar Transition*, *Phys. Lett. B* **180** (1986) 4 [[INSPIRE](#)].
- [107] A.L. Hallin, F.P. Calaprice, R.W. Dunford, A.B. McDonald and G.C. Paulson, *Limits on axions from nuclear decays*, *Nucl. Instrum. Meth. B* **24-25** (1987) 276.
- [108] V.M. Datar, S. Fortier, S. Gales, E. Hourani, H. Langevin, J.M. Maison et al., *Search for short lived neutral particle in the 15.1-MeV isovector transition of C-12*, *Phys. Rev. C* **37** (1988) 250 [[INSPIRE](#)].
- [109] S. Freedman, *Search for shortlived axions*, *Nucl. Instrum. Meth. A* **284** (1989) 50 [[INSPIRE](#)].
- [110] T. Asanuma, M. Minowa, T. Tsukamoto, S. Orito and T. Tsunoda, *A Search for correlated e^+e^- pairs in the decay of AM-241*, *Phys. Lett. B* **237** (1990) 588 [[INSPIRE](#)].
- [111] F.W.N. de Boer, J. Lehmann and J. Steyaert, *Search for a short lived isoscalar axion with mass between 4-GeV/c^{**(-2)} and 15-MeV/c^{**(-2)}*, *J. Phys. G* **16** (1990) L1 [[INSPIRE](#)].
- [112] F.W.N. de Boer et al., *A deviation in internal pair conversion*, *Phys. Lett. B* **388** (1996) 235 [[INSPIRE](#)].
- [113] F.W.N. de Boer and J. Steyaert, *Comment on a search for a short lived isoscalar axion with mass between 4-MeV/c^{**2} and 15-MeV/c^{**2}*, *J. Phys. G* **23** (1997) 729 [[INSPIRE](#)].
- [114] A.J. Krasznahorkay et al., *Observation of Anomalous Internal Pair Creation in Be8: A Possible Indication of a Light, Neutral Boson*, *Phys. Rev. Lett.* **116** (2016) 042501 [[arXiv:1504.01527](#)] [[INSPIRE](#)].
- [115] X. Zhang and G.A. Miller, *Can nuclear physics explain the anomaly observed in the internal pair production in the Beryllium-8 nucleus?*, *Phys. Lett. B* **773** (2017) 159 [[arXiv:1703.04588](#)] [[INSPIRE](#)].
- [116] J.L. Feng et al., *Protophobic Fifth-Force Interpretation of the Observed Anomaly in ⁸Be Nuclear Transitions*, *Phys. Rev. Lett.* **117** (2016) 071803 [[arXiv:1604.07411](#)] [[INSPIRE](#)].
- [117] P.-H. Gu and X.-G. He, *Realistic model for a fifth force explaining anomaly in ⁸Be* → ⁸Be e⁺e⁻ Decay*, *Nucl. Phys. B* **919** (2017) 209 [[arXiv:1606.05171](#)] [[INSPIRE](#)].
- [118] J.L. Feng et al., *Particle physics models for the 17 MeV anomaly in beryllium nuclear decays*, *Phys. Rev. D* **95** (2017) 035017 [[arXiv:1608.03591](#)] [[INSPIRE](#)].
- [119] U. Ellwanger and S. Moretti, *Possible Explanation of the Electron Positron Anomaly at 17 MeV in ⁸Be Transitions Through a Light Pseudoscalar*, *JHEP* **11** (2016) 039 [[arXiv:1609.01669](#)] [[INSPIRE](#)].

- [120] Y. Kahn, G. Krnjaic, S. Mishra-Sharma and T.M.P. Tait, *Light Weakly Coupled Axial Forces: Models, Constraints and Projections*, *JHEP* **05** (2017) 002 [[arXiv:1609.09072](#)] [[INSPIRE](#)].
- [121] P. Fayet, *The light U boson as the mediator of a new force, coupled to a combination of Q, B, L and dark matter*, *Eur. Phys. J. C* **77** (2017) 53 [[arXiv:1611.05357](#)] [[INSPIRE](#)].
- [122] J. Kozaczuk, D.E. Morrissey and S.R. Stroberg, *Light axial vector bosons, nuclear transitions and the ^8Be anomaly*, *Phys. Rev. D* **95** (2017) 115024 [[arXiv:1612.01525](#)] [[INSPIRE](#)].
- [123] L. Delle Rose, S. Khalil and S. Moretti, *Explanation of the 17 MeV Atomki anomaly in a U(1)'-extended two Higgs doublet model*, *Phys. Rev. D* **96** (2017) 115024 [[arXiv:1704.03436](#)] [[INSPIRE](#)].
- [124] A.J. Krasznahorkay et al., *New experimental results for the 17 MeV particle created in ^8Be* , *EPJ Web Conf.* **137** (2017) 08010 [[INSPIRE](#)].
- [125] A.J. Krasznahorkay et al., *On the creation of the 17 MeV X boson in the 17.6 MeV M1 transition of ^8Be* , *EPJ Web Conf.* **142** (2017) 01019 [[INSPIRE](#)].
- [126] S. Pastore, R.B. Wiringa, S.C. Pieper and R. Schiavilla, *Quantum Monte Carlo calculations of electromagnetic transitions in ^8Be with meson-exchange currents derived from chiral effective field theory*, *Phys. Rev. C* **90** (2014) 024321 [[arXiv:1406.2343](#)] [[INSPIRE](#)].
- [127] A. Bross, M. Crisler, S.H. Pordes, J. Volk, S. Errede and J. Wrbanek, *A Search for Shortlived Particles Produced in an Electron Beam Dump*, *Phys. Rev. Lett.* **67** (1991) 2942 [[INSPIRE](#)].
- [128] E.M. Riordan et al., *A Search for Short Lived Axions in an Electron Beam Dump Experiment*, *Phys. Rev. Lett.* **59** (1987) 755 [[INSPIRE](#)].
- [129] M. Davier, J. Jeanjean and H. Nguyen Ngoc, *Search for Axions in Electron Bremsstrahlung*, *Phys. Lett. B* **180** (1986) 295 [[INSPIRE](#)].
- [130] BABAR collaboration, J.P. Lees et al., *Search for a Dark Photon in e^+e^- Collisions at BaBar*, *Phys. Rev. Lett.* **113** (2014) 201801 [[arXiv:1406.2980](#)] [[INSPIRE](#)].
- [131] A. Anastasi et al., *Limit on the production of a low-mass vector boson in $e^+e^- \rightarrow U\gamma$, $U \rightarrow e^+e^-$ with the KLOE experiment*, *Phys. Lett. B* **750** (2015) 633 [[arXiv:1509.00740](#)] [[INSPIRE](#)].
- [132] D.J. Bechis et al., *Search for Axion Production in Low-energy Electron Bremsstrahlung*, *Phys. Rev. Lett.* **42** (1979) 1511 [Erratum *ibid.* **43** (1979) 90] [[INSPIRE](#)].
- [133] A. Rothenberg, *A search for unknown sources of neutrino like particles*, Ph.D. Thesis, SLAC, Stanford U.S.A. (1972).
- [134] C.N. Brown et al., *A New Limit on Axion Production in 800-GeV Hadronic Showers*, *Phys. Rev. Lett.* **57** (1986) 2101 [[INSPIRE](#)].
- [135] R. Guo et al., *Improved Limit on Axion Production in 800-GeV Hadronic Showers*, *Phys. Rev. D* **41** (1990) 2924 [[INSPIRE](#)].
- [136] A. Konaka et al., *Search for Neutral Particles in Electron Beam Dump Experiment*, *Phys. Rev. Lett.* **57** (1986) 659 [[INSPIRE](#)].

- [137] T. Aoyama, M. Hayakawa, T. Kinoshita and M. Nio, *Tenth-Order Electron Anomalous Magnetic Moment — Contribution of Diagrams without Closed Lepton Loops*, *Phys. Rev. D* **91** (2015) 033006 [Erratum *ibid.* **D 96** (2017) 019901] [[arXiv:1412.8284](#)] [[INSPIRE](#)].
- [138] D. Hanneke, S. Fogwell and G. Gabrielse, *New Measurement of the Electron Magnetic Moment and the Fine Structure Constant*, *Phys. Rev. Lett.* **100** (2008) 120801 [[arXiv:0801.1134](#)] [[INSPIRE](#)].
- [139] D. Hanneke, S.F. Hoogerheide and G. Gabrielse, *Cavity Control of a Single-Electron Quantum Cyclotron: Measuring the Electron Magnetic Moment*, *Phys. Rev. A* **83** (2011) 052122 [[arXiv:1009.4831](#)] [[INSPIRE](#)].
- [140] W.J. Marciano, A. Masiero, P. Paradisi and M. Passera, *Contributions of axionlike particles to lepton dipole moments*, *Phys. Rev. D* **94** (2016) 115033 [[arXiv:1607.01022](#)] [[INSPIRE](#)].
- [141] K. Melnikov and A. Vainshtein, *Theory of the muon anomalous magnetic moment*, *Springer Tracts Mod. Phys.* **216** (2006) 1.
- [142] D. Stöckinger, *The Muon Magnetic Moment and Supersymmetry*, *J. Phys. G* **34** (2007) R45 [[hep-ph/0609168](#)] [[INSPIRE](#)].
- [143] G.F. Giudice, P. Paradisi and M. Passera, *Testing new physics with the electron $g-2$* , *JHEP* **11** (2012) 113 [[arXiv:1208.6583](#)] [[INSPIRE](#)].
- [144] R. Essig, P. Schuster and N. Toro, *Probing Dark Forces and Light Hidden Sectors at Low-Energy e^+e^- Colliders*, *Phys. Rev. D* **80** (2009) 015003 [[arXiv:0903.3941](#)] [[INSPIRE](#)].
- [145] J.D. Bjorken, R. Essig, P. Schuster and N. Toro, *New Fixed-Target Experiments to Search for Dark Gauge Forces*, *Phys. Rev. D* **80** (2009) 075018 [[arXiv:0906.0580](#)] [[INSPIRE](#)].
- [146] B. Batell, M. Pospelov and A. Ritz, *Probing a Secluded $U(1)$ at B -factories*, *Phys. Rev. D* **79** (2009) 115008 [[arXiv:0903.0363](#)] [[INSPIRE](#)].
- [147] B. Batell, M. Pospelov and A. Ritz, *Exploring Portals to a Hidden Sector Through Fixed Targets*, *Phys. Rev. D* **80** (2009) 095024 [[arXiv:0906.5614](#)] [[INSPIRE](#)].
- [148] J.L. Hewett et al., *Fundamental Physics at the Intensity Frontier*, [arXiv:1205.2671](#) [[INSPIRE](#)].
- [149] R. Essig et al., *Working Group Report: New Light Weakly Coupled Particles*, in *Proceedings of 2013 Community Summer Study on the Future of U.S. Particle Physics: Snowmass on the Mississippi (CSS2013)*, Minneapolis U.S.A. (2013) [[arXiv:1311.0029](#)] [[INSPIRE](#)].
- [150] S. Alekhin et al., *A facility to Search for Hidden Particles at the CERN SPS: the SHiP physics case*, *Rept. Prog. Phys.* **79** (2016) 124201 [[arXiv:1504.04855](#)] [[INSPIRE](#)].
- [151] J. Alexander et al., *Dark Sectors 2016 Workshop: Community Report*, 2016, [arXiv:1608.08632](#) [[INSPIRE](#)].
- [152] Y.-S. Tsai, *Axion bremsstrahlung by an electron beam*, *Phys. Rev. D* **34** (1986) 1326 [[INSPIRE](#)].
- [153] Y.-S. Tsai, *Pair Production and Bremsstrahlung of Charged Leptons*, *Rev. Mod. Phys.* **46** (1974) 815 [Erratum *ibid.* **49** (1977) 521] [[INSPIRE](#)].
- [154] K.J. Kim and Y.-S. Tsai, *Improved Weizsacker-Williams method and its application to lepton and w boson pair production*, *Phys. Rev. D* **8** (1973) 3109 [[INSPIRE](#)].
- [155] B. Wojtsekhowski, *Searching for a U -boson with a positron beam*, *AIP Conf. Proc.* **1160** (2009) 149 [[arXiv:0906.5265](#)] [[INSPIRE](#)].

- [156] B. Wojtsekhowski, D. Nikolenko and I. Rachek, *Searching for a new force at VEPP-3*, [arXiv:1207.5089](#) [[INSPIRE](#)].
- [157] M. Raggi and V. Kozhuharov, *Proposal to Search for a Dark Photon in Positron on Target Collisions at DAΦNE Linac*, *Adv. High Energy Phys.* **2014** (2014) 959802 [[arXiv:1403.3041](#)] [[INSPIRE](#)].
- [158] BELLE collaboration, I. Jaegle, *Search for the dark photon and the dark Higgs boson at Belle*, *Phys. Rev. Lett.* **114** (2015) 211801 [[arXiv:1502.00084](#)] [[INSPIRE](#)].
- [159] M. Battaglieri et al., *The Heavy Photon Search Test Detector*, *Nucl. Instrum. Meth. A* **777** (2015) 91 [[arXiv:1406.6115](#)] [[INSPIRE](#)].
- [160] J. Balewski et al., *DarkLight: A Search for Dark Forces at the Jefferson Laboratory Free-Electron Laser Facility*, in *Proceedings of 2013 Community Summer Study on the Future of U.S. Particle Physics: Snowmass on the Mississippi (CSS2013)*, Minneapolis U.S.A. (2013) [[arXiv:1307.4432](#)] [[INSPIRE](#)].
- [161] J. Balewski et al., *The DarkLight Experiment: A Precision Search for New Physics at Low Energies*, [[arXiv:1412.4717](#)] [[INSPIRE](#)].
- [162] A. Denig, *Recent results from the Mainz Microtron MAMI and an outlook for the future*, *AIP Conf. Proc.* **1735** (2016) 020006 [[INSPIRE](#)].
- [163] A. Schaefer, *New Particles in Strong Fields?*, *J. Phys. G* **15** (1989) 373 [[INSPIRE](#)].
- [164] A. Ishida et al., *New Precision Measurement of Hyperfine Splitting of Positronium*, *Phys. Lett. B* **734** (2014) 338 [[arXiv:1310.6923](#)] [[INSPIRE](#)].
- [165] R. Barbieri and T.E.O. Ericson, *Evidence Against the Existence of a Low Mass Scalar Boson from Neutron-Nucleus Scattering*, *Phys. Lett.* **57B** (1975) 270 [[INSPIRE](#)].
- [166] V.V. Nesvizhevsky, G. Pignol and K.V. Protasov, *Neutron scattering and extra short range interactions*, *Phys. Rev. D* **77** (2008) 034020 [[arXiv:0711.2298](#)] [[INSPIRE](#)].
- [167] Y. Kamyshkov, J. Tithof and M. Vysotsky, *Bounds on new light particles from high-energy and very small momentum transfer np elastic scattering data*, *Phys. Rev. D* **78** (2008) 114029 [[arXiv:0810.4653](#)] [[INSPIRE](#)].
- [168] R. Daido and F. Takahashi, *The sign of the dipole-dipole potential by axion exchange*, *Phys. Lett. B* **772** (2017) 127 [[arXiv:1704.00155](#)] [[INSPIRE](#)].
- [169] A.A. Poblaguev et al., *Experimental study of the radiative decays $K^+ \rightarrow \mu^+ \nu e^+ e^-$ and $K^+ \rightarrow e^+ \nu e^+ e^-$* , *Phys. Rev. Lett.* **89** (2002) 061803 [[hep-ex/0204006](#)] [[INSPIRE](#)].
- [170] NA48 collaboration, A. Lai et al., *Investigation of $K(L, S) \rightarrow \pi^+ \pi^- e^+ e^-$ decays*, *Eur. Phys. J. C* **30** (2003) 33 [[INSPIRE](#)].
- [171] KTeV collaboration, J. Adams et al., *Measurement of the branching fraction of the decay $K(L) \rightarrow \pi^+ \pi^- e^+ e^-$* , *Phys. Rev. Lett.* **80** (1998) 4123 [[INSPIRE](#)].
- [172] KTeV collaboration, A. Alavi-Harati et al., *Search for the $K(L) \rightarrow \pi^0 \pi^0 e^+ e^-$ decay in the KTeV experiment*, *Phys. Rev. Lett.* **89** (2002) 211801 [[hep-ex/0210056](#)] [[INSPIRE](#)].
- [173] NA48/1 collaboration, J.R. Batley et al., *Observation of the rare decay $K(S) \rightarrow \pi^0 e^+ e^-$* , *Phys. Lett. B* **576** (2003) 43 [[hep-ex/0309075](#)] [[INSPIRE](#)].
- [174] NA48 collaboration, A. Lai et al., *Search for the decay $K(S) \rightarrow \pi^0 e^+ e^-$* , *Phys. Lett. B* **514** (2001) 253 [[INSPIRE](#)].

- [175] NA31 collaboration, G.D. Barr et al., *Search for the decay $K(S) \rightarrow \pi^0 e^+ e^-$. CERN-NA031 experiment*, *Phys. Lett. B* **304** (1993) 381 [INSPIRE].
- [176] CELSIUS/WASA collaboration, C. Bargholtz et al., *Measurement of the $\eta \rightarrow \pi^+ \pi^- e^+ e^-$ decay branching ratio*, *Phys. Lett. B* **644** (2007) 299 [hep-ex/0609007] [INSPIRE].
- [177] CMD-2 collaboration, R.R. Akhmetshin et al., *Study of conversion decays $\phi \rightarrow \eta e^+ e^-$, $\eta \rightarrow e^+ e^- \gamma$ and $\eta \rightarrow \pi^+ \pi^- e^+ e^-$ at CMD-2*, *Phys. Lett. B* **501** (2001) 191 [hep-ex/0012039] [INSPIRE].
- [178] CELSIUS/WASA collaboration, M. Berlowski et al., *Measurement of eta meson decays into lepton-antilepton pairs*, *Phys. Rev. D* **77** (2008) 032004 [arXiv:0711.3531] [INSPIRE].
- [179] KLOE collaboration, F. Ambrosino et al., *Measurement of the branching ratio and search for a CP-violating asymmetry in the $\eta \rightarrow \pi^+ \pi^- e^+ e^- (\gamma)$ decay at KLOE*, *Phys. Lett. B* **675** (2009) 283 [arXiv:0812.4830] [INSPIRE].
- [180] CLEO collaboration, P. Naik et al., *Observation of η' decays to $\pi^+ \pi^- \pi^0$ and $\pi^+ \pi^- e^+ e^-$* , *Phys. Rev. Lett.* **102** (2009) 061801 [arXiv:0809.2587] [INSPIRE].
- [181] BESIII collaboration, M. Ablikim et al., *Measurement of $\eta' \rightarrow \pi^+ \pi^- e^+ e^-$ and $\eta' \rightarrow \pi^+ \pi^- \mu^+ \mu^-$* , *Phys. Rev. D* **87** (2013) 092011 [arXiv:1303.7360] [INSPIRE].
- [182] BABAR collaboration, B. Aubert et al., *Direct CP, Lepton Flavor and Isospin Asymmetries in the Decays $B \rightarrow K^{(*)} \ell^+ \ell^-$* , *Phys. Rev. Lett.* **102** (2009) 091803 [arXiv:0807.4119] [INSPIRE].
- [183] BELLE collaboration, J.T. Wei et al., *Measurement of the Differential Branching Fraction and Forward-Backward Asymmetry for $B \rightarrow K^{(*)} \ell^+ \ell^-$* , *Phys. Rev. Lett.* **103** (2009) 171801 [arXiv:0904.0770] [INSPIRE].
- [184] CLEO collaboration, T.J.V. Bowcock et al., *Upper Limits for the Production of Light Shortlived Neutral Particles in Radiative Υ Decay*, *Phys. Rev. Lett.* **56** (1986) 2676 [INSPIRE].
- [185] G. Mageras et al., *Search for Shortlived Axions in Radiative Υ Decays*, *Phys. Rev. Lett.* **56** (1986) 2672 [INSPIRE].
- [186] FERMILAB E760 collaboration, T.A. Armstrong et al., *Observation of the radiative decay $J/\psi \rightarrow e^+ e^- \gamma$* , *Phys. Rev. D* **54** (1996) 7067 [INSPIRE].
- [187] V.P. Druzhinin et al., *Search for Rare Radiative Decays of ϕ Meson at Vepp-2m*, *Z. Phys. C* **37** (1987) 1 [INSPIRE].
- [188] T.V. Dimova, *Radiative and other rare ρ , ω , ϕ -decays from VEPP-2M*, *Nucl. Phys. Proc. Suppl.* **181-182** (2008) 204 [INSPIRE].
- [189] M. Suzuki, *Constraints on Couplings of a Fast Decaying Axion*, *Phys. Lett. B* **175** (1986) 364 [INSPIRE].
- [190] HADES collaboration, G. Agakishiev et al., *Searching a Dark Photon with HADES*, *Phys. Lett. B* **731** (2014) 265 [arXiv:1311.0216] [INSPIRE].
- [191] M.N. Achasov et al., *Search for the $\eta' \rightarrow e^+ e^-$ decay with the SND detector*, *Phys. Rev. D* **91** (2015) 092010 [arXiv:1504.01245] [INSPIRE].
- [192] M.J. Dolan, T. Ferber, C. Hearty, F. Kahlhoefer and K. Schmidt-Hoberg, *Revised constraints and Belle II sensitivity for visible and invisible axion-like particles*, *JHEP* **12** (2017) 094 [arXiv:1709.00009] [INSPIRE].

- [193] J.A. Dror, R. Lasenby and M. Pospelov, *New constraints on light vectors coupled to anomalous currents*, *Phys. Rev. Lett.* **119** (2017) 141803 [[arXiv:1705.06726](#)] [[INSPIRE](#)].
- [194] J.A. Dror, R. Lasenby and M. Pospelov, *Dark forces coupled to nonconserved currents*, *Phys. Rev. D* **96** (2017) 075036 [[arXiv:1707.01503](#)] [[INSPIRE](#)].
- [195] M. Drewes et al., *A White Paper on keV Sterile Neutrino Dark Matter*, *JCAP* **01** (2017) 025 [[arXiv:1602.04816](#)] [[INSPIRE](#)].
- [196] A.J. Buras, D. Buttazzo, J. Girrbach-Noe and R. Knegjens, *$K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and $K_L \rightarrow \pi^0 \nu \bar{\nu}$ in the Standard Model: status and perspectives*, *JHEP* **11** (2015) 033 [[arXiv:1503.02693](#)] [[INSPIRE](#)].
- [197] NA62 collaboration, M. Corvino, *Search for $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ and Future Searches for Exotic Processes at NA62*, *Int. J. Mod. Phys. Conf. Ser.* **46** (2018) 1860068 [[INSPIRE](#)].
- [198] D.S.M. Alves and N. Weiner, to appear.
- [199] PARTICLE DATA GROUP collaboration, C. Amsler et al., *Review of Particle Physics*, *Chin. Phys. C* **38** (2014) 090001, and *Note on Scalar Mesons Below 2 GeV*, update (2015).
- [200] PARTICLE DATA GROUP collaboration, C. Patrignani et al., *Review of Particle Physics*, *Chin. Phys. C* **40** (2016) 100001 [[INSPIRE](#)].
- [201] F.E. Close and A. Kirk, *A Glueball- $q\bar{q}$ filter in central hadron production*, *Phys. Lett. B* **397** (1997) 333 [[hep-ph/9701222](#)] [[INSPIRE](#)].
- [202] T. Barnes, F.E. Close, P.R. Page and E.S. Swanson, *Higher quarkonia*, *Phys. Rev. D* **55** (1997) 4157 [[hep-ph/9609339](#)] [[INSPIRE](#)].
- [203] T. Gutsche, V.E. Lyubovitskij and M.C. Tichy, *Strong decays of radially excited mesons in a chiral approach*, *Phys. Rev. D* **79** (2009) 014036 [[arXiv:0811.0668](#)] [[INSPIRE](#)].
- [204] F.E. Close, G.R. Farrar and Z.-p. Li, *Determining the gluonic content of isoscalar mesons*, *Phys. Rev. D* **55** (1997) 5749 [[hep-ph/9610280](#)] [[INSPIRE](#)].
- [205] G.R. Farrar and G.T. Gabadadze, *Light gluino mass and condensate from properties of η and η'* , *Phys. Lett. B* **397** (1997) 104 [[hep-ph/9608330](#)] [[INSPIRE](#)].
- [206] D.M. Li, H. Yu and S.S. Fang, *The quarkonia-gluon structure of η , η' and $\eta(1410)$ revisited*, *Eur. Phys. J. C* **28** (2003) 335 [[INSPIRE](#)].
- [207] L. Faddeev, A.J. Niemi and U. Wiedner, *Glueballs, closed flux tubes and $\eta(1440)$* , *Phys. Rev. D* **70** (2004) 114033 [[hep-ph/0308240](#)] [[INSPIRE](#)].
- [208] A. Masoni, C. Cicalo and G.L. Usai, *The case of the pseudoscalar glueball*, *J. Phys. G* **32** (2006) R293 [[INSPIRE](#)].
- [209] G. Li, Q. Zhao and C.-H. Chang, *Decays of J/ψ and ψ' into vector and pseudoscalar meson and the pseudoscalar glueball- $q\bar{q}$ mixing*, *J. Phys. G* **35** (2008) 055002 [[hep-ph/0701020](#)] [[INSPIRE](#)].
- [210] H.-Y. Cheng, H.-n. Li and K.-F. Liu, *Pseudoscalar glueball mass from η - η' - G mixing*, *Phys. Rev. D* **79** (2009) 014024 [[arXiv:0811.2577](#)] [[INSPIRE](#)].
- [211] T. Gutsche, V.E. Lyubovitskij and M.C. Tichy, *$\eta(1405)$ in a chiral approach based on mixing of the pseudoscalar glueball with the first radial excitations of η and η'* , *Phys. Rev. D* **80** (2009) 014014 [[arXiv:0904.3414](#)] [[INSPIRE](#)].

- [212] B.A. Li, *Chiral field theory of 0^\mp glueball*, *Phys. Rev. D* **81** (2010) 114002 [[arXiv:0912.2323](#)] [[INSPIRE](#)].
- [213] E. Klempt and A. Zaitsev, *Glueballs, Hybrids, Multiquarks. Experimental facts versus QCD inspired concepts*, *Phys. Rept.* **454** (2007) 1 [[arXiv:0708.4016](#)] [[INSPIRE](#)].
- [214] E. Klempt, *Hadron spectroscopy and strong QCD*, *Int. J. Mod. Phys. A* **21** (2006) 739 [[INSPIRE](#)].
- [215] F. Aceti, W.H. Liang, E. Oset, J.J. Wu and B.S. Zou, *Isospin breaking and $f_0(980)$ - $a_0(980)$ mixing in the $\eta(1405) \rightarrow \pi^0 f_0(980)$ reaction*, *Phys. Rev. D* **86** (2012) 114007 [[arXiv:1209.6507](#)] [[INSPIRE](#)].
- [216] J.-J. Wu, X.-H. Liu, Q. Zhao and B.-S. Zou, *The Puzzle of anomalously large isospin violations in $\eta(1405/1475) \rightarrow 3\pi$* , *Phys. Rev. Lett.* **108** (2012) 081803 [[arXiv:1108.3772](#)] [[INSPIRE](#)].
- [217] X.-G. Wu, J.-J. Wu, Q. Zhao and B.-S. Zou, *Understanding the property of $\eta(1405/1475)$ in the J/ψ radiative decay*, *Phys. Rev. D* **87** (2013) 014023 [[arXiv:1211.2148](#)] [[INSPIRE](#)].
- [218] B.A. Dobrescu and F. Yu, *Coupling-mass mapping of dijet peak searches*, *Phys. Rev. D* **88** (2013) 035021 [[arXiv:1306.2629](#)] [[INSPIRE](#)].
- [219] CMS collaboration, *Search for low mass vector resonances decaying into quark-antiquark pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **01** (2018) 097 [[arXiv:1710.00159](#)] [[INSPIRE](#)].
- [220] CMS collaboration, *Search for narrow resonances in dijet final states at $\sqrt{s} = 8$ TeV with the novel CMS technique of data scouting*, *Phys. Rev. Lett.* **117** (2016) 031802 [[arXiv:1604.08907](#)] [[INSPIRE](#)].
- [221] ATLAS collaboration, *Search for light resonances decaying to boosted quark pairs and produced in association with a photon or a jet in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [arXiv:1801.08769](#) [[INSPIRE](#)].
- [222] ATLAS collaboration, *Search for new light resonances decaying to jet pairs and produced in association with a photon or a jet in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, [ATLAS-CONF-2016-070](#) (2016).
- [223] N. Toro and I. Yavin, *Multiphotons and photon jets from new heavy vector bosons*, *Phys. Rev. D* **86** (2012) 055005 [[arXiv:1202.6377](#)] [[INSPIRE](#)].
- [224] S.D. Ellis, T.S. Roy and J. Scholtz, *Jets and Photons*, *Phys. Rev. Lett.* **110** (2013) 122003 [[arXiv:1210.1855](#)] [[INSPIRE](#)].
- [225] S.D. Ellis, T.S. Roy and J. Scholtz, *Phenomenology of Photon-Jets*, *Phys. Rev. D* **87** (2013) 014015 [[arXiv:1210.3657](#)] [[INSPIRE](#)].
- [226] N. Arkani-Hamed and N. Weiner, *LHC Signals for a SuperUnified Theory of Dark Matter*, *JHEP* **12** (2008) 104 [[arXiv:0810.0714](#)] [[INSPIRE](#)].
- [227] C. Cheung, J.T. Ruderman, L.-T. Wang and I. Yavin, *Lepton Jets in (Supersymmetric) Electroweak Processes*, *JHEP* **04** (2010) 116 [[arXiv:0909.0290](#)] [[INSPIRE](#)].
- [228] A. Katz and R. Sundrum, *Breaking the Dark Force*, *JHEP* **06** (2009) 003 [[arXiv:0902.3271](#)] [[INSPIRE](#)].
- [229] Y. Bai and Z. Han, *Measuring the Dark Force at the LHC*, *Phys. Rev. Lett.* **103** (2009) 051801 [[arXiv:0902.0006](#)] [[INSPIRE](#)].

- [230] M. Baumgart, C. Cheung, J.T. Ruderman, L.-T. Wang and I. Yavin, *Non-Abelian Dark Sectors and Their Collider Signatures*, *JHEP* **04** (2009) 014 [[arXiv:0901.0283](#)] [[INSPIRE](#)].
- [231] Y.F. Chan, M. Low, D.E. Morrissey and A.P. Spray, *LHC Signatures of a Minimal Supersymmetric Hidden Valley*, *JHEP* **05** (2012) 155 [[arXiv:1112.2705](#)] [[INSPIRE](#)].
- [232] T. Han, Z. Si, K.M. Zurek and M.J. Strassler, *Phenomenology of hidden valleys at hadron colliders*, *JHEP* **07** (2008) 008 [[arXiv:0712.2041](#)] [[INSPIRE](#)].
- [233] A. Falkowski, J.T. Ruderman, T. Volansky and J. Zupan, *Discovering Higgs Decays to Lepton Jets at Hadron Colliders*, *Phys. Rev. Lett.* **105** (2010) 241801 [[arXiv:1007.3496](#)] [[INSPIRE](#)].
- [234] D. Curtin et al., *Exotic decays of the 125 GeV Higgs boson*, *Phys. Rev. D* **90** (2014) 075004 [[arXiv:1312.4992](#)] [[INSPIRE](#)].
- [235] A. Gupta, R. Primulando and P. Saraswat, *A New Probe of Dark Sector Dynamics at the LHC*, *JHEP* **09** (2015) 079 [[arXiv:1504.01385](#)] [[INSPIRE](#)].
- [236] B. Dasgupta, J. Kopp and P. Schwaller, *Photons, photon jets and dark photons at 750 GeV and beyond*, *Eur. Phys. J. C* **76** (2016) 277 [[arXiv:1602.04692](#)] [[INSPIRE](#)].
- [237] Y. Tsai, L.-T. Wang and Y. Zhao, *Faking ordinary photons by displaced dark photon decays*, *Phys. Rev. D* **95** (2017) 015027 [[arXiv:1603.00024](#)] [[INSPIRE](#)].
- [238] M. Bauer, M. Neubert and A. Thamm, *LHC as an Axion Factory: Probing an Axion Explanation for $(g - 2)_\mu$ with Exotic Higgs Decays*, *Phys. Rev. Lett.* **119** (2017) 031802 [[arXiv:1704.08207](#)] [[INSPIRE](#)].
- [239] M. Bauer, M. Neubert and A. Thamm, *Collider Probes of Axion-Like Particles*, *JHEP* **12** (2017) 044 [[arXiv:1708.00443](#)] [[INSPIRE](#)].