

Hadronic vacuum polarization using gradient flow

Robert V. Harlander,^a Fabian Lange^a and Tobias Neumann^{b,c}

^a*TTK, RWTH Aachen University,
Sommerfeldstr. 16, 52056 Aachen, Germany*

^b*Department of Physics, Illinois Institute of Technology,
3101 S. Dearborn St., Chicago, IL 60616, U.S.A.*

^c*Fermilab,
P.O. Box 500, Batavia, IL 60510, U.S.A.*

E-mail: harlander@physik.rwth-aachen.de,
flange@physik.rwth-aachen.de, tneumann@fnal.gov

ABSTRACT: The gradient-flow operator product expansion for QCD current correlators including operators up to mass dimension four is calculated through NNLO. This paves an alternative way for efficient lattice evaluations of hadronic vacuum polarization functions. In addition, flow-time evolution equations for flowed composite operators are derived. Their explicit form for the non-trivial dimension-four operators of QCD is given through order α_s^3 .

KEYWORDS: Lattice QCD, Perturbative QCD, Renormalization Group

ARXIV EPRINT: [2007.01057](https://arxiv.org/abs/2007.01057)

Contents

1	Introduction	1
2	Current-current correlators	3
2.1	Operator product expansion	3
2.2	Coefficient functions	5
3	Flowed operator product expansion	6
3.1	Flowed operators	6
3.2	Small-flow-time expansion	8
4	Calculation of the mixing matrix	9
5	Flow-time evolution	13
6	Conclusions	16
A	Renormalization group functions	17
B	Perturbative coefficient functions	19
B.1	Vector and axial-vector currents	19
B.2	Scalar and pseudo-scalar currents	21
C	Renormalized mixing matrix	22
D	Supplementary material	23

1 Introduction

The vacuum polarization functions (VPFs) for (axial-)vector and (pseudo-)scalar particles are among the most important objects when studying QCD. On the one hand, this is because their imaginary part is directly related to physical observables such as the decay rates of the Z - or the Higgs boson, or the hadronic R-ratio. If the characteristic energy scale of these quantities is far above the QCD scale, a perturbative evaluation of the polarization functions is sufficient in these cases to arrive at high-precision results (see, e.g., ref. [1]).

But VPFs also contribute indirectly to physical observables such as anomalous magnetic moments [2, 3], the definition of short-distance quark masses [4], or hadronic contributions to the QED coupling [5, 6]. These applications involve an integration of the VPFs over the non-perturbative regime, which is typically achieved with the help of experimental data and dispersion relations. Only very recently, first-principle lattice calculations have

become competitive with these dispersive approaches. In the case of the hadronic vacuum polarization contribution to the muon’s anomalous magnetic moment, the two approaches turn out to lead to incompatible results [7].¹ It would therefore be highly desirable to have additional independent first-principle calculations of the VPF.

About ten years ago, the gradient-flow formalism (GFF) was suggested as a mechanism to improve the efficiency of lattice calculations [10–12] (see also refs. [13, 14]). Since then, it has become a standard for the scale-setting procedure [15, 16]. However, also other applications of the GFF have been studied, among them a new way to determine the energy-momentum tensor on the lattice. The underlying idea in this case is the small-flow-time expansion of composite operators [11], leading to the flowed Operator Product Expansion (OPE) (also named smeared OPE in refs. [17, 18]), where the regular operators are replaced by operators taken at finite flow time. Its main advantages with respect to (w.r.t.) the regular OPE is the absence of operator mixing, and the improved efficiency of the evaluation of operator matrix elements. The translation of the regular to the flowed operators can be done perturbatively. For the energy-momentum tensor, it is available through next-to-next-to-leading order (NNLO) [19–21]. Quite recently, the small-flow-time expansion was applied at next-to-leading order (NLO) to CP-violating operators [22], and to four-quark operators [23].

In this paper, we present the flowed OPE for the time-ordered product of two currents through NNLO QCD. Taking the vacuum expectation value (VEV) leads to the VPF. This should thus allow for an alternative first-principle evaluation of VPFs on the lattice. In addition, we derive a general logarithmic flow-time evolution equation for flowed operators which resembles the renormalization group (RG) equation of regular operators.

The remainder of this paper is organized as follows. Section 2 introduces the regular OPE of current correlators with operators up to mass dimension four. This includes the renormalization of these operators as well as an overview of the literature which provides the corresponding perturbative Wilson coefficients. (The coefficients for the dimension-four operators for various currents are reproduced in appendix B.) The transition to the flowed OPE is presented in section 3. Section 4 describes the calculation of the mixing matrix between regular and flowed operators in the small-flow-time limit. While a large part of this mixing matrix is already known [21] and recollected in appendix C, the missing components require higher order mass terms of the VEV of the flowed dimension-four operators and their renormalization with the help of the vacuum-energy renormalization constant. These results complete the ingredients required for the flowed OPE of the VPF through NNLO. In section 5, we derive a logarithmic evolution equation from the generic flow-time dependence of the mixing matrix. Section 6 presents our conclusions and gives a short outlook on possible extensions of this work.

¹The lattice calculation of the light-by-light contribution to $(g - 2)_\mu$ is in agreement with other determinations though [8, 9].

2 Current-current correlators

Our results are presented for a general non-Abelian gauge theory based on a simple compact Lie group with n_f quark fields $\psi_1, \dots, \psi_{n_f}$ in the fundamental representation, of which the first n_h are degenerate with mass m , while the remaining n_l are massless. The generators T^a of the fundamental representation are normalized as $\text{Tr}(T^a T^b) = -T_R \delta^{ab}$, and the structure constants f^{abc} are defined through the Lie algebra $[T^a, T^b] = f^{abc} T^c$. The dimensions of the fundamental and the adjoint representation are n_c and n_A , respectively, and their quadratic Casimir eigenvalues are denoted by C_F and C_A . For $\text{SU}(N)$, it is

$$n_c = N, \quad n_A = N^2 - 1, \quad C_F = T_R \frac{N^2 - 1}{N}, \quad C_A = 2T_R N, \quad (2.1)$$

and QCD is recovered for $T_R = 1/2$ and $N = 3$, i.e. $C_F = 4/3$ and $C_A = 3$. For brevity, we often use ‘‘QCD’’ also to refer to the more general gauge theory in the following.

2.1 Operator product expansion

The role of the perturbative and the non-perturbative regime of VPFs can be made most explicit through the OPE (see, e.g., ref. [24]):

$$T(Q) \equiv \int d^4x e^{iQx} \langle T j(x) j(0) \rangle \stackrel{Q^2 \rightarrow \infty}{\sim} \sum_{k,n} C_n^{(k),B}(Q) \langle O_n^{(k)}(x=0) \rangle, \quad (2.2)$$

where $j(x)$ generically stands for a scalar, pseudo-scalar, vector, axial-vector, or tensor current, and k labels the mass dimension. Figure 1 shows sample Feynman diagrams which arise from the perturbative evaluation of the current correlator in eq. (2.2). In the following, we only consider the so-called non-singlet diagrams, where the currents are connected by a common quark line. An example for a singlet-diagram, on the other hand, is shown in figure 1(e).

The coefficients $C_n^{(k),B}$ on the right-hand side of eq. (2.2) depend on the quantum numbers of the current and may thus carry Lorentz indices. Apart from the explicit results for specific currents in appendix B, we suppress these indices throughout the paper. We furthermore assume that, upon transition from the left- to the right-hand side, possible global divergences are subtracted off of $T(Q)$.

Up to mass dimension two, the only operators of QCD which contribute to physical matrix elements are proportional to unity, i.e.,

$$O_1^{(0)} \equiv O^{(0)} = \mathbb{1}, \quad O_1^{(2)} \equiv O^{(2)} = m_B^2 \mathbb{1}, \quad (2.3)$$

where m_B is the bare mass of the n_h degenerate massive quarks. This means that

$$C_1^{(0)} \equiv C^{(0)} \equiv C^{(0),B}, \quad C_1^{(2)} \equiv C^{(2)} \equiv Z_m^2 C^{(2),B} \quad (2.4)$$

are ultra-violet (UV)-finite, where Z_m is the renormalization constant of the quark mass defined in appendix A.




Figure 1. Sample diagrams contributing to the perturbative calculation of the VPF, i.e., the left-hand side of eq. (2.2). The currents are symbolized by wavy lines, gluons by spirals. We consider the case where n_l quarks are massless (thin straight lines), and n_h quarks are degenerate with mass m (thick straight lines). (a) One-loop contribution for non-diagonal currents; (b-e) sample three-loop diagrams for diagonal currents. In (d), the currents couple to massless quarks. (e) is a “singlet” diagram. All diagrams in this paper were produced with the help of FeynGame [25].

At mass dimension four, we choose the following basis of operators (the space-time argument is suppressed in most of what follows):

$$\begin{aligned}
 O_1^{(4)} &\equiv O_1 = \frac{1}{g_B^2} F_{\mu\nu}^a F_{\mu\nu}^a, \\
 O_2^{(4)} &\equiv O_2 = \sum_{q=1}^{n_l} \bar{\psi}_q \overleftrightarrow{D} \psi_q, \\
 O_3^{(4)} &\equiv O_3 = m_B^4,
 \end{aligned}
 \tag{2.5}$$

where

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + f^{abc} A_\mu^b A_\nu^c, \quad \overleftrightarrow{D}_\mu = \partial_\mu - \overleftarrow{\partial}_\mu + 2A_\mu^a T^a,
 \tag{2.6}$$

with the regular (as opposed to “flowed”, see section 3) quark and gluon fields $\psi_q(x)$ and $A_\mu^a(x)$, respectively, and the bare coupling g_B .

We employ Euclidean space-time, but translation of the intermediate formulas and the final results to Minkowski space is possible without difficulty. Working in

$$d = 4 - 2\epsilon
 \tag{2.7}$$

space-time dimensions, the mass dimensions of O_1 and O_2 are actually equal to d , while that of O_3 is equal to 4. Higher dimensional operators are neglected in the following.

The set in eq. (2.5) does not contain gauge dependent operators, or operators that vanish due to the equations of motion when acting on physical states, since they are irrelevant for the scope of this paper. In fact, in this respect the upper limit of the sum over q in O_2 could be replaced by n_h , because the terms with massless quarks vanish on-shell. For the same reason, one could use $O'_2 \equiv -2m_B \sum_{q=1}^{n_h} \bar{\psi}_q \psi_q$ instead of O_2 in the

definition of the operator basis (2.5). Other choices are possible as well, but the basis in eq. (2.5) is particularly suitable for our purposes, because it is most directly related to the operator basis used in refs. [20, 21, 26].

Matrix elements of the dimension-four operators are divergent in general. However, one may define “renormalized operators” O_n^R as linear combinations among them, for which physical matrix elements become finite:

$$O_n^R = \sum_k Z_{nk} O_k. \tag{2.8}$$

Analogously, one defines renormalized coefficient functions through the condition

$$\sum_n C_n^B O_n \stackrel{!}{=} \sum_n C_n O_n^R \Rightarrow C_n = \sum_m C_m^B (Z^{-1})_{mn}, \tag{2.9}$$

where $C_n^B \equiv C_n^{(4),B}$, cf. eqs. (2.2) and (2.5). It is well known that, since the operators of eq. (2.5) are part of the QCD Lagrangian, the renormalization matrix Z can be expressed in terms of the anomalous dimensions of QCD [27, 28]:

$$Z = \begin{pmatrix} Z_{2 \times 2} & \vec{Z}_3 \\ \vec{0}^T & Z_m^{-4} \end{pmatrix}, \quad \text{where } Z_{2 \times 2} = \begin{pmatrix} -\epsilon/\beta_\epsilon & -2\gamma_m/\beta_\epsilon \\ 0 & 1 \end{pmatrix}, \tag{2.10}$$

$$\vec{Z}_3 = 4\hat{\mu}^{-2\epsilon} Z_m^{-4} \begin{pmatrix} a_s \frac{\partial}{\partial a_s} Z_0 \\ 2 Z_0 \end{pmatrix}, \quad \hat{\mu} \equiv \frac{\mu}{\sqrt{4\pi}} e^{\gamma_E/2}.$$

The ’t Hooft mass μ ensures that each renormalized operator O_n^R in eq. (2.8) has the same mass dimension as the corresponding bare one, and $\hat{\mu}$ appears because we will adopt the $\overline{\text{MS}}$ scheme by default ($\gamma_E = -\Gamma'(1) = 0.577216\dots$). We also introduced the quantity $a_s = \alpha_s/\pi = g^2/(4\pi^2)$ here, where g is the renormalized strong coupling in the $\overline{\text{MS}}$ scheme. Z_0 is the $\overline{\text{MS}}$ renormalization constant for the vacuum energy [28]. It is given in appendix A, together with the anomalous quark mass dimension γ_m and the d -dimensional beta function β_ϵ .

2.2 Coefficient functions

The OPE form of the current correlators is usually inconvenient for their perturbative evaluation. Instead, one rather evaluates the l.h.s. of eq. (2.2) directly by calculating the relevant two-point functions to the appropriate order. The exact analytical result for general quark masses is known at the two-loop level [29–32], while higher orders up to the four-loop level have been reconstructed by combining various kinematical limits [33–43], or through integral reduction [44–46] and subsequent numerical evaluation of the resulting master integrals [47].

Since the dimension-zero and -two operators in eq. (2.2) are proportional to unity, the coefficients $C^{(0)}$ and $C^{(2)}$ are immediately determined from the small-mass expansion of

these perturbative results for the VPF. They are thus known up to the four-loop level at the moment [48–56].²

The Wilson coefficients C_n of the dimension-four operators, on the other hand, require a dedicated calculation which keeps track of the contributions from the individual operators. This has been done through $\mathcal{O}(a_s^3)$ for C_1 and C_2 , and through $\mathcal{O}(a_s^2)$ for C_3 in refs. [61–64]. For the purpose of this paper, only the $\mathcal{O}(a_s^2)$ results are required. For completeness, we include them in appendix B.

3 Flowed operator product expansion

Having introduced the setup in the “regular” theory, we now translate this to the flowed OPE for the current correlators.

3.1 Flowed operators

We introduce the flowed operators as

$$\begin{aligned}\tilde{O}_1(t, x) &= \frac{Z_s}{g_B^2} G_{\mu\nu}^a(t, x) G_{\mu\nu}^a(t, x) = \frac{\hat{\mu}^{-2\epsilon}}{g^2} G_{\mu\nu}^a(t, x) G_{\mu\nu}^a(t, x), \\ \tilde{O}_2(t, x) &= \dot{Z}_\chi \sum_{q=1}^{n_f} \bar{\chi}_q(t, x) \overleftrightarrow{\mathcal{D}}(t, x) \chi_q(t, x), \\ \tilde{O}_3(t, x) &= m^4,\end{aligned}\tag{3.1}$$

where

$$\overleftrightarrow{\mathcal{D}}_\mu = \mathcal{D}_\mu - \overleftarrow{\mathcal{D}}_\mu, \quad \mathcal{D}_\mu = \partial_\mu + B_\mu^a T^a, \quad \overleftarrow{\mathcal{D}}_\mu = \overleftarrow{\partial}_\mu - B_\mu^a T^a.\tag{3.2}$$

The flowed gauge and quark fields $B_\mu^a = B_\mu^a(x, t)$ and $\chi_q = \chi_q(x, t)$ obey the equations [10, 12]

$$\begin{aligned}\partial_t B_\mu^a &= \mathcal{D}_\nu^{ab} G_{\nu\mu}^b + \kappa \mathcal{D}_\mu^{ab} \partial_\nu B_\nu^b, \\ \partial_t \chi_q &= \Delta \chi_q - \kappa \partial_\mu B_\mu^a T^a \chi_q, \\ \partial_t \bar{\chi}_q &= \bar{\chi}_q \overleftarrow{\Delta} + \kappa \bar{\chi}_q \partial_\mu B_\mu^a T^a,\end{aligned}\tag{3.3}$$

with the initial conditions

$$B_\mu^a(t=0, x) = A_\mu^a(x), \quad \chi_q(t=0, x) = \psi_q(x), \quad q \in \{1, \dots, n_f\}.\tag{3.4}$$

Here we used the flowed covariant derivative in the adjoint representation,

$$\mathcal{D}_\mu^{ab} = \delta^{ab} \partial_\mu - f^{abc} B_\mu^c,\tag{3.5}$$

and the flowed d’Alembert operators

$$\Delta = \mathcal{D}_\mu \mathcal{D}_\mu, \quad \overleftarrow{\Delta} = \overleftarrow{\mathcal{D}}_\mu \overleftarrow{\mathcal{D}}_\mu,\tag{3.6}$$

²The imaginary parts of the VPFs are known even at the five-loop level in the phenomenologically most relevant cases [57–60].

where the flowed covariant derivatives in the fundamental representation are given in eq. (3.2). For our three-loop calculations, we set the gauge parameter $\kappa = 1$ in eq. (3.3), because this simplifies the intermediate expressions. All our results are independent of this choice though.

\mathring{Z}_χ is the non-minimal renormalization constant for the flowed quark fields χ_q , defined by the all-order condition [20]

$$\langle \tilde{\mathcal{O}}_2(t) \rangle \Big|_{m=0} \equiv -\frac{2n_c n_f}{(4\pi t)^2}, \quad (3.7)$$

where $\langle \cdot \rangle$ denotes the VEV. It reads

$$\mathring{Z}_\chi = \zeta_\chi Z_\chi, \quad (3.8)$$

where Z_χ is the $\overline{\text{MS}}$ part,

$$Z_\chi = 1 + a_s \frac{\gamma_{\chi,0}}{2\epsilon} + a_s^2 \left[\frac{\gamma_{\chi,0}}{4\epsilon^2} \left(\frac{\gamma_{\chi,0}}{2} - \beta_0 \right) + \frac{\gamma_{\chi,1}}{4\epsilon} \right] + \mathcal{O}(a_s^3), \quad (3.9)$$

and

$$\begin{aligned} \zeta_\chi = 1 + a_s \left(\frac{\gamma_{\chi,0}}{2} L_{\mu t} - \frac{3}{4} C_F \ln 3 - C_F \ln 2 \right) \\ + a_s^2 \left\{ \frac{\gamma_{\chi,0}}{4} \left(\beta_0 + \frac{\gamma_{\chi,0}}{2} \right) L_{\mu t}^2 + \left[\frac{\gamma_{\chi,1}}{2} - \frac{\gamma_{\chi,0}}{2} \left(\beta_0 + \frac{\gamma_{\chi,0}}{2} \right) \ln 3 \right. \right. \\ \left. \left. - \frac{2}{3} \gamma_{\chi,0} \left(\beta_0 + \frac{\gamma_{\chi,0}}{2} \right) \ln 2 \right] L_{\mu t} + \frac{c_\chi^{(2)}}{16} \right\} + \mathcal{O}(a_s^3). \end{aligned} \quad (3.10)$$

The short-hand notation

$$L_{\mu t} = \ln \frac{\mu^2}{\mu_t^2}, \quad \mu_t = \frac{1}{\sqrt{2t\epsilon^{\gamma_E}}} \quad (3.11)$$

reflects our choice of the ‘‘central’’ renormalization scale μ_t [21].

The minimal renormalization constant Z_χ is known analytically through NNLO [12, 21],

$$\begin{aligned} \gamma_{\chi,0} &= \frac{3}{2} C_F, \\ \gamma_{\chi,1} &= C_F \left[C_A \left(\frac{223}{48} - \ln 2 \right) - C_F \left(\frac{3}{16} + \ln 2 \right) - \frac{11}{12} T_R n_f \right]. \end{aligned} \quad (3.12)$$

The finite coefficient in eq. (3.10) has been obtained numerically in ref. [65]:

$$c_\chi^{(2)} = C_A C_F c_{\chi,A} + C_F^2 c_{\chi,F} + C_F T_R n_f c_{\chi,R}, \quad (3.13)$$

with³

$$\begin{aligned} c_{\chi,A} &= -23.7947, \quad c_{\chi,F} = 30.3914, \\ c_{\chi,R} &= -\frac{131}{18} + \frac{46}{3} \zeta(2) + \frac{944}{9} \ln 2 + \frac{160}{3} \ln^2 2 - \frac{172}{3} \ln 3 + \frac{104}{3} \ln 2 \ln 3 \\ &\quad - \frac{178}{3} \ln^2 3 + \frac{8}{3} \text{Li}_2(1/9) - \frac{400}{3} \text{Li}_2(1/3) + \frac{112}{3} \text{Li}_2(3/4) = -3.92255\dots, \end{aligned} \quad (3.14)$$

³The sign on the r.h.s. of equation (B.3) in ref. [65] is incorrect.

with Riemann's zeta function $\zeta(s) \equiv \sum_{n=1}^{\infty} n^{-s}$ and the di-logarithm $\text{Li}_2(z) = \sum_{k=1}^{\infty} z^k/k^2$. The strong coupling renormalization constant Z_s in eq. (3.1) ensures that matrix elements of $\tilde{O}_1(t)$ are finite [10, 11]. The reason for keeping track of the non-integer mass dimension of $\tilde{O}_1(t)$ is clarified later.

Eq. (3.14) displays only the first six leading digits in numerical results. Results with higher accuracy are provided in the supplementary material, which also includes the $L_{\mu t}$ terms, see appendix D. We expect that these floating-point numbers can be considered equivalent to their exact results for all practical purposes. This is why we often use the numerical values for the coefficients in what follows, even if the exact result is available.

Similar to the regular operators in eq. (2.5), one could trade the flowed operator $\tilde{O}_2(t)$ for $\tilde{O}'_2(t) = -2m\tilde{Z}_\chi \sum_{q=1}^{n_f} \tilde{\chi}_q(t)\chi_q(t)$. However, in this case the final results to be derived below would be different, because the equations of motion for the flowed operators relate $\tilde{O}'_2(t)$ to both $\tilde{O}_2(t)$ and $\tilde{O}_1(t)$ (see refs. [20, 21]). A transformation of the results in this paper to $\tilde{O}'_2(t)$ is straightforward though.

3.2 Small-flow-time expansion

The small-flow-time expansion allows us to relate the QCD operators and coefficients with the flowed quantities as follows:

$$\begin{aligned} \tilde{O}_n(t) &= \zeta_n^{(0)}(t)\mathbf{1} + \zeta_n^{(2),\text{B}}(t) m_{\text{B}}^2\mathbf{1} + \sum_k \zeta_{nk}^{\text{B}}(t) O_k + \dots \\ &\equiv \zeta_n^{(0)}(t)\mathbf{1} + \zeta_n^{(2)}(t) m^2\mathbf{1} + \sum_k \zeta_{nk}(t) O_k^{\text{R}} + \dots, \end{aligned} \tag{3.15}$$

where the ellipsis denotes terms that vanish as $t \rightarrow 0$, and

$$\zeta_n^{(2)}(t) = \zeta_n^{(2),\text{B}}(t) Z_m^2, \quad \zeta_{nk}(t) = \sum_l \zeta_{nl}^{\text{B}}(t) Z_{lk}^{-1} \tag{3.16}$$

are the renormalized, finite mixing coefficients. Inversion of eq. (3.15) gives

$$\begin{aligned} O_n^{\text{R}} &= \sum_k \zeta_{nk}^{-1}(t) \tilde{O}_k(t) + \dots, \\ \tilde{O}_n(t) &\equiv \tilde{O}_n(t) - \zeta_n^{(0)}(t)\mathbf{1} - \zeta_n^{(2)}(t)m^2\mathbf{1}, \end{aligned} \tag{3.17}$$

where ζ_{nk}^{-1} is the nk -element of the inverse of the mixing matrix ζ . This lets one define the “flowed OPE” for the current correlator:

$$T(Q) \stackrel{Q^2 \rightarrow \infty}{\sim} \tilde{C}^{(0)}(Q^2, t)\mathbf{1} + \tilde{C}^{(2)}(Q^2, t)m^2\mathbf{1} + \sum_n \tilde{C}_n(Q^2, t)\tilde{O}_n(t) + \dots \tag{3.18}$$

where the corresponding coefficient functions are related to the regular Wilson coefficients through

$$\begin{aligned} \tilde{C}_n(Q^2, t) &= \sum_k C_k(Q^2)\zeta_{kn}^{-1}(t), \\ \tilde{C}^{(0,2)}(Q^2, t) &= C^{(0,2)}(Q^2) - \sum_n \tilde{C}_n(t, Q^2)\zeta_n^{(0,2)}(t). \end{aligned} \tag{3.19}$$

The regular QCD coefficients $C^{(0)}$ and $C^{(2)}$ are given by the first two terms in m^2/Q^2 of the large- Q^2 expansion of the VPFs. Through the required order, they can be found in ref. [53] for diagonal vector- and axial-vector currents, and in ref. [54] for scalar- and pseudo-scalar currents, for example. The dimension-four coefficients can be found in refs. [61, 64]. For convenience of the reader, they are also collected in appendix B.

4 Calculation of the mixing matrix

We now determine the mixing matrix ζ in a perturbative calculation through NNLO. By using the known results for the regular Wilson coefficients given in appendix B, one can determine the flowed coefficients of eq. (3.19) to the same order. Together with an evaluation of the flowed operator matrix elements on the lattice, the VPFs can be extracted and used in the determination of various physical quantities.

The bare mixing matrix ζ^{B} can be determined with the help of the *method of projectors*:

$$\zeta_n^{(0,2),\text{B}}(t) = P^{(0,2)}[\tilde{O}_n(t)], \quad \zeta_{nk}^{\text{B}}(t) = P_k^{(4)}[\tilde{O}_n(t)], \quad (4.1)$$

where the action of $P^{(n)}$ is to take suitable derivatives of a specific Green's function of the operator such that

$$P^{(n)}[O^{(m)}] = \delta_{nm}, \quad P^{(n)}[O_k] = P_k^{(4)}[O^{(n)}] = 0, \quad P_k^{(4)}[O_l] = \delta_{kl}, \quad (4.2)$$

for $n, m \in \{0, 2\}$ and $k, l \in \{1, 2, 3\}$. For details, see refs. [21, 66, 67].

Specifically, the projectors onto $\mathbb{1}$, $m_{\text{B}}^2 \mathbb{1}$, and O_3^{B} are given by derivatives of vacuum matrix elements w.r.t. m_{B} :

$$\begin{aligned} \zeta_n^{(0)}(t) &= P^{(0)}[\tilde{O}_n(t)] \equiv \langle \tilde{O}_n(t) \rangle \Big|_{m_{\text{B}}=0}, \\ \zeta_n^{(2)}(t) &= Z_m^2 P^{(2)}[\tilde{O}_n(t)] \equiv Z_m^2 \frac{1}{2!} \frac{\partial^2}{\partial m_{\text{B}}^2} \langle \tilde{O}_n(t) \rangle \Big|_{m_{\text{B}}=0}, \\ \zeta_{n3}^{\text{B}}(t) &= P_3^{(4)}[\tilde{O}_n(t)] \equiv \frac{1}{4!} \frac{\partial^4}{\partial m_{\text{B}}^4} \langle \tilde{O}_n(t) \rangle \Big|_{m_{\text{B}}=0}. \end{aligned} \quad (4.3)$$

A crucial point of the *method of projectors* is that the derivatives and limits are taken *before* loop integration [66, 67]. This guarantees that all loop contributions of projections on the r.h.s. of eq. (3.15) lead to scaleless integrals and thus vanish in dimensional regularization. On the other hand, it implies that the projections on the l.h.s. of eq. (3.15) can be divergent, even though physical matrix elements of $\tilde{O}_n(t)$ are finite. This is why we need to carefully account for a possible non-integer mass dimension of these operators, see eq. (3.1).

We directly obtain

$$\zeta_{33}^{\text{B}} = \frac{1}{4!} \frac{\partial^4}{\partial m_{\text{B}}^4} m^4 = Z_m^{-4}, \quad \zeta_3^{(0),\text{B}} = \zeta_3^{(2),\text{B}} = 0, \quad \zeta_{31}^{\text{B}} = \zeta_{32}^{\text{B}} = 0, \quad (4.4)$$

where the third set of equations follows from $\tilde{O}_3(t) = m^4 = O_3$ and the projector property $P_1^{(4)}[O_3] = P_2^{(4)}[O_3] = 0$, see eq. (4.2).

The bare and renormalized mixing matrices for the dimension-four operators thus take the form

$$\zeta^{\text{B}} = \begin{pmatrix} \zeta_{2 \times 2}^{\text{B}} & \vec{\zeta}_3^{\text{B}} \\ \vec{0}^T & Z_m^{-4} \end{pmatrix}, \quad \zeta = \begin{pmatrix} \zeta_{2 \times 2} & \vec{\zeta}_3 \\ \vec{0}^T & 1 \end{pmatrix}, \quad (4.5)$$

where $\vec{0}^T = (0, 0)$, and

$$\begin{aligned} \zeta_{2 \times 2}^{\text{B}} &= \begin{pmatrix} \zeta_{11}^{\text{B}} & \zeta_{12}^{\text{B}} \\ \zeta_{21}^{\text{B}} & \zeta_{22}^{\text{B}} \end{pmatrix}, & \zeta_{2 \times 2} &= \zeta_{2 \times 2}^{\text{B}} Z_{2 \times 2}^{-1}, \\ \vec{\zeta}_3^{\text{B}} &= (\zeta_{13}^{\text{B}}, \zeta_{23}^{\text{B}})^T, & \vec{\zeta}_3 &= (\vec{\zeta}_3^{\text{B}} - \zeta_{2 \times 2}^{\text{B}} \vec{Z}_3) Z_m^4. \end{aligned} \quad (4.6)$$

$\zeta_{2 \times 2}$ can be obtained from the mixing matrix of the operators occurring in the energy-momentum tensor and is accordingly known through NNLO [21]. Explicit results are given in appendix C.

The coefficient $\zeta_n^{(0)}$ is simply the VEV of $\tilde{O}_n(t)$ for $m = 0$. For $\tilde{O}_1(t)$ it has been calculated through NNLO in refs. [10, 65, 68]:⁴

$$\begin{aligned} \zeta_1^{(0)}(t) = \langle \tilde{O}_1(t) \rangle \Big|_{m=0} &= \frac{3}{4\pi^2 t^2} \frac{n_A}{8} \left\{ 1 + \bar{a}_s \left[C_A \left(\frac{13}{9} + \frac{11}{6} \ln 2 - \frac{3}{4} \ln 3 \right) - \frac{2}{9} T_{\text{R}} n_f \right] \right. \\ &\quad \left. + \bar{a}_s^2 \left[1.74865 C_A^2 - 1.97283 C_A T_{\text{R}} n_f \right. \right. \\ &\quad \left. \left. + 0.306224 C_{\text{F}} T_{\text{R}} n_f + 0.121042 T_{\text{R}}^2 n_f^2 \right] \right\} + \mathcal{O}(\bar{a}_s^3), \end{aligned} \quad (4.7)$$

where $\bar{a}_s = a_s(\mu_t)$. Due to the RG invariance of $a_s \tilde{O}_1(t)$ [10, 11], the result for general values of the 't Hooft mass μ can be obtained by multiplying the result given in eq. (4.7) by \bar{a}_s/a_s , replacing

$$\bar{a}_s = a_s \left[1 + a_s \beta_0 L_{\mu t} + a_s^2 L_{\mu t} (\beta_1 + \beta_0^2 L_{\mu t}) \right] + \mathcal{O}(a_s^3), \quad (4.8)$$

and re-expanding in a_s .

For \tilde{O}_2 , on the other hand, we have eq. (3.7) to all orders in perturbation theory by definition, i.e.

$$\zeta_2^{(0)}(t) = \langle \tilde{O}_2(t) \rangle \Big|_{m=0} \equiv -\frac{2n_c n_f}{(4\pi t)^2}. \quad (4.9)$$

Therefore, the only coefficients that are not yet known through NNLO are

$$\zeta_1^{(2)}, \zeta_2^{(2)}, \zeta_{13} \text{ and } \zeta_{23}. \quad (4.10)$$

According to eq. (4.3), they require the calculation of derivatives w.r.t. m_{B} in $\langle \tilde{O}_1(t) \rangle$ and $\langle \tilde{O}_2(t) \rangle$ up to three loops (see figure 2 for a set of associated diagrams). We achieved this using the setup described in ref. [65], which employs `qgraf` [69, 70] and `q2e/exp` [71, 72] for the generation and subsequent categorization of the Feynman diagrams, FORM [73, 74]

⁴The coefficient of the C_A^2 term in ref. [65] contains a typo: instead of 27.9786, it should read 27.9784.




Figure 2. Sample diagrams contributing to $\zeta_n^{(m)}$ ($m \in \{0, 2\}$) and ζ_{n3} , for $n = 1$ (a-d) and $n = 2$ (e-h). The notation is the same as in figure 1; in addition, white circles denote flow-vertices, and lines with arrows next to them denote flow-lines (see ref. [65] for details). Diagrams (a) and (b) only contribute to $\zeta_1^{(0)}$.

for the manipulation of the ensuing algebraic expressions, the `color` package [75] of `FORM` for the calculation of the gauge group factors, and `Kira`⊕`FireFly` [76–80] for the Feynman integral reduction using integration-by-parts identities and the Laporta algorithm [44–46] over finite fields [81–84]. For the evaluation of the master integrals, we adopt the method described in ref. [68], which performs sector decomposition [85] with the help of `FIESTA` [86, 87] in order to extract the UV poles, along with a fully symmetric integration rule of order 13 for the numerical evaluation of their coefficients [88], implemented with high precision arithmetics by using the `MPFR` library [89]. Some intermediate steps of the calculation are done within `Mathematica` [90].

Multiplying the result by Z_m^2 suffices to obtain the renormalized expressions for $\zeta_n^{(2)}$, and we find, setting $\mu = \mu_t$,

$$\zeta_1^{(2)}(t) = \frac{3n_A n_h}{8\pi^2 t} T_R \bar{a}_s \left[1 + \bar{a}_s \left(7.43789 C_A + 2 C_F - \frac{10}{9} T_R n_f \right) \right] + \mathcal{O}(a_s^3), \quad (4.11a)$$

$$\zeta_2^{(2)}(t) = \frac{n_c n_h}{4\pi^2 t} \left[1 + \bar{a}_s C_F \left(\frac{7}{4} - \frac{3}{4} \ln 3 \right) + \bar{a}_s^2 C_F (2.10889 C_A + 2.05158 C_F - 0.268909 T_R n_f) \right] + \mathcal{O}(a_s^3), \quad (4.11b)$$

where again $\bar{a}_s = a_s(\mu_t)$. Since quark loops appear in $\langle \tilde{O}_1(t) \rangle$ only at the two-loop level, $\zeta_1^{(2)}(t)$ starts at $\mathcal{O}(a_s)$. In the case of $\zeta_2^{(2)}(t)$, the result for general μ can be obtained

through multiplication by

$$\frac{m^2(\mu_t)}{m^2(\mu)} = 1 + a_s \gamma_{m,0} L_{\mu t} + a_s^2 L_{\mu t} \left[\gamma_{m,1} + \frac{1}{2} (\beta_0 \gamma_{m,0} + \gamma_{m,0}^2) L_{\mu t} \right] + \mathcal{O}(a_s^3), \quad (4.12)$$

expressing \bar{a}_s by a_s through eq. (4.8), and re-expanding in a_s . For $\zeta_1^{(2)}(t)$ one needs to multiply by eq. (4.12), and in addition by \bar{a}_s/a_s .

ζ_{13} and ζ_{23} require the more sophisticated renormalization given in eq. (4.6). It is important here to work consistently in d space-time dimensions. Since \vec{Z}_3 contains a $1/\epsilon$ pole already at $\mathcal{O}(a_s^0)$, we need to keep the $\mathcal{O}(\epsilon^2)$ terms of $\zeta_{2 \times 2}$ at NLO, and the $\mathcal{O}(\epsilon)$ terms at NNLO. They were not required in the calculation of ref. [21], so we recalculated $\zeta_{2 \times 2}$, keeping these higher terms in ϵ . Using the identity $n_A T_R = n_c C_F$ [75], our final result for $\vec{\zeta}_3$ reads:

$$\begin{aligned} \zeta_{13}(t) = & \frac{n_h n_c C_F}{16\pi^2} a_s \left\{ 5 - 6\zeta(2) - 6L_{\mu t} - 6L_{\mu t}^2 \right. \\ & + a_s \left[-3.31445 C_A - 27.5707 C_F - 15.0886 T_R n_h + 19.5780 T_R n_l \right. \\ & \quad \left. + L_{\mu t} \left(-5.68293 C_A - 32.7594 C_F - 4.17386 T_R n_h + 19.8261 T_R n_l \right) \right. \\ & \quad \left. + L_{\mu t}^2 \left(-\frac{433}{12} C_A - \frac{33}{2} C_F + \frac{26}{3} T_R n_f \right) \right. \\ & \quad \left. \left. + L_{\mu t}^3 \left(-\frac{22}{3} C_A - 6 C_F + \frac{8}{3} T_R n_f \right) \right] \right\} + \mathcal{O}(a_s^3), \end{aligned} \quad (4.13a)$$

$$\begin{aligned} \zeta_{23}(t) = & \frac{n_c n_h}{2\pi^2} \left\{ 1 + L_{\mu t} + a_s C_F \left[\frac{67}{16} + \frac{13}{2} \ln 2 - \frac{15}{2} \ln 3 - \frac{9}{4} \text{Li}_2(1/4) \right. \right. \\ & \quad \left. \left. + \left(\frac{33}{8} - \ln 2 - \frac{3}{4} \ln 3 \right) L_{\mu t} + \frac{3}{2} L_{\mu t}^2 \right] \right. \\ & + a_s^2 C_F \left[-0.710509 C_A + 6.97943 C_F - 6.43804 T_R n_h - 2.87689 T_R n_l \right. \\ & \quad \left. + L_{\mu t} \left(1.53754 C_A + 4.22899 C_F - 4.47865 T_R n_h - 1.47865 T_R n_l \right) \right. \\ & \quad \left. + L_{\mu t}^2 \left(2.57807 C_A + 4.09934 C_F - 0.931798 T_R n_f \right) \right. \\ & \quad \left. \left. + L_{\mu t}^3 \left(\frac{11}{24} C_A + \frac{3}{2} C_F - \frac{1}{6} T_R n_f \right) \right] \right\} + \mathcal{O}(a_s^3). \end{aligned} \quad (4.13b)$$

The logarithmic terms at $\mathcal{O}(a_s^n)$ are determined by the RG equation derived in section 5. Nevertheless, for the convenience of the reader, we provide the result for general μ in this case.

This, together with the results for $\zeta_{2 \times 2}$ obtained in refs. [21] and explicitly given in eq. (C.5), completes the result for the small-flow-time coefficients of the OPE up to dimension four of eq. (3.18).

5 Flow-time evolution

In the final section of this paper we derive a general flow-time evolution equation for flowed operators. It resembles the RG equation for regular operators but with a “flowed anomalous dimension matrix”. While studies of the relation between the RG and the flow-time evolution have also been performed elsewhere in the literature (see, e.g., refs. [26, 91–94]), to our knowledge the treatment described here has not been discussed before.

Let us return to the small-flow-time expansion of the operators \bar{O} defined in eqs. (3.15), (3.17), employing a matrix rather than component-wise notation for the sake of clarity:

$$\bar{O}(t) = \zeta^{\text{B}}(t)O = \zeta(t)O^{\text{R}}. \quad (5.1)$$

Since we work in the small-flow-time limit, the dependence of $\zeta(t)$ on t can be only through $L_{\mu t}$, defined in eq. (3.11). Taking the logarithmic derivative w.r.t. t of eq. (5.1), one thus obtains

$$t\partial_t\bar{O}(t) = (t\partial_t\zeta(t))O^{\text{R}}. \quad (5.2)$$

Using eq. (5.1) to eliminate the regular operators O^{R} , we find the flow-equation for flowed composite operators:

$$t\partial_t\bar{O}(t) = \gamma^{\text{f}}(t)\bar{O}(t), \quad \text{where} \quad \gamma^{\text{f}}(t) \equiv (t\partial_t\zeta(t))\zeta^{-1}(t). \quad (5.3)$$

So far the discussion is general and holds for any flowed OPE. Specializing to our case of the QCD dimension-four operators, we can write the “flowed anomalous dimension” matrix as

$$\begin{aligned} \gamma^{\text{f}} &= \begin{pmatrix} \gamma_{2\times 2}^{\text{f}} & \vec{\gamma}_3^{\text{f}} \\ 0 & 0 \end{pmatrix}, & \gamma_{2\times 2}^{\text{f}}(t) &= (t\partial_t\zeta_{2\times 2}(t))\zeta_{2\times 2}^{-1}(t), \\ & & \vec{\gamma}_3^{\text{f}}(t) &= -\gamma_{2\times 2}^{\text{f}}(t)\vec{\zeta}_3(t) + t\partial_t\vec{\zeta}_3(t). \end{aligned} \quad (5.4)$$

Through $\mathcal{O}(a_s^2)$, the result can be directly evaluated from eqs. (C.5) and (4.13). A consistency check is obtained by noting that $\zeta(t)$ depends on t only through $L_{\mu t}$:

$$t\partial_t\zeta(t) = \mu^2\frac{\partial}{\partial\mu^2}\zeta(t) = \mu^2\frac{\text{d}}{\text{d}\mu^2}\zeta(t) - a_s\beta\frac{\partial}{\partial a_s}\zeta(t). \quad (5.5)$$

On the other hand, we know that $a_s\tilde{O}_1(t)$ and $\tilde{O}_2(t)$ are RG invariant [10, 11, 20] and therefore, with eq. (3.15),

$$\begin{aligned} \begin{pmatrix} 0 \\ 0 \\ 4m^4\gamma_m \end{pmatrix} &= \mu^2\frac{\text{d}}{\text{d}\mu^2}H^{-1}(a_s)\tilde{O}(t) = \\ &= \mu^2\frac{\text{d}}{\text{d}\mu^2}H^{-1}(a_s)\left(\zeta^{(0)}(t)\mathbb{1} + \zeta^{(2)}(t)m^2\mathbb{1} + \zeta(t)O^{\text{R}}\right), \end{aligned} \quad (5.6)$$

where

$$H(x) = \begin{pmatrix} H_{2\times 2}(x) & \vec{0} \\ \vec{0}^T & 1 \end{pmatrix}, \quad \text{with} \quad H_{2\times 2}(x) = \begin{pmatrix} (4\pi^2x)^{-1} & 0 \\ 0 & 1 \end{pmatrix}. \quad (5.7)$$

Since operators of different mass dimensions do not mix under RG evolution and $\zeta_3^{(0,2)}(t) = 0$, we can drop the first two terms in the brackets on the r.h.s. of eq. (5.6).⁵ We thus arrive at

$$\mu^2 \frac{d}{d\mu^2} \zeta(t) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 4\gamma_m \end{pmatrix} - \begin{pmatrix} \beta & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \zeta(t) - \zeta(t) \gamma^O, \quad (5.8)$$

where γ^O is the anomalous dimension of the operators O^R , defined through

$$\mu^2 \frac{d}{d\mu^2} O^R = \gamma^O(a_s) O^R. \quad (5.9)$$

It can be written as

$$\gamma^O = \left(\mu^2 \frac{d}{d\mu^2} Z \right) Z^{-1} = \begin{pmatrix} \gamma_{2 \times 2}^O & \vec{\gamma}_3^O \\ \vec{0}^T & 4\gamma_m \end{pmatrix}, \quad (5.10)$$

with Z from eq. (2.10). Using the expressions of section 2.1, one derives [27, 28]

$$\begin{aligned} \gamma_{2 \times 2}^O &= \left(\mu^2 \frac{d}{d\mu^2} Z_{2 \times 2} \right) Z_{2 \times 2}^{-1} = \begin{pmatrix} -a_s \frac{\partial}{\partial a_s} \beta & -2 a_s \frac{\partial}{\partial a_s} \gamma_m \\ 0 & 0 \end{pmatrix}, \\ \vec{\gamma}_3^O &= Z_m^4 \left(\mu^2 \frac{d}{d\mu^2} \vec{Z}_3 - \gamma_{2 \times 2}^O \vec{Z}_3 \right) = \begin{pmatrix} 4 a_s \frac{\partial}{\partial a_s} \gamma_0 \\ 8 \gamma_0 \end{pmatrix}. \end{aligned} \quad (5.11)$$

The QCD renormalization group functions β and γ_m are defined in eqs. (A.2) and (A.4), respectively. Since they are of $\mathcal{O}(a_s)$, the explicit μ -dependence of $\zeta_{2 \times 2}(t)$ can be derived through $\mathcal{O}(a_s^3)$ from the results of ref. [21]. Thus, for $\gamma_{2 \times 2}^f$, eq. (5.5) is not just a consistency check, but a means to derive higher order terms. In our case, we can obtain the result through $\mathcal{O}(a_s^3)$:

$$\begin{aligned} \gamma_{11}^f &= a_s^2 \left[\frac{3}{32} C_A^2 + \frac{1}{8} C_A T_R n_f + \frac{7}{8} C_F T_R n_f \right] + a_s^3 \left[-\frac{7}{48} C_A T_R^2 n_f^2 \right. \\ &\quad - \frac{35}{36} C_F T_R^2 n_f^2 + C_A C_F T_R n_f \left(\frac{11891}{2880} + \frac{27}{40} \ln 2 - \frac{81}{160} \ln 3 \right) \\ &\quad + C_A^2 T_R n_f \left(\frac{1687}{2880} + \frac{29}{40} \ln 2 - \frac{9}{20} \ln 3 \right) + C_A^3 \left(\frac{6643}{11520} - \frac{319}{160} \ln 2 + \frac{99}{80} \ln 3 \right) \\ &\quad + C_F^2 T_R n_f \left(\frac{25}{64} + \frac{45}{8} \ln 2 - \frac{111}{32} \ln 3 + \frac{3}{4} \text{Li}_2(1/4) - \frac{3}{8} \zeta(2) \right) \\ &\quad \left. + \left(\frac{1}{6} C_A^2 T_R n_f + \frac{11}{64} C_A^3 + \frac{77}{48} C_A C_F T_R n_f - \frac{1}{12} C_A T_R^2 n_f^2 - \frac{7}{12} C_F T_R^2 n_f^2 \right) L_{\mu t} \right] \\ &\quad + \mathcal{O}(a_s^4), \end{aligned} \quad (5.12a)$$

⁵This can also be seen by noting that these two terms, multiplied by $H^{-1}(a_s)$, are $a_s \langle \tilde{O}_1(t) \rangle$ and $\langle \tilde{O}_2(t) \rangle$, expanded through order m^2 .

$$\gamma_{12}^f = -\frac{3}{2} a_s C_F + a_s^2 \left\{ -\frac{367}{48} C_A C_F + \frac{5}{3} C_F T_{Rn_f} + C_F^2 \left[-\frac{3}{16} - \frac{3}{2} \ln 2 - \frac{9}{8} \ln 3 \right] \right. \quad (5.12b)$$

$$\left. + \left[-\frac{11}{4} C_A C_F + C_F T_{Rn_f} \right] L_{\mu t} \right\}$$

$$+ a_s^3 \left\{ C_F^2 C_A \left[-\frac{391}{768} - \frac{431}{24} \ln 2 + \frac{3}{8} \ln^2 2 + \frac{11}{64} \ln 3 + \frac{9}{8} \text{Li}_2(1/4) - \frac{33}{64} \zeta(2) \right] \right.$$

$$+ C_F T_{Rn_f}^2 \left[-\frac{25}{18} - \frac{1}{2} \zeta(2) \right] + C_F^3 \left[-\frac{1401}{256} + \frac{339}{16} \ln 2 - \frac{9}{8} \ln^2 2 \right.$$

$$\left. - \frac{657}{64} \ln 3 - \frac{9}{4} \ln 2 \ln 3 - \frac{27}{32} \ln^2 3 + \frac{153}{32} \text{Li}_2(1/4) + \frac{39}{64} \zeta(2) \right]$$

$$+ C_A^2 C_F \left[-\frac{5291}{144} - \frac{2311}{24} \ln 2 + \frac{4641}{64} \ln 3 + \frac{11}{32} \text{Li}_2(1/4) + \frac{11}{8} \zeta(2) \right]$$

$$+ C_F^2 T_{Rn_f} \left[\frac{8827}{960} - \frac{2089}{120} \ln 2 + \frac{847}{80} \ln 3 + \frac{3}{8} \text{Li}_2(1/4) - \frac{3}{16} \zeta(2) - \frac{9}{2} \zeta(3) \right]$$

$$+ C_A C_F T_{Rn_f} \left[\frac{5861}{360} + \frac{4273}{120} \ln 2 - \frac{2139}{80} \ln 3 - \frac{1}{8} \text{Li}_2(1/4) + \frac{7}{8} \zeta(2) + \frac{9}{2} \zeta(3) \right]$$

$$+ \frac{3}{32} C_F c_\chi^{(2)} + \left[-\frac{4445}{192} C_A^2 C_F + \frac{647}{48} C_A C_F T_{Rn_f} - \frac{5}{3} C_F T_{Rn_f}^2 \right.$$

$$+ C_F^2 C_A \left(-\frac{33}{64} - \frac{33}{8} \ln 2 - \frac{99}{32} \ln 3 \right) + C_F^2 T_{Rn_f} \left(\frac{15}{16} + \frac{3}{2} \ln 2 + \frac{9}{8} \ln 3 \right) \left. \right] L_{\mu t}$$

$$+ \left[-\frac{121}{32} C_A^2 C_F + \frac{11}{4} C_A C_F T_{Rn_f} - \frac{1}{2} C_F T_{Rn_f}^2 \right] L_{\mu t}^2 \left. \right\} + \mathcal{O}(a_s^4),$$

$$\gamma_{21}^f = a_s^3 \left[C_A^2 T_{Rn_f} \left(\frac{599}{5760} + \frac{33}{80} \ln 2 - \frac{99}{320} \ln 3 \right) + C_A T_{Rn_f}^2 \left(\frac{41}{1440} \right. \quad (5.12c)$$

$$\left. - \frac{3}{20} \ln 2 + \frac{9}{80} \ln 3 \right) + C_A C_F T_{Rn_f} \left(\frac{209}{1152} + \frac{55}{16} \ln 2 - \frac{407}{192} \ln 3 \right. \right.$$

$$+ \frac{11}{24} \text{Li}_2(1/4) - \frac{11}{48} \zeta(2) \left. \right) + C_F T_{Rn_f}^2 \left(\frac{43}{144} - \frac{5}{4} \ln 2 + \frac{37}{48} \ln 3 \right. \right.$$

$$\left. - \frac{1}{6} \text{Li}_2(1/4) + \frac{1}{12} \zeta(2) \right) \left. \right] + \mathcal{O}(a_s^4),$$

$$\gamma_{22}^f = a_s^2 \left[C_A C_F \left(\frac{11}{96} - \frac{11}{12} \ln 2 - \frac{11}{16} \ln 3 \right) + C_F T_{Rn_f} \left(-\frac{2}{3} + \frac{1}{3} \ln 2 \right. \quad (5.12d)$$

$$\left. + \frac{1}{4} \ln 3 \right) \right] + a_s^3 \left\{ C_A C_F T_{Rn_f} \left[\frac{2521}{960} - \frac{1709}{90} \ln 2 - \frac{1}{6} \ln^2 2 \right. \right.$$

$$+ \frac{1537}{160} \ln 3 - \frac{1}{24} \text{Li}_2(1/4) \left. \right] + C_A^2 C_F \left[-\frac{7397}{2304} - \frac{213}{16} \ln 2 + \frac{11}{24} \ln^2 2 \right. \right.$$

$$+ \frac{107}{16} \ln 3 + \frac{11}{8} \text{Li}_2(1/4) - \frac{121}{192} \zeta(2) \left. \right] + C_F T_{Rn_f}^2 \left[-\frac{749}{720} + \frac{763}{90} \ln 2 \right. \right.$$

$$\left. - \frac{83}{20} \ln 3 - \frac{1}{6} \text{Li}_2(1/4) + \frac{1}{12} \zeta(2) \right] + C_F^2 T_{Rn_f} \left[-\frac{139}{192} - \frac{119}{8} \ln 2 \right.$$

$$\begin{aligned}
 & + \frac{1}{6} \ln^2 2 + \frac{261}{32} \ln 3 + \frac{1}{2} \ln 2 \ln 3 + \frac{3}{16} \ln^2 3 - \frac{23}{8} \text{Li}_2(1/4) + \frac{5}{48} \zeta(2) \Big] \\
 & + C_F^2 C_A \left[\frac{253}{384} + \frac{209}{8} \ln 2 - \frac{11}{24} \ln^2 2 - \frac{99}{8} \ln 3 - \frac{11}{8} \ln 2 \ln 3 - \frac{33}{64} \ln^2 3 \right. \\
 & \left. + \frac{187}{32} \text{Li}_2(1/4) + \frac{143}{192} \zeta(2) \right] + \left(\frac{11}{96} C_A - \frac{1}{24} T_R n_f \right) c_\chi^{(2)} \\
 & + \left[C_A^2 C_F \left(\frac{121}{576} - \frac{121}{72} \ln 2 - \frac{121}{96} \ln 3 \right) + C_F T_R^2 n_f^2 \left(\frac{4}{9} - \frac{2}{9} \ln 2 - \frac{1}{6} \ln 3 \right) \right. \\
 & \left. + C_A C_F T_R n_f \left(-\frac{187}{144} + \frac{11}{9} \ln 2 + \frac{11}{12} \ln 3 \right) \right] L_{\mu t} \Big\} + \mathcal{O}(a_s^4).
 \end{aligned}$$

We verified that this agrees through $\mathcal{O}(a_s^2)$ with the result which is obtained by directly inserting eq. (C.5) into eq. (5.4). Due to the factor of $1/g^2$ in \tilde{O}_1 (see eq. (3.1)), $\gamma_{2 \times 2}^f$ is not RG invariant, while $H_{2 \times 2}^{-1} \gamma_{2 \times 2}^f H_{2 \times 2}$ is. It may be useful to note that, by subtracting the VEVs off of \tilde{O}_1 and \tilde{O}_2 ,

$$\begin{aligned}
 \tilde{O}_{1,\text{sub}}(t, x) &= \tilde{O}_1(t, x) - \langle \tilde{O}_1(t, x) \rangle, \\
 \tilde{O}_{2,\text{sub}}(t, x) &= \tilde{O}_2(t, x) - \langle \tilde{O}_2(t, x) \rangle,
 \end{aligned} \tag{5.13}$$

the resulting operators do not mix with \tilde{O}_3 under t -evolution. Rather, their logarithmic t -evolution is fully governed by $\gamma_{2 \times 2}^f$ and thus known through $\mathcal{O}(a_s^3)$.

Eq. (5.8) does not analogously allow one to derive the $\mathcal{O}(a_s^3)$ terms of $\tilde{\gamma}_3^f$, because it involves γ_0 which, in contrast to β and γ_m , starts at $\mathcal{O}(a_s^0)$ rather than $\mathcal{O}(a_s)$, see eq. (A.7). Therefore, we can only give the result through $\mathcal{O}(a_s^2)$ for $\tilde{\gamma}_3^f$:

$$\begin{aligned}
 \gamma_{13}^f &= \frac{3n_c n_h}{8\pi^2} a_s C_F \left\{ 1 + a_s \left[9.24729 C_A - 2.47340 C_F \right. \right. \\
 & \left. \left. - 2.91787 T_R n_h + 1.08213 T_R n_l + L_{\mu t} \left(\frac{11}{6} C_A + 3 C_F - \frac{2}{3} T_R n_f \right) \right] \right\}, \tag{5.14a}
 \end{aligned}$$

$$\begin{aligned}
 \gamma_{23}^f &= \frac{n_c n_h}{2\pi^2} \left\{ 1 + a_s C_F \left[\frac{33}{8} - \ln 2 - \frac{3}{4} \ln 3 + 3 L_{\mu t} \right] + a_s^2 C_F \left[2.81363 C_A + 4.22899 C_F \right. \right. \\
 & \left. \left. - 4.31769 T_R n_h - 1.31769 T_R n_l + L_{\mu t} \left(6.43224 C_A + 8.19868 C_F - 1.70263 T_R n_f \right) \right] \right. \\
 & \left. + L_{\mu t}^2 \left(\frac{11}{8} C_A + \frac{9}{2} C_F - \frac{1}{2} T_R n_f \right) \right\}. \tag{5.14b}
 \end{aligned}$$

We checked, of course, that eq. (5.8) is consistent with the results for ζ_{13} and ζ_{23} of eq. (4.13).

6 Conclusions

We presented the flowed OPE for general current correlators and its matching to regular QCD through NNLO in the strong coupling α_s and through mass dimension four by using the small-flow-time expansion. Our calculation is based on the renormalization procedure

for the regular QCD dimension-four operators worked out in refs. [27, 28], the mixing matrix between flowed and regular operators derived in ref. [21], the method of projectors [66], and the tools and results for perturbative calculations in the GFF presented in ref. [65].

Overall, our results allow to combine the known perturbative results for the regular QCD current correlators from the literature to gradient-flow lattice calculations. This lays out the path for an alternative determination of hadronic contributions to observables such as the anomalous magnetic moment of the muon. In addition, we derived a general logarithmic flow-time evolution equation for flowed operators and presented its explicit form for the dimension-four operators considered in this paper.

Our methods are sufficiently general to be applied to similar problems at higher orders in perturbation theory, such as CP violating operators [22] relevant for the electric dipole moment of the neutron, or four-quark operators occurring in flavor physics [23].

Acknowledgments

We would like to thank A. Hoang for confirmative comments on the validity of the approach pursued in this paper, and Y. Kluth for helpful comments and clarifications. RVH would like to thank K. Chetyrkin for past initiation and collaboration on the material of section 2.

This work was supported by *Deutsche Forschungsgemeinschaft (DFG)* through project HA 2990/9-1, and by the U.S. Department of Energy under award No. DE-SC0008347. This document was prepared using the resources of the Fermi National Accelerator Laboratory (Fermilab), a U.S. Department of Energy, Office of Science, HEP User Facility. Fermilab is managed by Fermi Research Alliance, LLC (FRA), acting under Contract No. DE-AC02-07CH11359.

A Renormalization group functions

The d -dimensional beta function is defined as

$$\mu^2 \frac{d}{d\mu^2} a_s(\mu) = a_s(\mu) \beta_\epsilon(a_s(\mu)), \tag{A.1}$$

where $a_s \equiv \alpha_s/\pi \equiv g^2/(4\pi^2)$. The renormalized coupling $g = g(\mu)$ is related to the bare one through $g_B = \hat{\mu}^\epsilon Z_s^{1/2} g$, where Z_s is the $\overline{\text{MS}}$ renormalization constant. From this follow the relations

$$\begin{aligned} \beta_\epsilon(a_s) &= -\epsilon \left(1 + a_s \frac{\partial}{\partial a_s} \ln Z_s(a_s) \right)^{-1} = -\epsilon + \beta(a_s) \equiv -\epsilon - \sum_{n \geq 0} a_s^{n+1} \beta_n, \\ Z_s(a_s) &= 1 - \frac{a_s}{\epsilon} \beta_0 + a_s^2 \left(\frac{1}{\epsilon^2} \beta_0^2 - \frac{1}{2\epsilon} \beta_1 \right) + \mathcal{O}(a_s^3). \end{aligned} \tag{A.2}$$

Through section 5, we only need the first two perturbative coefficients, while β_2 is required in order to derive the $\mathcal{O}(a_s^3)$ terms of γ^f in eq. (5.12):

$$\begin{aligned}\beta_0 &= \frac{1}{4} \left(\frac{11}{3} C_A - \frac{4}{3} T_R n_f \right), & \beta_1 &= \frac{1}{16} \left(\frac{34}{3} C_A^2 - 4 C_F T_R n_f - \frac{20}{3} C_A T_R n_f \right), \\ \beta_2 &= \frac{1}{64} \left(\frac{2857}{54} C_A^3 + 2 C_F^2 T_R n_f - \frac{205}{9} C_F C_A T_R n_f - \frac{1415}{27} C_A^2 T_R n_f \right. \\ &\quad \left. + \frac{44}{9} C_F T_R^2 n_f^2 + \frac{158}{27} C_A T_R^2 n_f^2 \right).\end{aligned}\tag{A.3}$$

The anomalous dimension of the quark mass is defined through

$$\gamma_m(a_s) = -a_s \beta_\epsilon(a_s) \frac{\partial}{\partial a_s} \ln Z_m(a_s) \equiv - \sum_{n \geq 0} a_s^{n+1} \gamma_{m,n},\tag{A.4}$$

with the first three perturbative coefficients given by

$$\begin{aligned}\gamma_{m,0} &= \frac{3}{4} C_F, & \gamma_{m,1} &= \frac{3}{32} C_F^2 + \frac{97}{96} C_A C_F - \frac{5}{24} C_F T_R n_f, \\ \gamma_{m,2} &= \frac{1}{64} \left[\frac{129}{2} C_F^3 - \frac{129}{4} C_F^2 C_A + \frac{11413}{108} C_F C_A^2 \right. \\ &\quad \left. + C_F^2 T_R n_f (-46 + 48 \zeta(3)) + C_F C_A T_R n_f \left(-\frac{556}{27} - 48 \zeta(3) \right) - \frac{140}{27} C_F T_R^2 n_f^2 \right].\end{aligned}\tag{A.5}$$

It determines the $\overline{\text{MS}}$ renormalized mass m through

$$m_B = Z_m m, \quad Z_m = 1 - \frac{a_s}{\epsilon} \gamma_{m,0} + a_s^2 \left[\frac{\gamma_{m,0}}{2\epsilon^2} (\gamma_{m,0} + \beta_0) - \frac{1}{2\epsilon} \gamma_{m,1} \right] + \mathcal{O}(a_s^3).\tag{A.6}$$

Similarly to β_ϵ , the third coefficient $\gamma_{m,2}$ is needed only in section 5.

The renormalization constant of the vacuum energy Z_0 is related to the corresponding anomalous dimension γ_0 through

$$\gamma_0(a_s) = [4\gamma_m(a_s) - \epsilon] Z_0(a_s) + \beta_\epsilon(a_s) a_s \frac{\partial}{\partial a_s} Z_0(a_s) \equiv - \frac{n_c n_h}{(4\pi)^2} \sum_{n \geq 0} a_s^n \gamma_{0,n},\tag{A.7}$$

which leads to

$$\begin{aligned}Z_0(a_s) &= \frac{n_c n_h}{(4\pi)^2 \epsilon} \left\{ 1 + a_s \left(\frac{\gamma_{0,1}}{2} - \frac{2\gamma_{m,0}}{\epsilon} \right) + a_s^2 \left[\frac{2}{3\epsilon^2} (\beta_0 \gamma_{m,0} + 4\gamma_{m,0}^2) \right. \right. \\ &\quad \left. \left. - \frac{1}{6\epsilon} (\beta_0 \gamma_{0,1} + 4\gamma_{0,1} \gamma_{m,0} + 8\gamma_{m,1}) + \frac{1}{3} \gamma_{0,2} \right] \right\} + \mathcal{O}(a_s^3).\end{aligned}\tag{A.8}$$

The first three perturbative coefficients are given by [28, 62]⁶

$$\begin{aligned}\gamma_{0,0} &= 1, & \gamma_{0,1} &= C_F, \\ \gamma_{0,2} &= -C_F^2 \left(\frac{131}{32} - 3\zeta(3) \right) - C_F C_A \left(-\frac{109}{32} + \frac{3}{2} \zeta(3) \right) - C_F T_R \left(\frac{5}{8} n_f + 3n_h \right).\end{aligned}\tag{A.9}$$

⁶Higher orders have been computed in ref. [95].

B Perturbative coefficient functions

This appendix cites the results for the coefficient functions C_n of the regular dimension-four operators appearing in the OPE of the current correlators defined in eq. (2.2). We consider scalar, pseudo-scalar, vector- and axial-vector currents, both diagonal and non-diagonal, i.e. the currents assume the form

$$j(x) = \bar{\psi}_k(x)\Gamma\psi_l(x), \quad \Gamma \in \{1, i\gamma_5, \gamma^\mu, \gamma^\mu\gamma_5\}, \quad (\text{B.1})$$

$$k, l \in N \cup M, \quad M = \{1, \dots, n_h\}, \quad N = \{n_h + 1, \dots, n_f\}.$$

This means that ψ_k and ψ_l can be either both massive with mass m ($k, l \in M$), or both massless ($k, l \in N$), or one of them is massless, the other massive (e.g. $k \in M, l \in N$). While C_1 is independent of k and l , the coefficient C_2 of the quark operator takes the form

$$C_2 = C_{2,N} + \frac{1}{n_h} (\delta_{kM} + \delta_{lM}) (C_{2,M} + C_{2,\text{nd}}), \quad (\text{B.2})$$

where $\delta_{kM} = 1$ for $k \in M$, and 0 otherwise. Also the results for C_3 depend on whether the quarks k and l are massive or not. This dependence will be indicated explicitly below, using the δ_{kM} symbol defined above.

For convenience, we introduce the short-hand notation

$$l_{\mu Q} \equiv \ln \frac{Q^2}{\mu^2}, \quad (\text{B.3})$$

and the dimensionless quantities

$$\hat{C}_1 = Q^4 C_1, \quad \hat{C}_2 = -2 Q^4 C_2, \quad \hat{C}_3 = Q^4 C_3. \quad (\text{B.4})$$

The extra factor (-2) between C_2 and \hat{C}_2 arises from using O'_2 in ref. [64] rather than O_2 from eq. (2.5). For the sake of brevity, we insert the SU(3) color factors.

B.1 Vector and axial-vector currents

In this case, the current correlator can be decomposed into a transversal and a longitudinal part, each associated with a set of coefficient functions:

$$\int d^4x e^{iqx} j_\mu^{v/a}(x) j_\nu^{v/a}(0) \xrightarrow{Q^2 \rightarrow \infty} \sum_n \left((g_{\mu\nu} Q^2 - Q_\mu Q_\nu) C_n^{v/a,\Gamma} - Q_\mu Q_\nu C_n^{v/a,L} \right) O_n^R. \quad (\text{B.5})$$

The upper sign refers to the vector, the lower sign to the axial-vector case. The results have been taken from ref. [64]:

$$\hat{C}_1^{v/a,\Gamma} = \frac{1}{12} a_s + \frac{7}{72} a_s^2, \quad (\text{B.6a})$$

$$\hat{C}_{2,N}^{v/a,\Gamma} = a_s^2 \left(-1 + \frac{1}{3} l_{\mu Q} + \frac{4}{3} \zeta(3) \right), \quad (\text{B.6b})$$

$$\hat{C}_{2,M}^{v/a,\Gamma} = -a_s + a_s^2 \left[-\frac{29}{6} + \frac{1}{6} n_f + l_{\mu Q} \left(-\frac{11}{4} + \frac{1}{6} n_f \right) \right], \quad (\text{B.6c})$$

$$\hat{C}_{2,\text{nd}}^{v/a,\Gamma} = \pm \left(1 + \frac{4}{3} a_s + a_s^2 \left[\frac{191}{18} - \frac{7}{27} n_f + l_{\mu Q} \left(\frac{11}{3} - \frac{2}{9} n_f \right) \right] \right), \quad (\text{B.6d})$$

$$\hat{C}_1^{v/a,\text{L}} = 0, \quad \hat{C}_{2,N}^{v/a,\text{L}} = 0, \quad \hat{C}_{2,M}^{v/a,\text{L}} = 1, \quad \hat{C}_{2,\text{nd}}^{v/a,\text{L}} = \mp 1, \quad (\text{B.6e})$$

$$\hat{C}_3^{v/a,\Gamma} = \frac{3}{16\pi^2} \left\{ \delta_{kM} \delta_{lM} \left[a_s \left(\frac{152}{9} - \frac{32}{3} \zeta(3) \right) \right. \right. \quad (\text{B.6f})$$

$$\left. + a_s^2 \left[\frac{1295}{9} - \frac{524}{3} \zeta(3) + 120 \zeta(5) + n_f \left(-\frac{362}{81} + \frac{8}{27} \zeta(3) \right) \right. \right.$$

$$\left. + l_{\mu Q} \left(114 - 72 \zeta(3) + n_f \left(-\frac{76}{27} + \frac{16}{9} \zeta(3) \right) \right) \right]$$

$$\pm 2\delta_{kM} \delta_{lM} \left[-4l_{\mu Q} + a_s \left(-\frac{56}{3} + 16 \zeta(3) - \frac{32}{3} l_{\mu Q} - 8l_{\mu Q}^2 \right) \right.$$

$$\left. + a_s^2 \left[-\frac{18967}{108} + \frac{4588}{27} \zeta(3) + \frac{4}{3} \zeta(4) - \frac{280}{27} \zeta(5) + n_f \left(\frac{337}{54} - \frac{40}{9} \zeta(3) \right) \right. \right.$$

$$\left. + l_{\mu Q} \left(-\frac{3617}{18} + \frac{332}{3} \zeta(3) + n_f \left(\frac{157}{27} - \frac{8}{3} \zeta(3) \right) \right) \right]$$

$$\left. + l_{\mu Q}^2 \left(-77 + \frac{22}{9} n_f \right) + l_{\mu Q}^3 \left(-18 + \frac{4}{9} n_f \right) \right]$$

$$+ (\delta_{kM} + \delta_{lM}) \left[-2 + a_s \left(-4 - 4l_{\mu Q} \right) \right.$$

$$\left. + a_s^2 \left[-\frac{776}{27} + \frac{2996}{27} \zeta(3) - \frac{3440}{27} \zeta(5) + n_f \left(\frac{23}{27} - \frac{4}{9} \zeta(3) \right) \right. \right.$$

$$\left. + l_{\mu Q} \left(-36 + \frac{10}{9} n_f \right) - 8l_{\mu Q}^2 \right]$$

$$+ n_h a_s^2 \left[-\frac{80}{9} + \frac{16}{3} \zeta(3) + l_{\mu Q} \left(\frac{40}{9} - \frac{16}{3} \zeta(3) \right) - \frac{2}{3} l_{\mu Q}^2 \right]$$

$$+ n_h a_s^2 \left[(\delta_{kM} + \delta_{lM}) \left(\frac{100}{9} - \frac{16}{3} \zeta(3) \right) \right.$$

$$\left. \pm \delta_{kM} \delta_{lM} \left(-\frac{64}{9} + 16l_{\mu Q} + \frac{32}{3} \zeta(3) \right) \right] \left. \right\},$$

$$\hat{C}_3^{v/a,\text{L}} = \frac{3}{16\pi^2} \left\{ \delta_{kM} \delta_{lM} \left[8 + a_s \left(\frac{152}{3} + 32l_{\mu Q} \right) \right. \right. \quad (\text{B.6g})$$

$$\left. + a_s^2 \left[\frac{7306}{9} - \frac{1292}{9} \zeta(3) - \frac{640}{9} \zeta(5) + n_f \left(-\frac{212}{9} + \frac{8}{3} \zeta(3) \right) \right. \right.$$

$$\left. + l_{\mu Q} \left(\frac{1430}{3} - \frac{116}{9} n_f \right) + l_{\mu Q}^2 \left(108 - \frac{8}{3} n_f \right) \right]$$

$$\pm 2\delta_{kM} \delta_{lM} \left[4l_{\mu Q} + a_s \left(8 - 16 \zeta(3) + \frac{16}{3} l_{\mu Q} + 8l_{\mu Q}^2 \right) \right]$$

$$\begin{aligned}
 & + a_s^2 \left[-\frac{6713}{108} - \frac{464}{3} \zeta(3) - \frac{4}{3} \zeta(4) + \frac{940}{9} \zeta(5) + n_f \left(\frac{31}{54} + \frac{8}{9} \zeta(3) \right) \right. \\
 & + l_{\mu Q} \left(\frac{1429}{18} - \frac{332}{3} \zeta(3) + n_f \left(-3 + \frac{8}{3} \zeta(3) \right) \right) + l_{\mu Q}^2 \left(\frac{155}{3} - \frac{14}{9} n_f \right) \\
 & \left. + l_{\mu Q}^3 \left(18 - \frac{4}{9} n_f \right) \right] \\
 & + (\delta_{kM} + \delta_{lM}) \left[-4 - 4l_{\mu Q} + a_s \left(-\frac{100}{3} + 16 \zeta(3) - \frac{64}{3} l_{\mu Q} - 8 l_{\mu Q}^2 \right) \right. \\
 & + a_s^2 \left[-\frac{37123}{108} + \frac{2038}{9} \zeta(3) + \frac{4}{3} \zeta(4) - \frac{620}{9} \zeta(5) + n_f \left(\frac{605}{54} - \frac{20}{9} \zeta(3) \right) \right. \\
 & + l_{\mu Q} \left(-\frac{5719}{18} + n_f \left(\frac{85}{9} - \frac{8}{3} \zeta(3) \right) + \frac{332}{3} \zeta(3) \right) \\
 & \left. + l_{\mu Q}^2 \left(-\frac{317}{3} + \frac{26}{9} n_f \right) + l_{\mu Q}^3 \left(-18 + \frac{4}{9} n_f \right) \right] \\
 & \left. + a_s^2 n_h \left[\left(\frac{32}{9} + 8 l_{\mu Q} \right) (\delta_{kM} + \delta_{lM}) \pm \left(-\frac{64}{9} - 16 l_{\mu Q} \right) \delta_{kM} \delta_{lM} \right] \right\}.
 \end{aligned}$$

B.2 Scalar and pseudo-scalar currents

Also the results for the scalar and the pseudo-scalar currents (upper and lower signs, respectively) are taken from ref. [64]:

$$\hat{C}_1^{s/p} = \frac{1}{8} a_s + a_s^2 \left[\frac{11}{16} + \frac{1}{4} l_{\mu Q} \right], \quad (\text{B.7a})$$

$$\hat{C}_{2,N}^{s/p} = a_s^2 \left[-\frac{5}{6} + \frac{1}{2} l_{\mu Q} \right], \quad (\text{B.7b})$$

$$\begin{aligned}
 \hat{C}_{2,M}^{s/p} &= \frac{1}{2} + a_s \left[\frac{11}{6} + l_{\mu Q} \right] \\
 & + a_s^2 \left[\frac{5437}{288} - \frac{17}{4} \zeta(3) - \frac{79}{144} n_f + l_{\mu Q} \left(\frac{155}{12} - \frac{4}{9} n_f \right) + l_{\mu Q}^2 \left(\frac{19}{8} - \frac{1}{12} n_f \right) \right], \quad (\text{B.7c})
 \end{aligned}$$

$$\begin{aligned}
 \hat{C}_{2,\text{nd}}^{s/p} &= \pm \left(1 + a_s \left[\frac{14}{3} + 2 l_{\mu Q} \right] \right. \\
 & \left. + a_s^2 \left[\frac{7549}{144} - \frac{15}{2} \zeta(3) - \frac{41}{24} n_f + l_{\mu Q} \left(\frac{367}{12} - \frac{19}{18} n_f \right) + l_{\mu Q}^2 \left(\frac{19}{4} - \frac{1}{6} n_f \right) \right] \right), \quad (\text{B.7d})
 \end{aligned}$$

$$\begin{aligned}
 \hat{C}_3^{s/p} &= \frac{3}{16\pi^2} \left\{ \delta_{kM} \delta_{lM} \left[4 + a_s \left[\frac{32}{3} + 24 l_{\mu Q} + 16 \zeta(3) \right] \right. \right. \\
 & + a_s^2 \left[\frac{9955}{36} + \frac{724}{9} \zeta(3) - \frac{610}{9} \zeta(5) + n_f \left(-\frac{463}{54} + \frac{32}{9} \zeta(3) \right) \right. \\
 & \left. \left. + l_{\mu Q} \left(\frac{583}{3} + 140 \zeta(3) + n_f \left(-\frac{46}{9} - \frac{8}{3} \zeta(3) \right) \right) + l_{\mu Q}^2 (105 - 2 n_f) \right] \right] \\
 & \pm 2 \delta_{kM} \delta_{lM} \left[-4 l_{\mu Q} + a_s \left[-16 + 16 \zeta(3) - 24 l_{\mu Q} - 16 l_{\mu Q}^2 \right] \right]
 \end{aligned} \quad (\text{B.7e})$$

$$\begin{aligned}
 & + a_s^2 \left[-\frac{19003}{108} + \frac{574}{3} \zeta(3) + \frac{4}{3} \zeta(4) + \frac{50}{9} \zeta(5) + n_f \left(\frac{245}{54} + \frac{16}{9} \zeta(3) \right) \right. \\
 & + l_{\mu Q} \left(-\frac{14399}{36} + \frac{518}{3} \zeta(3) + n_f \left(\frac{67}{6} - \frac{8}{3} \zeta(3) \right) \right) \\
 & \left. + l_{\mu Q}^2 \left(-222 + \frac{52}{9} n_f \right) + l_{\mu Q}^3 \left(-53 + \frac{10}{9} n_f \right) \right] \\
 & + (\delta_{kM} + \delta_{lM}) \left[1 - 2l_{\mu Q} + a_s \left[-6 + 8\zeta(3) - 4l_{\mu Q} - 8l_{\mu Q}^2 \right] \right. \\
 & + a_s^2 \left[-\frac{13343}{432} + \frac{155}{6} \zeta(3) + \frac{2}{3} \zeta(4) + \frac{190}{9} \zeta(5) + n_f \left(\frac{205}{216} + \frac{20}{9} \zeta(3) \right) \right. \\
 & + l_{\mu Q}^2 \left(-\frac{285}{4} + \frac{37}{18} n_f \right) + l_{\mu Q}^3 \left(-\frac{53}{2} + \frac{5}{9} n_f \right) \\
 & \left. + l_{\mu Q} \left(-\frac{9017}{72} + \frac{265}{3} \zeta(3) + n_f \left(\frac{115}{36} - \frac{4}{3} \zeta(3) \right) \right) \right] \\
 & + n_h a_s^2 \left[-5 + 4\zeta(3) + 4l_{\mu Q} - l_{\mu Q}^2 \right] \\
 & + n_h a_s^2 \left[(\delta_{kM} + \delta_{lM}) \left(-\frac{86}{9} - 8\zeta(3) + 4l_{\mu Q} \right) \right. \\
 & \left. \pm \delta_{kM} \delta_{lM} \left(-\frac{32}{9} - 16\zeta(3) + 16l_{\mu Q} \right) \right] \Bigg\}.
 \end{aligned}$$

C Renormalized mixing matrix

For the reader's convenience, we provide the relation between the mixing matrix ζ used in this paper and its definition in ref. [21], referred to as ‘‘HKL’’ in what follows. This is easily derived from the relation between the operators O_n defined in eq. (2.5) and the $O_{n,\mu\nu}$ of HKL:

$$\begin{pmatrix} O_1 \\ O_2 \end{pmatrix} \delta_{\mu\nu} = H_{2 \times 2}(a_s^B) \begin{pmatrix} O_{2,\mu\nu} \\ O_{4,\mu\nu} \end{pmatrix}, \quad (\text{C.1})$$

with $H_{2 \times 2}$ defined in eq. (5.7). The mixing matrix between regular and flowed operators in HKL, restricted to the two operators which are relevant for this paper, is defined through

$$\begin{pmatrix} \tilde{O}_{2,\mu\nu}(t) \\ \tilde{O}_{4,\mu\nu}(t) \end{pmatrix} = \zeta_{2 \times 2}^{\text{HKL}}(t) \begin{pmatrix} O_{2,\mu\nu} \\ O_{4,\mu\nu} \end{pmatrix}, \quad \zeta_{2 \times 2}^{\text{HKL}} = \begin{pmatrix} \zeta_{22}^{\text{HKL}} & \zeta_{24}^{\text{HKL}} \\ \zeta_{42}^{\text{HKL}} & \zeta_{44}^{\text{HKL}} \end{pmatrix}, \quad (\text{C.2})$$

where the ζ_{ij}^{HKL} are the entries of the full 4×4 mixing matrix of HKL. Inserting eq. (C.1) and

$$\begin{pmatrix} \tilde{O}_1(t) \\ \tilde{O}_2(t) \end{pmatrix} \delta_{\mu\nu} = H_{2 \times 2}(\hat{\mu}^{2\epsilon} a_s) \chi(t) \begin{pmatrix} \tilde{O}_{2,\mu\nu}(t) \\ \tilde{O}_{4,\mu\nu}(t) \end{pmatrix}, \quad \text{with} \quad \chi(t) = \begin{pmatrix} 1 & 0 \\ 0 & \zeta_\chi(t) \end{pmatrix}, \quad (\text{C.3})$$

with $\zeta_\chi(t)$ from eq. (3.10), gives the renormalized mixing matrix used in the current paper in terms of the bare mixing matrix of HKL:

$$\zeta_{2 \times 2}(t) = H_{2 \times 2}(a_s \hat{u}^{2\epsilon}) \chi(t) \zeta_{2 \times 2}^{\text{HKL}}(t) H_{2 \times 2}^{-1}(a_s^{\text{B}}) Z_{2 \times 2}^{-1}(a_s). \quad (\text{C.4})$$

Explicitly, one finds:

$$\zeta_{11}(t) = 1 + \frac{7}{8} a_s C_A + a_s^2 \left\{ C_A^2 \left[\frac{227}{180} - \frac{87}{80} \ln 2 + \frac{27}{40} \ln 3 + \frac{3}{32} L_{\mu t} \right] \right. \\ \left. + C_A T_{\text{R}n_f} \left[-\frac{1}{18} + \frac{1}{8} L_{\mu t} \right] + C_F T_{\text{R}n_f} \left[\frac{3}{16} + \frac{1}{4} L_{\mu t} \right] \right\}, \quad (\text{C.5a})$$

$$\zeta_{12}(t) = a_s C_F \left[-\frac{5}{4} - \frac{3}{2} L_{\mu t} \right] + a_s^2 \left[C_F^2 \left(-\frac{17}{32} - \frac{3}{8} L_{\mu t} \right) \right. \\ \left. + C_A C_F \left(-\frac{431}{60} + \frac{1}{2} \zeta(2) - \frac{4273}{120} \ln 2 + \frac{2139}{80} \ln 3 + \frac{1}{8} \text{Li}_2(1/4) \right. \right. \\ \left. \left. - \frac{367}{48} L_{\mu t} - \frac{11}{8} L_{\mu t}^2 \right) + C_F T_{\text{R}n_f} \left(\frac{15}{8} + \frac{1}{2} \zeta(2) + \frac{5}{3} L_{\mu t} + \frac{1}{2} L_{\mu t}^2 \right) \right], \quad (\text{C.5b})$$

$$\zeta_{21}(t) = \frac{5}{12} a_s T_{\text{R}n_f} + a_s^2 \left[C_A T_{\text{R}n_f} \left(\frac{209}{480} + \frac{9}{20} \ln 2 - \frac{27}{80} \ln 3 \right) \right. \\ \left. + C_F T_{\text{R}n_f} \left(\frac{1}{4} + \frac{1}{2} \text{Li}_2(1/4) - \frac{1}{4} \zeta(2) + \frac{10}{3} \ln 2 - \frac{21}{8} \ln 3 \right) \right], \quad (\text{C.5c})$$

$$\zeta_{22}(t) = 1 + a_s C_F \left[\frac{1}{8} - \ln 2 - \frac{3}{4} \ln 3 \right] \\ + a_s^2 \left\{ C_A C_F \left[-\frac{691}{384} - \frac{55}{8} \ln 2 + \frac{1}{4} \ln^2 2 + \frac{63}{16} \ln 3 \right. \right. \\ \left. \left. + \left(\frac{11}{96} - \frac{11}{12} \ln 2 - \frac{11}{16} \ln 3 \right) L_{\mu t} + \frac{3}{4} \text{Li}_2(1/4) - \frac{11}{32} \zeta(2) \right] \right. \\ \left. + C_F T_{\text{R}n_f} \left[\frac{333}{160} - \frac{763}{60} \ln 2 + \frac{249}{40} \ln 3 + \left(-\frac{2}{3} + \frac{1}{3} \ln 2 + \frac{1}{4} \ln 3 \right) L_{\mu t} \right. \right. \\ \left. \left. + \frac{1}{4} \text{Li}_2(1/4) - \frac{1}{8} \zeta(2) \right] + C_F^2 \left[\frac{47}{128} + \frac{113}{8} \ln 2 + \frac{1}{4} \ln^2 2 - \frac{219}{32} \ln 3 \right. \right. \\ \left. \left. + \frac{51}{16} \text{Li}_2(1/4) + \frac{13}{32} \zeta(2) \right] + \frac{1}{16} c_\chi^{(2)} \right\}, \quad (\text{C.5d})$$

where $c_\chi^{(2)}$ is given in eq. (3.13). The results including higher orders in ϵ are provided in the supplementary material, see appendix D.

D Supplementary material

The main results of this paper are provided in computer readable format (e.g. with `Mathematica` [90]). The notation is described in table 1. All coefficients are represented by floating-point numbers in this file. The relative uncertainty for our numerically evaluated coefficients is estimated to be 10^{-10} or better.

	$\zeta_1^{(0)}$	$\zeta_1^{(2)}$	ζ_{13}	$\zeta_2^{(0)}$	$\zeta_2^{(2)}$	ζ_{23}
Eq.	(4.7)	(4.11a)	(4.13a)	(4.9)	(4.11b)	(4.13b)
code	zeta01anc	zeta21anc	zeta13anc	zeta02anc	zeta22anc	zeta23anc

	$\gamma_{2 \times 2}^f$	$\vec{\gamma}_3^f$	$\zeta_{2 \times 2}$
Eq.	(5.12)	(5.14)	(C.5)
code	gamma22anc	gamma3anc	ZetaMatrix22anc

Table 1. Notation in the supplementary material.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] K.G. Chetyrkin, J.H. Kühn and A. Kwiatkowski, *QCD corrections to the e^+e^- cross-section and the Z boson decay rate*, *Phys. Rept.* **277** (1996) 189 [[hep-ph/9503396](#)] [[INSPIRE](#)].
- [2] F. Jegerlehner, *The anomalous magnetic moment of the muon*, Springer, Berlin, Heidelberg, Germany (2017).
- [3] T. Aoyama et al., *The anomalous magnetic moment of the muon in the Standard Model*, [arXiv:2006.04822](#) [[INSPIRE](#)].
- [4] K. Chetyrkin et al., *Precise charm- and bottom-quark masses: theoretical and experimental uncertainties*, *Theor. Math. Phys.* **170** (2012) 217 [[arXiv:1010.6157](#)] [[INSPIRE](#)].
- [5] A. Crivellin, M. Hoferichter, C.A. Manzari and M. Montull, *Hadronic vacuum polarization: $(g-2)_\mu$ versus global electroweak fits*, [arXiv:2003.04886](#) [[INSPIRE](#)].
- [6] A. Keshavarzi, W.J. Marciano, M. Passera and A. Sirlin, *The muon $g-2$ and $\Delta\alpha$ connection*, [arXiv:2006.12666](#) [[INSPIRE](#)].
- [7] S. Borsányi et al., *Leading-order hadronic vacuum polarization contribution to the muon magnetic moment from lattice QCD*, [arXiv:2002.12347](#) [[INSPIRE](#)].
- [8] T. Blum et al., *Hadronic light-by-light scattering contribution to the muon anomalous magnetic moment from lattice QCD*, *Phys. Rev. Lett.* **124** (2020) 132002 [[arXiv:1911.08123](#)] [[INSPIRE](#)].
- [9] E.-H. Chao, A. Gérardin, J.R. Green, R.J. Hudspith and H.B. Meyer, *Hadronic light-by-light contribution to $(g-2)_\mu$ from lattice QCD with SU(3) flavor symmetry*, [arXiv:2006.16224](#) [[INSPIRE](#)].
- [10] M. Lüscher, *Properties and uses of the Wilson flow in lattice QCD*, *JHEP* **08** (2010) 071 [*Erratum ibid.* **03** (2014) 092] [[arXiv:1006.4518](#)] [[INSPIRE](#)].
- [11] M. Lüscher and P. Weisz, *Perturbative analysis of the gradient flow in non-Abelian gauge theories*, *JHEP* **02** (2011) 051 [[arXiv:1101.0963](#)] [[INSPIRE](#)].

- [12] M. Lüscher, *Chiral symmetry and the Yang-Mills gradient flow*, *JHEP* **04** (2013) 123 [[arXiv:1302.5246](#)] [[INSPIRE](#)].
- [13] R. Narayanan and H. Neuberger, *Infinite N phase transitions in continuum Wilson loop operators*, *JHEP* **03** (2006) 064 [[hep-th/0601210](#)] [[INSPIRE](#)].
- [14] M. Lüscher, *Trivializing maps, the Wilson flow and the HMC algorithm*, *Commun. Math. Phys.* **293** (2010) 899 [[arXiv:0907.5491](#)] [[INSPIRE](#)].
- [15] S. Borsányi et al., *High-precision scale setting in lattice QCD*, *JHEP* **09** (2012) 010 [[arXiv:1203.4469](#)] [[INSPIRE](#)].
- [16] R. Sommer, *Scale setting in lattice QCD*, *PoS(LATTICE2013)015* (2014) [[arXiv:1401.3270](#)] [[INSPIRE](#)].
- [17] C. Monahan and K. Orginos, *Locally-smearred operator product expansions*, *PoS(LATTICE2014)330* (2014) [[arXiv:1410.3393](#)] [[INSPIRE](#)].
- [18] C. Monahan and K. Orginos, *Locally smeared operator product expansions in scalar field theory*, *Phys. Rev. D* **91** (2015) 074513 [[arXiv:1501.05348](#)] [[INSPIRE](#)].
- [19] H. Suzuki, *Energy-momentum tensor from the Yang-Mills gradient flow*, *PTEP* **2013** (2013) 083B03 [*Erratum ibid.* **2015** (2015) 079201] [[arXiv:1304.0533](#)] [[INSPIRE](#)].
- [20] H. Makino and H. Suzuki, *Lattice energy-momentum tensor from the Yang-Mills gradient flow — inclusion of fermion fields*, *PTEP* **2014** (2014) 063B02 [*Erratum ibid.* **2015** (2015) 079202] [[arXiv:1403.4772](#)] [[INSPIRE](#)].
- [21] R.V. Harlander, Y. Kluth and F. Lange, *The two-loop energy-momentum tensor within the gradient-flow formalism*, *Eur. Phys. J. C* **78** (2018) 944 [*Erratum ibid.* **79** (2019) 858] [[arXiv:1808.09837](#)] [[INSPIRE](#)].
- [22] M.D. Rizik, C.J. Monahan and A. Shindler, *Short flow-time coefficients of CP-violating operators*, [arXiv:2005.04199](#) [[INSPIRE](#)].
- [23] A. Suzuki, Y. Taniguchi, H. Suzuki and K. Kanaya, *Four quark operators for kaon bag parameter with gradient flow*, [arXiv:2006.06999](#) [[INSPIRE](#)].
- [24] C.A. Dominguez, *Analytical determination of the QCD quark masses*, *Int. J. Mod. Phys. A* **29** (2014) 1430069 [[arXiv:1411.3462](#)] [[INSPIRE](#)].
- [25] R.V. Harlander, S.Y. Klein and M. Lipp, *FeynGame*, *Comput. Phys. Commun.* **256** (2020) 107465 [[arXiv:2003.00896](#)] [[INSPIRE](#)].
- [26] H. Makino, O. Morikawa and H. Suzuki, *Gradient flow and the Wilsonian renormalization group flow*, *PTEP* **2018** (2018) 053B02 [[arXiv:1802.07897](#)] [[INSPIRE](#)].
- [27] V.P. Spiridonov, *Anomalous dimension of $G_{\mu\nu}^2$ and β function*, report no. IYAI-P-0378, (1984) [[INSPIRE](#)].
- [28] V.P. Spiridonov and K.G. Chetyrkin, *Nonleading mass corrections and renormalization of the operators $m\bar{\psi}\psi$ and $G_{\mu\nu}^2$* , *Sov. J. Nucl. Phys.* **47** (1988) 522 [*Yad. Fiz.* **47** (1988) 818] [[INSPIRE](#)].
- [29] A.O.G. Källén and A. Sabry, *Fourth order vacuum polarization*, *Kong. Dan. Vid. Sel. Mat. Fys. Med.* **29** (1955) 1 [[INSPIRE](#)].
- [30] D.J. Broadhurst, *Chiral symmetry breaking and perturbative QCD*, *Phys. Lett. B* **101** (1981) 423 [[INSPIRE](#)].

- [31] B.A. Kniehl, *Two loop corrections to the vacuum polarizations in perturbative QCD*, *Nucl. Phys. B* **347** (1990) 86 [INSPIRE].
- [32] A. Djouadi and P. Gambino, *Electroweak gauge bosons selfenergies: complete QCD corrections*, *Phys. Rev. D* **49** (1994) 3499 [Erratum *ibid.* **53** (1996) 4111] [hep-ph/9309298] [INSPIRE].
- [33] D.J. Broadhurst, J. Fleischer and O.V. Tarasov, *Two loop two point functions with masses: asymptotic expansions and Taylor series, in any dimension*, *Z. Phys. C* **60** (1993) 287 [hep-ph/9304303] [INSPIRE].
- [34] J. Fleischer and O.V. Tarasov, *Calculation of Feynman diagrams from their small momentum expansion*, *Z. Phys. C* **64** (1994) 413 [hep-ph/9403230] [INSPIRE].
- [35] D.J. Broadhurst, P.A. Baikov, V.A. Ilyin, J. Fleischer, O.V. Tarasov and V.A. Smirnov, *Two loop gluon condensate contributions to heavy quark current correlators: exact results and approximations*, *Phys. Lett. B* **329** (1994) 103 [hep-ph/9403274] [INSPIRE].
- [36] P.A. Baikov and D.J. Broadhurst, *Three loop QED vacuum polarization and the four loop muon anomalous magnetic moment*, in *4th International Workshop on Software Engineering and Artificial Intelligence for high-energy and nuclear physics*, (1995), pg. 0167 [hep-ph/9504398] [INSPIRE].
- [37] K.G. Chetyrkin, J.H. Kühn and M. Steinhauser, *Heavy quark vacuum polarization to three loops*, *Phys. Lett. B* **371** (1996) 93 [hep-ph/9511430] [INSPIRE].
- [38] K.G. Chetyrkin, J.H. Kühn and M. Steinhauser, *Heavy quark current correlators to $O(\alpha_s^2)$* , *Nucl. Phys. B* **505** (1997) 40 [hep-ph/9705254] [INSPIRE].
- [39] K.G. Chetyrkin, R. Harlander and M. Steinhauser, *Singlet polarization functions at $O(\alpha_s^2)$* , *Phys. Rev. D* **58** (1998) 014012 [hep-ph/9801432] [INSPIRE].
- [40] A.H. Hoang, V. Mateu and S. Mohammad Zebarjad, *Heavy quark vacuum polarization function at $O(\alpha_s^2)O(\alpha_s^3)$* , *Nucl. Phys. B* **813** (2009) 349 [arXiv:0807.4173] [INSPIRE].
- [41] Y. Kiyo, A. Maier, P. Maierhöfer and P. Marquard, *Reconstruction of heavy quark current correlators at $O(\alpha_s^3)$* , *Nucl. Phys. B* **823** (2009) 269 [arXiv:0907.2120] [INSPIRE].
- [42] D. Greynat and S. Peris, *Resummation of threshold, low- and high-energy expansions for heavy-quark correlators*, *Phys. Rev. D* **82** (2010) 034030 [Erratum *ibid.* **82** (2010) 119907] [arXiv:1006.0643] [INSPIRE].
- [43] D. Greynat, P. Masjuan and S. Peris, *Analytic reconstruction of heavy-quark two-point functions at $O(\alpha_s^3)$* , *Phys. Rev. D* **85** (2012) 054008 [arXiv:1104.3425] [INSPIRE].
- [44] F.V. Tkachov, *A theorem on analytical calculability of four loop renormalization group functions*, *Phys. Lett. B* **100** (1981) 65 [INSPIRE].
- [45] K.G. Chetyrkin and F.V. Tkachov, *Integration by parts: the algorithm to calculate β -functions in 4 loops*, *Nucl. Phys. B* **192** (1981) 159 [INSPIRE].
- [46] S. Laporta, *High precision calculation of multiloop Feynman integrals by difference equations*, *Int. J. Mod. Phys. A* **15** (2000) 5087 [hep-ph/0102033] [INSPIRE].
- [47] A. Maier and P. Marquard, *Validity of Padé approximations in vacuum polarization at three- and four-loop order*, *Phys. Rev. D* **97** (2018) 056016 [arXiv:1710.03724] [INSPIRE].
- [48] K.G. Chetyrkin, A.L. Kataev and F.V. Tkachov, *Higher order corrections to $\sigma(e^+e^- \rightarrow \text{hadrons})$ in quantum chromodynamics*, *Phys. Lett. B* **85** (1979) 277 [INSPIRE].

- [49] M. Dine and J.R. Sapirstein, *Higher order QCD corrections in e^+e^- annihilation*, *Phys. Rev. Lett.* **43** (1979) 668 [INSPIRE].
- [50] W. Celmaster and R.J. Gonsalves, *QCD perturbation expansions in a coupling constant renormalized by momentum space subtraction*, *Phys. Rev. Lett.* **42** (1979) 1435 [INSPIRE].
- [51] S.G. Gorishnii, A.L. Kataev and S.A. Larin, *Three loop corrections of order $O(M^2)$ to the correlator of electromagnetic quark currents*, *Nuovo Cim. A* **92** (1986) 119 [INSPIRE].
- [52] K.G. Chetyrkin, R. Harlander, J.H. Kühn and M. Steinhauser, *Mass corrections to the vector current correlator*, *Nucl. Phys. B* **503** (1997) 339 [hep-ph/9704222] [INSPIRE].
- [53] R. Harlander and M. Steinhauser, *$O(\alpha_s^2)$ corrections to top quark production at e^+e^- colliders*, *Eur. Phys. J. C* **2** (1998) 151 [hep-ph/9710413] [INSPIRE].
- [54] R. Harlander and M. Steinhauser, *Higgs decay to top quarks at $O(\alpha_s^2)$* , *Phys. Rev. D* **56** (1997) 3980 [hep-ph/9704436] [INSPIRE].
- [55] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, *Vacuum polarization in pQCD: first complete $O(\alpha_s^4)$ result*, *Nucl. Phys. B Proc. Suppl.* **135** (2004) 243 [INSPIRE].
- [56] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, *$R(s)$ and hadronic τ -decays in order α_s^4 : technical aspects*, *Nucl. Phys. B Proc. Suppl.* **189** (2009) 49 [arXiv:0906.2987] [INSPIRE].
- [57] K.G. Chetyrkin and A. Khodjamirian, *Strange quark mass from pseudoscalar sum rule with $O(\alpha_s^4)$ accuracy*, *Eur. Phys. J. C* **46** (2006) 721 [hep-ph/0512295] [INSPIRE].
- [58] P.A. Baikov, K.G. Chetyrkin and J.H. Kühn, *Scalar correlator at $O(\alpha_s^4)$, Higgs decay into b-quarks and bounds on the light quark masses*, *Phys. Rev. Lett.* **96** (2006) 012003 [hep-ph/0511063] [INSPIRE].
- [59] P.A. Baikov, K.G. Chetyrkin, J.H. Kühn and J. Rittinger, *Complete $O(\alpha_s^4)$ QCD corrections to hadronic Z-decays*, *Phys. Rev. Lett.* **108** (2012) 222003 [arXiv:1201.5804] [INSPIRE].
- [60] P.A. Baikov, K.G. Chetyrkin, J.H. Kühn and J. Rittinger, *Adler function, sum rules and Crewther relation of order $O(\alpha_s^4)$: the singlet case*, *Phys. Lett. B* **714** (2012) 62 [arXiv:1206.1288] [INSPIRE].
- [61] K.G. Chetyrkin, V.P. Spiridonov and S.G. Gorishnii, *Wilson expansion for correlators of vector currents at the two loop level: dimension four operators*, *Phys. Lett. B* **160** (1985) 149 [INSPIRE].
- [62] K.G. Chetyrkin and J.H. Kühn, *Quartic mass corrections to R_{had}* , *Nucl. Phys. B* **432** (1994) 337 [hep-ph/9406299] [INSPIRE].
- [63] K.G. Chetyrkin, R.V. Harlander and J.H. Kühn, *Quartic mass corrections to R_{had} at $O(\alpha_s^3)$* , *Nucl. Phys. B* **586** (2000) 56 [Erratum *ibid.* **634** (2002) 413] [hep-ph/0005139] [INSPIRE].
- [64] R. Harlander, *Quarkmasseneffekte in der Quantenchromodynamik und asymptotische Entwicklung von Feynman-Integralen* (in German), Shaker Verlag, Germany (1998).
- [65] J. Artz, R.V. Harlander, F. Lange, T. Neumann and M. Prausa, *Results and techniques for higher order calculations within the gradient-flow formalism*, *JHEP* **06** (2019) 121 [Erratum *ibid.* **10** (2019) 032] [arXiv:1905.00882] [INSPIRE].
- [66] S.G. Gorishnii, S.A. Larin and F.V. Tkachov, *The algorithm for OPE coefficient functions in the \overline{MS} scheme*, *Phys. Lett. B* **124** (1983) 217 [INSPIRE].
- [67] S.G. Gorishnii and S.A. Larin, *Coefficient functions of asymptotic operator expansions in minimal subtraction scheme*, *Nucl. Phys. B* **283** (1987) 452 [INSPIRE].

- [68] R.V. Harlander and T. Neumann, *The perturbative QCD gradient flow to three loops*, *JHEP* **06** (2016) 161 [[arXiv:1606.03756](#)] [[INSPIRE](#)].
- [69] P. Nogueira, *Automatic Feynman graph generation*, *J. Comput. Phys.* **105** (1993) 279 [[INSPIRE](#)].
- [70] P. Nogueira, *Abusing QGRAF*, *Nucl. Instrum. Meth. A* **559** (2006) 220 [[INSPIRE](#)].
- [71] R. Harlander, T. Seidensticker and M. Steinhauser, *Complete corrections of $O(\alpha_s)$ to the decay of the Z boson into bottom quarks*, *Phys. Lett. B* **426** (1998) 125 [[hep-ph/9712228](#)] [[INSPIRE](#)].
- [72] T. Seidensticker, *Automatic application of successive asymptotic expansions of Feynman diagrams*, in *6th International Workshop on New Computing Techniques in Physics Research: software engineering, artificial intelligence neural nets, genetic algorithms, symbolic algebra, automatic calculation*, (1999) [[hep-ph/9905298](#)] [[INSPIRE](#)].
- [73] J.A.M. Vermaseren, *New features of FORM*, [math-ph/0010025](#) [[INSPIRE](#)].
- [74] J. Kuipers, T. Ueda, J.A.M. Vermaseren and J. Vollinga, *FORM version 4.0*, *Comput. Phys. Commun.* **184** (2013) 1453 [[arXiv:1203.6543](#)] [[INSPIRE](#)].
- [75] T. van Ritbergen, A.N. Schellekens and J.A.M. Vermaseren, *Group theory factors for Feynman diagrams*, *Int. J. Mod. Phys. A* **14** (1999) 41 [[hep-ph/9802376](#)] [[INSPIRE](#)].
- [76] P. Maierhöfer, J. Usovitsch and P. Uwer, *Kira — a Feynman integral reduction program*, *Comput. Phys. Commun.* **230** (2018) 99 [[arXiv:1705.05610](#)] [[INSPIRE](#)].
- [77] P. Maierhöfer and J. Usovitsch, *Kira 1.2 release notes*, [arXiv:1812.01491](#) [[INSPIRE](#)].
- [78] J. Klappert, F. Lange, P. Maierhöfer and J. Usovitsch, *Integral reduction with Kira 2.0 and finite field methods*, [arXiv:2008.06494](#) [[INSPIRE](#)].
- [79] J. Klappert and F. Lange, *Reconstructing rational functions with FireFly*, *Comput. Phys. Commun.* **247** (2020) 106951 [[arXiv:1904.00009](#)] [[INSPIRE](#)].
- [80] J. Klappert, S.Y. Klein and F. Lange, *Interpolation of dense and sparse rational functions and other improvements in FireFly*, [arXiv:2004.01463](#) [[INSPIRE](#)].
- [81] M. Kauers, *Fast solvers for dense linear systems*, *Nucl. Phys. B Proc. Suppl.* **183** (2008) 245 [[INSPIRE](#)].
- [82] P. Kant, *Finding linear dependencies in integration-by-parts equations: a Monte Carlo approach*, *Comput. Phys. Commun.* **185** (2014) 1473 [[arXiv:1309.7287](#)] [[INSPIRE](#)].
- [83] A. von Manteuffel and R.M. Schabinger, *A novel approach to integration by parts reduction*, *Phys. Lett. B* **744** (2015) 101 [[arXiv:1406.4513](#)] [[INSPIRE](#)].
- [84] T. Peraro, *Scattering amplitudes over finite fields and multivariate functional reconstruction*, *JHEP* **12** (2016) 030 [[arXiv:1608.01902](#)] [[INSPIRE](#)].
- [85] T. Binoth and G. Heinrich, *Numerical evaluation of multiloop integrals by sector decomposition*, *Nucl. Phys. B* **680** (2004) 375 [[hep-ph/0305234](#)] [[INSPIRE](#)].
- [86] A.V. Smirnov, *FIESTA 3: cluster-parallelizable multiloop numerical calculations in physical regions*, *Comput. Phys. Commun.* **185** (2014) 2090 [[arXiv:1312.3186](#)] [[INSPIRE](#)].
- [87] A.V. Smirnov, *FIESTA 4: optimized Feynman integral calculations with GPU support*, *Comput. Phys. Commun.* **204** (2016) 189 [[arXiv:1511.03614](#)] [[INSPIRE](#)].

- [88] A.C. Genz and A.A. Malik, *An imbedded family of fully symmetric numerical integration rules*, *SIAM J. Numer. Anal.* **20** (1983) 580.
- [89] L. Fousse, G. Hanrot, V. Lefèvre, P. Pélicier and P. Zimmermann, *MPFR: a multiple-precision binary floating-point library with correct rounding*, *ACM Trans. Math. Softw.* **33** (2007) 13.
- [90] Wolfram Research Inc., *Mathematica*, version 12, <https://www.wolfram.com/mathematica>.
- [91] A. Carosso, A. Hasenfratz and E.T. Neil, *Renormalization group properties of scalar field theories using gradient flow*, *PoS(LATTICE2018)248* (2018) [[arXiv:1811.03182](https://arxiv.org/abs/1811.03182)] [[INSPIRE](https://inspirehep.net/literature/1811031)].
- [92] Y. Abe and M. Fukuma, *Gradient flow and the renormalization group*, *PTEP* **2018** (2018) 083B02 [[arXiv:1805.12094](https://arxiv.org/abs/1805.12094)] [[INSPIRE](https://inspirehep.net/literature/1805120)].
- [93] A. Carosso, *Stochastic renormalization group and gradient flow*, *JHEP* **01** (2020) 172 [[arXiv:1904.13057](https://arxiv.org/abs/1904.13057)] [[INSPIRE](https://inspirehep.net/literature/1904130)].
- [94] H. Sonoda and H. Suzuki, *Derivation of a gradient flow from the exact renormalization group*, *PTEP* **2019** (2019) 033B05 [[arXiv:1901.05169](https://arxiv.org/abs/1901.05169)] [[INSPIRE](https://inspirehep.net/literature/1901051)].
- [95] K. Chetyrkin and P. Baikov, *QCD vacuum energy in 5 loops*, *PoS(RADCOR2017)025* (2018).