



Matrices Over Nondivision Algebras Without Eigenvalues

Drahoslava Janovská and Gerhard Opfer*

Abstract. We are concerned with matrices over nondivision algebras and show by an example from an \mathbb{R}^4 algebra that these matrices do not necessarily have eigenvalues, even if these matrices are invertible. The standard condition for eigenvectors $\mathbf{x} \neq \mathbf{0}$ will be replaced by the condition that \mathbf{x} contains at least one invertible component which is the same as $\mathbf{x} \neq \mathbf{0}$ for division algebras. The topic is of principal interest, and leads to the question what qualifies a matrix over a nondivision algebra to have eigenvalues. And connected with this problem is the question, whether these matrices are diagonalizable or triangulizable and allow a Schur decomposition. There is a last section where the question whether a specific matrix \mathbf{A} has eigenvalues is extended to all eight \mathbb{R}^4 algebras by applying numerical means. As a curiosity we found that the considered matrix \mathbf{A} over the algebra of *tessarines*, which is a commutative algebra, introduced by Cockle (Phil Mag 35(3):434–437, 1849; <http://www.oocities.org/cocklebio/>), possesses eigenvalues.

Mathematics Subject Classification. 11E88, 15A66, 20G20, 65F15, 65H17.

Keywords. Eigenvalues of matrices over nondivision algebras, Eigenvalues of matrices over coquaternions, Split-quaternions, Eigenvalues of matrices over a general \mathbb{R}^4 algebra, Diagonalizable and triangulizable matrices over nondivision algebras.

1. Introduction

By *standard matrices* we understand square matrices with entries from \mathbb{R} , \mathbb{C} , \mathbb{H} , the fields of real numbers, complex numbers, and quaternions, respectively. They have in common that they form a division algebra. By algebras in general we will understand the vector space \mathbb{R}^N equipped with an associative multiplication $\mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}^N$, with a one, often abbreviated as 1, where $N \in \mathbb{N}$, the set of positive integers. For algebras in this general sense, we will

*Corresponding author.

use the notation \mathcal{A} . More details can be found in a book by Garling [5]. The name *geometric algebra* is often used for these algebras [6].

A *division algebra* is an algebra, where the zero element is the only noninvertible element. Square matrices which have n rows and n columns are called matrices of order n . This paper will deal with matrices of order $n \in \mathbb{N}$, where the entries are from some algebra in the sense mentioned. In this context we also speak of matrices over an algebra \mathcal{A} . We also write $\mathbf{A} \in \mathcal{A}^{n \times n}$ to denote a square matrix of order n with entries from an algebra \mathcal{A} . At some occasions we also have to deal with nonsquare matrices which are denoted by $\mathbf{A} \in \mathcal{A}^{m \times n}$. If $m = 1$, \mathbf{A} is called *row vector*, if $n = 1$, \mathbf{A} is called *column vector*.

The question whether standard matrices have eigenvalues does not come up, since it is known, that these matrices have always eigenvalues and the number of eigenvalues never exceeds the order n . See Horn and Johnson [7] for matrices over \mathbb{R}, \mathbb{C} , and Brenner [1], and Zhang [19] for matrices over \mathbb{H} . For matrices over \mathcal{A} the term *number of eigenvalues* will be made more precise a little later.

However, for matrices over nondivision algebras the answer to the question *are there eigenvalues at all* is not so clear, and in the first place one should make clear what an eigenvalue is. Let $\mathbf{A} \in \mathcal{A}^{n \times n}$ where \mathcal{A} is a division algebra. The standard definition is

$$\mathbf{A}\mathbf{x} = \mathbf{x}\lambda, \quad \mathbf{x} \neq \mathbf{0} \tag{1.1}$$

for an *eigenvalue* $\lambda \in \mathcal{A}$ with respect to an *eigenvector* $\mathbf{x} \in \mathcal{A}^{n \times 1}$. Even in the standard case it is reasonable, to write $\mathbf{x}\lambda$ instead of $\lambda\mathbf{x}$, because λ may be regarded as a 1×1 matrix. The vector $\mathbf{0}$ is the zero column vector. In a division algebra, the condition $\mathbf{x} \neq \mathbf{0}$ is equivalent to saying that the vector \mathbf{x} contains at least one invertible component. However, if we transfer this definition also to nondivision algebras, then $\mathbf{x} \neq \mathbf{0}$ would allow that all components of \mathbf{x} are noninvertible. We will see a little later, that this definition does not lead to a reasonable theory of eigenvalues.

Definition 1.1. Let \mathcal{A} be a nondivision algebra. We denote the set of noninvertible elements of \mathcal{A} by $\mathcal{N}_{\mathcal{A}}$. An element $\lambda \in \mathcal{A}$ will be called an *eigenvalue with respect to an eigenvector \mathbf{x} of a matrix $\mathbf{A} \in \mathcal{A}^{n \times n}$* if

$$\mathbf{A}\mathbf{x} = \mathbf{x}\lambda, \quad \mathbf{x} \notin (\mathcal{N}_{\mathcal{A}})^{n \times 1}. \tag{1.2}$$

The set of all eigenvalues of a matrix \mathbf{A} will be denoted by $\sigma(\mathbf{A})$. If (λ, \mathbf{x}) only solves (1.1), then, λ will be called a *candidate for an eigenvalue with respect to \mathbf{x}* .

Definition 1.2. Let \mathcal{A} be an algebra. Two elements $a, b \in \mathcal{A}$ are called *similar*, denoted by $a \sim b$, if there is an invertible element $h \in \mathcal{A}$ such that $a = h^{-1}bh$. The set of all elements similar to a is denoted by $[a]$ and

$$[a] := \{b = h^{-1}ah \text{ for all } h \notin \mathcal{N}_{\mathcal{A}}\}$$

is called the *similarity class* of a . See also [18] for an early paper in this context. We will see in the next theorem, that the existence of one eigenvalue λ will imply that the whole similarity class $[\lambda]$ consists of eigenvalues.

Theorem 1.3. *Let λ be an eigenvalue of \mathbf{A} with respect to the eigenvector \mathbf{x} . Then, $h^{-1}\lambda h$ is also an eigenvalue of \mathbf{A} with respect to $\mathbf{x}h$ for all $h \notin \mathcal{N}_{\mathcal{A}}$.*

Proof. Multiply the defining Eq. (1.2) from the right by h . Then,

$$\mathbf{A}(\mathbf{x}h) = \mathbf{x}\lambda h = (\mathbf{x}h)(h^{-1}\lambda h).$$

Since $\mathbf{x} \notin (\mathcal{N}_{\mathcal{A}})^{n \times 1}$, the same is true for $\mathbf{x}h$, since h is invertible. □

In view of Theorem 1.3 the *number of eigenvalues* of a matrix \mathbf{A} is defined as the number of similarity classes which contain eigenvalues. We see in the next theorem that one of the essential requirements for eigenvalues is valid in the framework of this definition of eigenvalues.

Theorem 1.4. *Let \mathbf{A} be a matrix over an arbitrary algebra \mathcal{A} and let there be two eigenvalues λ_1 and λ_2 of \mathbf{A} with respect to the same eigenvector \mathbf{x} . Then, $\lambda_1 = \lambda_2$.*

Proof. By definition, $\mathbf{A}\mathbf{x} = \mathbf{x}\lambda_j$, $j = 1, 2$. This implies $\mathbf{0} = \mathbf{x}(\lambda_1 - \lambda_2)$. Since, also by definition, \mathbf{x} contains an invertible component, we have $\lambda_1 - \lambda_2 = 0$. □

We also see here already that the assumption $\mathbf{x} \neq \mathbf{0}$ alone would not allow the conclusion of the last theorem.

Let us turn to the question, whether all matrices over nondivision algebras have eigenvalues. There are two strategies. If one believes that there are always eigenvalues, one has to prove it. Otherwise, life is a little easier, one has to find one example of a matrix which does not have eigenvalues. We will pursue this line.

2. Some Elementary Properties of \mathbb{R}^4 Algebras

In order to find an example we will restrict our search to the eight algebras of \mathbb{R}^4 . These algebras with their multiplication rules are summarized in Table 1. The four units in these algebras are denoted by $1, \mathbf{i}, \mathbf{j}, \mathbf{k}$.

The full multiplication table of the eight listed algebras can be obtained by multiplying the last three columns in Table 1 from the right by $\mathbf{j}, \mathbf{k}, \mathbf{i}$, respectively. We note that the four algebras $\mathbb{H}_{\text{tes}}, \mathbb{H}_{\text{cotes}}, \mathbb{H}_{\text{tan}}, \mathbb{H}_{\text{cotan}}$ with the numbers 3, 4, 7, 8 are commutative. See [8, Table 12.1].

Elements in all these algebras will be denoted by $a = (a_1, a_2, a_3, a_4)$ which is the same as $a = a_1 + a_2\mathbf{i} + a_3\mathbf{j} + a_4\mathbf{k}$, where $a_1, a_2, a_3, a_4 \in \mathbb{R}$. The first component a_1 of a is called *real part* of a and it is denoted by $\Re(a)$. An algebra element of the form $(a_1, 0, 0, 0)$, $a_1 \in \mathbb{R}$ is called *real* and the set of real elements is also identified with \mathbb{R} . The set \mathbb{R} plays a special role in all noncommutative \mathbb{R}^4 algebras. All elements of \mathbb{R} (and no others) commute with all algebra elements. Let $a, b \in \mathcal{A}$ where \mathcal{A} is one of the eight \mathbb{R}^4 algebras. Then,

$$\Re(ab) = \Re(ba).$$

For the four noncommutative algebras (No 1,2,5,6) we define the *conjugate of an element* $a = (a_1, a_2, a_3, a_4)$ in the notation \bar{a} or $\text{conj}(a)$ by

TABLE 1. Eight \mathbb{R}^4 algebras with their essential multiplication rules and names¹

No	Name of algebra	Short name	i^2	j^2	k^2	ij	jk	ki
1	Quaternions	\mathbb{H}	-1	-1	-1	\mathbf{k}	\mathbf{i}	\mathbf{j}
2	Coquaternions	\mathbb{H}_{coq}	-1	1	1	\mathbf{k}	$-\mathbf{i}$	\mathbf{j}
3	Tessarines	\mathbb{H}_{tes}	-1	1	-1	\mathbf{k}	\mathbf{i}	$-\mathbf{j}$
4	Cotessarines	$\mathbb{H}_{\text{cotes}}$	1	1	1	\mathbf{k}	\mathbf{i}	\mathbf{j}
5	Nectarines	\mathbb{H}_{nec}	1	-1	1	\mathbf{k}	\mathbf{i}	$-\mathbf{j}$
6	Conectarines	\mathbb{H}_{con}	1	1	-1	\mathbf{k}	$-\mathbf{i}$	$-\mathbf{j}$
7	Tangerines	\mathbb{H}_{tan}	1	-1	-1	\mathbf{k}	$-\mathbf{i}$	\mathbf{j}
8	Cotangerines	$\mathbb{H}_{\text{cotan}}$	-1	-1	1	\mathbf{k}	$-\mathbf{i}$	$-\mathbf{j}$

$$\bar{a} = \text{conj}(a) := (a_1, -a_2, -a_3, -a_4),$$

and we use the notation

$$\text{abs}_2(a) := a\bar{a}$$

for the product $a\bar{a}$. The essential properties are given in the next theorem.

Theorem 2.1. *Let \mathcal{A} be one of the four noncommutative algebras of Table 1. Then $\text{abs}_2(a) = a\bar{a} = \bar{a}a$ is real, and a is invertible if and only if $\text{abs}_2(a) \neq 0$. For the inverse of a there is the formula*

$$a^{-1} = \frac{\bar{a}}{\text{abs}_2(a)}, \quad \text{if } \text{abs}_2(a) \neq 0.$$

Let $a, b \in \mathcal{A}$. Then, there is the property that abs_2 is multiplicative:

$$\text{abs}_2(ab) = \text{abs}_2(ba) = \text{abs}_2(a)\text{abs}_2(b).$$

Two nonreal elements $a, b \in \mathcal{A}$ are similar if and only if

$$\Re(a) = \Re(b) \text{ and } \text{abs}_2(a) = \text{abs}_2(b). \tag{2.1}$$

For $\text{abs}_2(a)$ there is the formula

$$\text{abs}_2(a) = \begin{cases} a_1^2 + a_2^2 + a_3^2 + a_4^4 & \text{for } a \in \mathbb{H}, \\ a_1^2 + a_2^2 - a_3^2 - a_4^4 & \text{for } a \in \mathbb{H}_{\text{coq}}, \\ a_1^2 - a_2^2 + a_3^2 - a_4^4 & \text{for } a \in \mathbb{H}_{\text{nec}}, \\ a_1^2 - a_2^2 - a_3^2 + a_4^4 & \text{for } a \in \mathbb{H}_{\text{con}}. \end{cases} \tag{2.2}$$

Proof. See [8]. □

It should be noted, that the additional condition in (2.1) that a, b are nonreal, cannot be omitted for all three noncommutative \mathbb{R}^4 algebras different from quaternions \mathbb{H} . Otherwise $a = (1, 0, 0, 0)$ and $b = (1, 5, 4, 3)$ would be similar in \mathbb{H}_{coq} , $a = (1, 0, 0, 0)$ and $b = (1, 4, 5, 3)$ would be similar in \mathbb{H}_{nec} and $a = (1, 0, 0, 0)$ and $b = (1, 3, 4, 5)$ would be similar in \mathbb{H}_{con} . However, in all four noncommutative \mathbb{R}^4 algebras a real element is never similar to a nonreal element. In Pogoruy and Rodríguez-Dagnino [15], this condition is missing.

¹The names of the algebras with the numbers 5–8 where given by Schmeikal [16].

TABLE 2. Multiplication table of $\mathbf{i}a, \mathbf{j}a, \mathbf{k}a, a = (1, 2, 3, 4)$

\mathcal{A}	$\mathbf{i}a$	$\mathbf{j}a$	$\mathbf{k}a$
\mathbb{H}	$(-2, 1, -4, 3)$	$(-3, 4, 1, -2)$	$(-4, -3, 2, 1)$
\mathbb{H}_{coq}	$(-2, 1, -4, 3)$	$(3, -4, 1, -2)$	$(4, 3, 2, 1)$
\mathbb{H}_{tes}	$(-2, 1, -4, 3)$	$(3, 4, 1, 2)$	$(-4, 3, -2, 1)$
$\mathbb{H}_{\text{cotes}}$	$(2, 1, 4, 3)$	$(3, 4, 1, 2)$	$(4, 3, 2, 1)$
\mathbb{H}_{nec}	$(2, 1, 4, 3)$	$(-3, 4, 1, -2)$	$(4, -3, -2, 1)$
\mathbb{H}_{con}	$(2, 1, 4, 3)$	$(3, -4, 1, -2)$	$(-4, 3, -2, 1)$
\mathbb{H}_{tan}	$(2, 1, 4, 3)$	$(-3, -4, 1, 2)$	$(-4, -3, 2, 1)$
$\mathbb{H}_{\text{cotan}}$	$(-2, 1, -4, 3)$	$(-3, -4, 1, 2)$	$(4, -3, -2, 1)$

TABLE 3. Multiplication table of $a\mathbf{i}, a\mathbf{j}, a\mathbf{k}, a = (1, 2, 3, 4)$

\mathcal{A}	$a\mathbf{i}$	$a\mathbf{j}$	$a\mathbf{k}$
\mathbb{H}	$(-2, 1, 4, -3)$	$(-3, -4, 1, 2)$	$(-4, 3, -2, 1)$
\mathbb{H}_{coq}	$(-2, 1, 4, -3)$	$(3, 4, 1, 2)$	$(4, -3, -2, 1)$
\mathbb{H}_{tes}	$(-2, 1, -4, 3)$	$(3, 4, 1, 2)$	$(-4, 3, -2, 1)$
$\mathbb{H}_{\text{cotes}}$	$(2, 1, 4, 3)$	$(3, 4, 1, 2)$	$(4, 3, 2, 1)$
\mathbb{H}_{nec}	$(2, 1, -4, -3)$	$(-3, -4, 1, 2)$	$(4, 3, 2, 1)$
\mathbb{H}_{con}	$(2, 1, -4, -3)$	$(3, 4, 1, 2)$	$(-4, -3, 2, 1)$
\mathbb{H}_{tan}	$(2, 1, 4, 3)$	$(-3, -4, 1, 2)$	$(-4, -3, 2, 1)$
$\mathbb{H}_{\text{cotan}}$	$(-2, 1, -4, 3)$	$(-3, -4, 1, 2)$	$(4, -3, -2, 1)$

Since eigenvalues appear always in similarity classes, we can reduce the eigenvalues to certain representatives of the similarity classes.

Lemma 2.2. *All eigenvalues of matrices over \mathbb{H} can be given the form $a + b\mathbf{i}$. All eigenvalues of matrices over \mathbb{H}_{coq} and over \mathbb{H}_{nec} can be given the form $a + b\mathbf{i}$ or $a + b\mathbf{j}$. All eigenvalues of matrices over \mathbb{H}_{con} can be given the form $a + b\mathbf{i}$ or $a + b\mathbf{k}$, where in all cases $b \geq 0$.*

Proof. This follows from (2.2) in dependence of the sign of abs_2 together with (2.1). □

For this reason all eigenvalues of matrices over \mathbb{H} can be represented as complex numbers, see Brenner [1] and therefore, the eigenvalue theory of matrices over \mathbb{H} is very similar to the standard case of matrices over \mathbb{R} or over \mathbb{C} .

For later use we will provide two tables for simple multiplication rules.

Lemma 2.3. *Let $a \in \mathcal{A}$ where \mathcal{A} is one of the eight \mathbb{R}^4 algebras of Table 1. Then, the multiplication of a by $\mathbf{i}, \mathbf{j}, \mathbf{k}$ from the left and from the right results apart from the signs in a permutation of the four components of a . Let $a = (1, 2, 3, 4)$, then the corresponding permutations (inclusive the signs) are given in Tables 2 and 3.*

Proof. Apply the multiplication rules. □

3. One Dimensional Eigenvalue Problems and Matrix Representations

All algebras (at least those considered here) are isomorphic to certain real matrix spaces. So it is reasonable in an eigenvalue problem over an algebra \mathcal{A} to replace the algebra elements by isomorphic matrices so that the eigenvalue problem over \mathcal{A} appears as a problem in real matrices.² Since these matrices can have only real or complex eigenvalues, it is of interest to study the connection between these eigenvalues and the eigenvalues of the underlying matrices over \mathcal{A} . For this purpose we study the simplest, one dimensional case

$$ax = x\lambda, \quad a, x, \lambda \in \mathcal{A}, \quad x \text{ invertible.} \quad (3.1)$$

A (1×1) matrix a may be regarded simultaneously as a diagonal and a triangular matrix. Therefore, the results of the Lemma 4.7, p. 606 are valid for the problem (3.1).

Lemma 3.1. *Let \mathcal{A} be an arbitrary algebra. The eigenvalue problem (3.1) is always solvable and the solution is*

$$\lambda \in [a]. \quad (3.2)$$

Proof. The invertible eigenvector $x = 1$ solves the problem. Multiplying Eq. (3.1) from the left by x^{-1} , yields $\lambda = x^{-1}ax = a$. The remaining part follows from Theorem 1.3. \square

How to find the matrix equivalents of an algebra element $a \in \mathcal{A}$. The simplest idea is, to regard the mapping

$$\ell_a : \mathcal{A} \rightarrow \mathcal{A} \text{ defined by } \ell_a(x) := ax, \quad x \in \mathcal{A} \quad (3.3)$$

as a linear mapping over \mathbb{R} . Let \mathcal{A} have dimension N and $a = (a_1, a_2, \dots, a_N) \in \mathcal{A}$. Define

$$\text{col}(a) := (a_1, a_2, \dots, a_N)^T \in \mathbb{R}^N,$$

where T defines *transposition*, such that $\text{col}(a)$ is a real column vector of length N . Then according to a well known theory (see Horn and Johnson [7, Chapter 0]) all linear transformations over \mathbb{R} can be represented by a real $N \times N$ matrix \mathbf{M} such that the linearity of ℓ_a defined in (3.3) implies that

$$\text{col}(\ell_a(x)) = \mathbf{M}\text{col}(x). \quad (3.4)$$

Details for computing the matrix \mathbf{M} are in [8, 9, 12]. Now, we compare the (known) real or complex eigenvalues of \mathbf{M} with the corresponding representatives of $[a]$. For all \mathbb{R}^4 algebras the eight 4×4 matrix representations are given in [8, Table 12.2]. The four matrices belonging to the noncommutative cases have a minimal polynomial of degree only 2, such that there are (at most) two distinct eigenvalues. The eigenvalues of all eight matrices are listed in Table 4.

² For some algebras, also complex matrix representations are possible. For quaternions see [17, p. 55]. Here, we will restrict ourselves to real matrix representations.

In the following Theorem we give a reconstruction scheme for representatives of eigenvalues in $[a]$, from the real and complex eigenvalues λ_ℓ , $\ell = 1, 2, 3, 4$ of \mathbf{M} (see Table 4), where a is a member of one of the eight \mathbb{R}^4 algebras.

Theorem 3.2. *Let $a = (a_1, a_2, a_3, a_4) \in \mathcal{A}$ where \mathcal{A} is one of the eight \mathbb{R}^4 algebras. Let the real or complex eigenvalues of \mathbf{M} be λ_ℓ , $\ell = 1, 2, 3, 4$, which are listed in Table 4. A representative of $[a]$ is given by the entries of Table 5.*

Proof. For the noncommutative cases we use Lemma 2.2 by keeping the similarity class of $[a]$ with the help of formula (2.1). For the commutative cases we reconstruct the four values of a_j from the four eigenvalues λ_j , $j = 1, 2, 3, 4$. □

The cases $A < 0$ in the first part of Table 5 always mean that the eigenvalues λ_j , $j = 1, 2$ are real. We will provide two simple examples, one for the noncommutative, one for the commutative case, which show, that allowing eigenvectors with all components noninvertible may lead to not uniquely determined eigenvalues and to other strange situations.

Example 3.3. Let $\mathcal{A} = \mathbb{H}_{\text{coq}}$. Define

$$\mathbf{A} = (1, 3, 2, 3) \in \mathbb{H}_{\text{coq}}^{1 \times 1} \text{ and } \mathbf{x} = (1, 3, -1, 3) \in \mathbb{H}_{\text{coq}}^{1 \times 1}.$$

Then, $\mathbf{Ax} = (-1, -3, 1, -3)$. Note, that \mathbf{x} is noninvertible. The equation $\mathbf{Ax} = \mathbf{x}\lambda$ has the following solutions for λ : choose λ_1, λ_2 arbitrarily and determine λ_3, λ_4 by

$$\lambda_3 = \frac{-4 - 4\lambda_1 - 3\lambda_2}{5}, \quad \lambda_4 = \frac{-3 - 3\lambda_1 + 4\lambda_2}{5}.$$

Examples for λ are $\lambda = (-4, 4, 0, 5)$, $\lambda = \frac{1}{10}(-5, 0, -4, -3)$. The first one is invertible, the second one is noninvertible. This example shows, that there are infinitely many similarity classes which contain eigenvalues and

TABLE 4. List of all eigenvalues $\lambda_{1,2,3,4}$ of \mathbf{M} in all eight \mathbb{R}^4 algebras. The eigenvalues $\lambda_{1,3}$ correspond to the +sign, and the eigenvalues $\lambda_{2,4}$ to the -sign. The abbreviation $A := \text{abs}_2(a) - a_1^2$ with $\text{abs}_2(a)$ from (2.2) is used

No	Name of algebra	$\lambda_{1,2}$	$\lambda_{3,4}$
1	Quaternions	$a_1 \pm \sqrt{A} \mathbf{i}$	$a_1 \pm \sqrt{A} \mathbf{i}$
2	Coquaternions	$a_1 \pm \sqrt{A} \mathbf{i}$	$a_1 \pm \sqrt{A} \mathbf{i}$
3	Tessarines	$a_1 + a_3 \pm (a_2 + a_4) \mathbf{i}$	$a_1 - a_3 \pm (a_2 - a_4) \mathbf{i}$
4	Cotessarines	$a_1 - a_2 \pm (a_3 - a_4)$	$a_1 + a_2 \pm (a_3 + a_4)$
5	Nectarines	$a_1 \pm \sqrt{A} \mathbf{i}$	$a_1 \pm \sqrt{A} \mathbf{i}$
6	Conectarines	$a_1 \pm \sqrt{A} \mathbf{i}$	$a_1 \pm \sqrt{A} \mathbf{i}$
7	Tangerines	$a_1 + a_2 \pm (a_3 + a_4) \mathbf{i}$	$a_1 - a_2 \pm (a_3 - a_4) \mathbf{i}$
8	Cotangerines	$a_1 + a_4 \pm (a_2 - a_3) \mathbf{i}$	$a_1 - a_4 \pm (a_2 + a_3) \mathbf{i}$

TABLE 5. List of representatives of $[a]$ in terms of matrix eigenvalues $\lambda_j, j = 1, 2, 3, 4$ for all eight \mathbb{R}^4 algebras, $A := \text{abs}_2(a) - a_1^2$

Algebra	Representatives of $[a]$	Case
\mathbb{H}	$a_1 = \Re(\lambda_1), \quad a_2 = \Im(\lambda_1), \quad a_3 = a_4 = 0$	$A \geq 0$
\mathbb{H}_{coq}	$a_1 = \Re(\lambda_1), \quad a_2 = \Im(\lambda_1), \quad a_3 = a_4 = 0$	$A \geq 0$
\mathbb{H}_{coq}	$a_1 = (\lambda_1 + \lambda_2)/2, \quad a_3 = \text{abs}((\lambda_1 - \lambda_2)/2), \quad a_2 = a_4 = 0$	$A < 0$
\mathbb{H}_{nec}	$a_1 = \Re(\lambda_1), \quad a_3 = \Im(\lambda_1), \quad a_2 = a_4 = 0$	$A \geq 0$
\mathbb{H}_{nec}	$a_1 = (\lambda_1 + \lambda_2)/2, \quad a_2 = \text{abs}((\lambda_1 - \lambda_2)/2), \quad a_3 = a_4 = 0$	$A < 0$
\mathbb{H}_{con}	$a_1 = \Re(\lambda_1), \quad a_4 = \Im(\lambda_1), \quad a_2 = a_3 = 0$	$A \geq 0$
\mathbb{H}_{con}	$a_1 = (\lambda_1 + \lambda_2)/2, \quad a_2 = \text{abs}((\lambda_1 - \lambda_2)/2), \quad a_3 = a_4 = 0$	$A < 0$
Commutative algebras		
\mathbb{H}_{tes}	$a_1 = \frac{1}{4}(\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4), \quad a_2 = -\frac{i}{4}(\lambda_1 - \lambda_2 + \lambda_3 - \lambda_4)$ $a_3 = \frac{1}{4}(\lambda_1 - \lambda_2 - \lambda_3 - \lambda_4), \quad a_4 = -\frac{i}{4}(\lambda_1 - \lambda_2 - \lambda_3 + \lambda_4)$	
$\mathbb{H}_{\text{cotes}}$	$a_1 = \frac{1}{4}(\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4), \quad a_2 = \frac{1}{4}(-\lambda_1 - \lambda_2 + \lambda_3 + \lambda_4)$ $a_3 = \frac{1}{4}(\lambda_1 - \lambda_2 + \lambda_3 - \lambda_4), \quad a_4 = \frac{1}{4}(-\lambda_1 + \lambda_2 + \lambda_3 - \lambda_4)$	
\mathbb{H}_{tan}	$a_1 = \frac{1}{4}(\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4), \quad a_2 = \frac{i}{4}(\lambda_1 + \lambda_2 - \lambda_3 - \lambda_4)$ $a_3 = \frac{1}{4}(\lambda_1 - \lambda_2 - \lambda_3 - \lambda_4), \quad a_4 = \frac{i}{4}(\lambda_1 - \lambda_2 - \lambda_3 + \lambda_4)$	
$\mathbb{H}_{\text{cotan}}$	$a_1 = \frac{1}{4}(\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4), \quad a_2 = -\frac{i}{4}(\lambda_1 - \lambda_2 + \lambda_3 - \lambda_4)$ $a_3 = \frac{i}{4}(\lambda_1 - \lambda_2 - \lambda_3 + \lambda_4), \quad a_4 = \frac{1}{4}(\lambda_1 + \lambda_2 - \lambda_3 - \lambda_4)$	

the eigenvalues may be invertible and not invertible. The inverse of \mathbf{A} is $\mathbf{A}^{-1} = \frac{1}{3}(-1, 3, 2, 3)$. If we would exclude noninvertible eigenvectors, the solutions of $\mathbf{Ax} = \mathbf{x}\lambda$ would be only $\lambda \sim \mathbf{A}$.

Example 3.4. Let $\mathcal{A} = \mathbb{H}_{\text{cotes}}$. Note, that $\mathbb{H}_{\text{cotes}}$ is commutative. Define

$$\mathbf{A} = (2, 3, 2, -1) \in \mathbb{H}_{\text{cotes}}^{1 \times 1} \text{ and } \mathbf{x} = (1, -1, 1, -1) \in \mathbb{H}_{\text{cotes}}^{1 \times 1}.$$

Then, $\mathbf{Ax} = (2, -2, 2, -2)$. The solutions of $\mathbf{Ax} = \mathbf{x}\lambda$ are all $\lambda \in \mathbb{H}_{\text{cotes}}$ with

$$\lambda_1 - \lambda_2 + \lambda_3 - \lambda_4 = 2.$$

The matrix \mathbf{A} is invertible, $\mathbf{A}^{-1} = \frac{1}{24}(4, 1, 4, -5)$, the eigenvector \mathbf{x} is not invertible and in the set of possible eigenvalues λ there are invertible and noninvertible eigenvalues. An invertible example is $\lambda = (3, -1, 4, 6)$, a noninvertible example is $\lambda = (0, -1, 0, -1)$. This shows also, that an invertible

matrix \mathbf{A} may have noninvertible eigenvalues. On the other hand, if noninvertible eigenvectors are excluded, then, the only eigenvalue of \mathbf{A} is $\lambda = \mathbf{A}$.

Now we can apply these results also to larger matrices over noncommutative algebras by replacing the algebra elements in the matrix by real matrices according to formula (3.4). And by distinguishing between real and complex, nonreal eigenvalues we can obtain information on the eigenvalues of the original matrix by applying the results of Table 5. However, there is a serious drawback. Since we are using standard techniques to find the eigenvalues of real matrices, the standard condition $\mathbf{x} \neq \mathbf{0}$ is invoked. Thus, whether the eigenvalues computed by standard techniques are also eigenvalues of the underlying algebraic matrix is a priori unknown. We will use one example to demonstrate the details. Let

$$\mathbf{A} := \begin{bmatrix} 1 & \mathbf{i} \\ \mathbf{j} & \mathbf{k} \end{bmatrix} \tag{3.5}$$

be a matrix in \mathbb{H}_{coq} that will play a central role in this paper. If we replace the four entries of \mathbf{A} by corresponding real 4×4 matrices we obtain a real 8×8 matrix which has the (double) eigenvalues $\pm\sqrt{2}$ and $1 \pm \mathbf{i}$. Now the second and third row of Table 5 tells us, that the eigenvalues of \mathbf{A} are (possibly)

$$\lambda_1 := 1 + \mathbf{i} \text{ and } \lambda_2 := \sqrt{2}\mathbf{j}. \tag{3.6}$$

If we have a look at the standard eigenvectors we see that they all have the form (a, b, b, a, c, d, d, c) . If we speculate that the first and second four elements are coquaternionic elements, then they are noninvertible. We will see in the next section, that the true eigenvectors belonging to \mathbf{A} are indeed all noninvertible.

4. A Counterexample

We will eventually show, that the matrix \mathbf{A} defined in (3.5) will have no eigenvalues in \mathbb{H}_{coq} . We note, that \mathbf{A} is invertible in \mathbb{H}_{coq} and

$$\mathbf{A}^{-1} := \frac{1}{2} \begin{bmatrix} 1 & \mathbf{j} \\ -\mathbf{i} & \mathbf{k} \end{bmatrix}. \tag{4.1}$$

We denote the eigenvectors of \mathbf{A} by

$$\mathbf{x} := \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} := \begin{bmatrix} (u_1, u_2, u_3, u_4) \\ (v_1, v_2, v_3, v_4) \end{bmatrix}, \tag{4.2}$$

and require for a $\lambda \in \mathbb{H}_{\text{coq}}$ to qualify as eigenvalue of \mathbf{A} with respect to an eigenvector \mathbf{x} that

$$\mathbf{A}\mathbf{x} = \mathbf{x}\lambda, \quad \mathbf{x} \text{ contains at least one invertible component.} \tag{4.3}$$

See the Definition 1.1. We will first show, that the two candidates for eigenvalues $\lambda_j, j = 1, 2$ obtained in (3.6) are not eigenvalues of \mathbf{A} . Then we will show that in general no eigenvalues exist.

Lemma 4.1. $\lambda = 1 + \mathbf{i}$ is not an eigenvalue of \mathbf{A} , where \mathbf{A} is defined in (3.5).

Proof. Let $\lambda := 1 + \mathbf{i}$. Then, we have to find the eigenvectors $\mathbf{x} := (x_1, x_2)^T$ (T means transposition) from the defining equations

$$\begin{aligned} x_1 + \mathbf{i}x_2 &= x_1(1 + \mathbf{i}), \\ \mathbf{j}x_1 + \mathbf{k}x_2 &= x_2(1 + \mathbf{i}). \end{aligned}$$

By using the Tables 2, 3, the first equation can be written as

$$(u_1, u_2, u_3, u_4) + (-v_2, v_1, -v_4, v_3) = (u_1, u_2, u_3, u_4) + (-u_2, u_1, u_4, -u_3),$$

which implies

$$u_1 = v_1, \quad u_2 = v_2, \quad u_3 = -v_3, \quad u_4 = -v_4,$$

and the second equation as

$$(u_3, -u_4, u_1, -u_2) + (v_4, v_3, v_2, v_1) = (v_1, v_2, v_3, v_4) + (-v_2, v_1, v_4, -v_3).$$

Comparing all four components and using the results from the first equation, we obtain eventually the pair of equations

$$v_1 + v_2 = v_3 + v_4, \quad v_1 - v_2 = -v_3 + v_4.$$

Adding and subtracting these equations yields

$$v_1 = v_4, \quad v_2 = v_3,$$

Thus, altogether with $v_1, v_2 \in \mathbb{R}, |v_1| + |v_2| > 0$, the eigenvector components are

$$x_1 = (v_1, v_2, -v_2, -v_1), \quad x_2 = (v_1, v_2, v_2, v_1),$$

and both components are not invertible. Thus, $1 + \mathbf{i}$ is not an eigenvalue of \mathbf{A} , however, it is a candidate for an eigenvalue. \square

Lemma 4.2. $\lambda = \sqrt{2}\mathbf{j}$ is not an eigenvalue of \mathbf{A} , where \mathbf{A} is defined in (3.5).

Proof. In this case $\mathbf{x} := (x_1, x_2)^T$ is wanted which solves

$$\begin{aligned} x_1 + \mathbf{i}x_2 &= \sqrt{2}x_1\mathbf{j}, \\ \mathbf{j}x_1 + \mathbf{k}x_2 &= \sqrt{2}x_2\mathbf{j}. \end{aligned}$$

The first and second equation can be written as (see Tables 2, 3)

$$\begin{aligned} (u_1, u_2, u_3, u_4) + (-v_2, v_1, -v_4, v_3) &= \sqrt{2}(u_3, u_4, u_1, u_2), \\ (u_3, -u_4, u_1, -u_2) + (v_4, v_3, v_2, v_1) &= \sqrt{2}(v_3, v_4, v_1, v_2), \end{aligned}$$

and these equations imply, respectively

$$\begin{aligned} v_1 &= \sqrt{2}u_4 - u_2, & v_2 &= u_1 - \sqrt{2}u_3, & v_3 &= \sqrt{2}u_2 - u_4, & v_4 &= u_3 - \sqrt{2}u_1, \\ u_1 &= \sqrt{2}v_1 - v_2, & u_2 &= v_1 - \sqrt{2}v_2, & u_3 &= \sqrt{2}v_3 - v_4, & u_4 &= v_3 - \sqrt{2}v_4. \end{aligned}$$

From the first set of equations, we deduce

$$v_1 + \sqrt{2}v_3 = u_2, \quad \sqrt{2}v_1 + v_3 = u_4, \quad \sqrt{2}v_2 + v_4 = -u_3, \quad v_2 + \sqrt{2}v_4 = -u_1.$$

Comparing with the second set of equations yields

$$v_1 = -v_4, \quad v_2 = -v_3, \quad u_3 = u_2, \quad u_1 = u_4.$$

This finally yields with $u_1 = \sqrt{2}v_1 - v_2, u_2 = v_1 - \sqrt{2}v_2$ and arbitrary $v_1, v_2 \in \mathbb{R}$ with $|v_1| + |v_2| > 0$:

$$x_1 = (u_1, u_2, u_2, u_1), \quad x_2 = (v_1, v_2, -v_2, -v_1).$$

Both components of \mathbf{x} are noninvertible, and $\sqrt{2}\mathbf{j}$ is not an eigenvalue of \mathbf{A} , but it is a candidate for an eigenvalue. \square

Now we come to the general case. In the first place we show that the eigenvalue problem can be expressed by two real, linear homogeneous 4×4 systems.

Lemma 4.3. *Let $\mathcal{A} = \mathbb{H}_{\text{coq}}$, \mathbf{A} defined in (3.5), \mathbf{x} defined in (4.2) and $\lambda = (\lambda_1, \lambda_2, \lambda_3, \lambda_4)$. The eigenvalue problem*

$$\mathbf{A}\mathbf{x} = \mathbf{x}\lambda$$

in \mathcal{A} can be given the real form

$$\mathbf{M}\mathbf{m} = \mathbf{0}, \quad \mathbf{N}\mathbf{n} = \mathbf{0}, \tag{4.4}$$

where \mathbf{M} and \mathbf{N} are two parameter dependent, real 4×4 matrices as follows:

$$\mathbf{M} := \begin{bmatrix} -1 & 0 & \lambda p_{2,3} & \lambda m_{1,4} \\ 0 & -1 & -\lambda p_{1,4} & \lambda m_{2,3} \\ \lambda m_{2,3} & \lambda p_{1,4} & -1 & 0 \\ -\lambda m_{1,4} & \lambda p_{2,3} & 0 & -1 \end{bmatrix}, \tag{4.5}$$

$$\mathbf{N} := \begin{bmatrix} -1 & 0 & -\lambda m_{2,3} & -(\lambda p_{1,4} + 2) \\ 0 & -1 & \lambda m_{1,4} + 2 & -\lambda p_{2,3} \\ \lambda p_{2,3} & \lambda m_{1,4} & -1 & 0 \\ -\lambda p_{1,4} & \lambda m_{2,3} & 0 & -1 \end{bmatrix}, \tag{4.6}$$

where

$$\mathbf{m} := (u_1 - u_4, u_2 - u_3, v_1 + v_4, v_2 + v_3)^T, \tag{4.7}$$

$$\mathbf{n} := (u_1 + u_4, u_2 + u_3, v_1 - v_4, v_2 - v_3)^T, \tag{4.8}$$

$$\lambda m_{1,4} := \lambda_1 - \lambda_4 - 1, \quad \lambda m_{2,3} := \lambda_2 - \lambda_3, \tag{4.9}$$

$$\lambda p_{1,4} := \lambda_1 + \lambda_4 - 1, \quad \lambda p_{2,3} := \lambda_2 + \lambda_3. \tag{4.10}$$

Proof. We apply the rules given in Table 2 for \mathbb{H}_{coq} and obtain

$$\begin{aligned} \mathbf{A}\mathbf{x} =: \mathbf{y} &= \begin{bmatrix} (u_1 - v_2, u_2 + v_1, u_3 - v_4, u_4 + v_3) \\ (u_3 + v_4, -u_4 + v_3, u_1 + v_2, -u_2 + v_1) \end{bmatrix} \\ &= \begin{bmatrix} (u_1, u_2, u_3, u_4)\lambda \\ (v_1, v_2, v_3, v_4)\lambda \end{bmatrix} \end{aligned} \tag{4.11}$$

The multiplication rule in \mathbb{H}_{coq} is

$$\begin{aligned}
 (u_1, u_2, u_3, u_4)\lambda &= (u_1\lambda_1 - u_2\lambda_2 + u_3\lambda_3 + u_4\lambda_4, \\
 &\quad u_1\lambda_2 + u_2\lambda_1 - u_3\lambda_4 + u_4\lambda_3, \\
 &\quad u_1\lambda_3 - u_2\lambda_4 + u_3\lambda_1 + u_4\lambda_2, \\
 &\quad u_1\lambda_4 + u_2\lambda_3 - u_3\lambda_2 + u_4\lambda_1).
 \end{aligned}$$

For $(v_1, v_2, v_3, v_4)\lambda$ the same rule applies. Comparing the four components of the two parts of (4.11) yields (after some reordering)

$$u_1 = v_1\lambda_3 - v_2(\lambda_4 + 1) + v_3\lambda_1 + v_4\lambda_2, \tag{4.12}$$

$$-u_2 = v_1(\lambda_4 - 1) + v_2\lambda_3 - v_3\lambda_2 + v_4\lambda_1, \tag{4.13}$$

$$u_3 = v_1\lambda_1 - v_2\lambda_2 + v_3\lambda_3 + v_4(\lambda_4 - 1), \tag{4.14}$$

$$-u_4 = v_1\lambda_2 + v_2\lambda_1 - v_3(\lambda_4 + 1) + v_4\lambda_3, \tag{4.15}$$

$$v_1 = u_1\lambda_2 + u_2(\lambda_1 - 1) - u_3\lambda_4 + u_4\lambda_3, \tag{4.16}$$

$$-v_2 = u_1(\lambda_1 - 1) - u_2\lambda_2 + u_3\lambda_3 + u_4\lambda_4, \tag{4.17}$$

$$v_3 = u_1\lambda_4 + u_2\lambda_3 - u_3\lambda_2 + u_4(\lambda_1 - 1), \tag{4.18}$$

$$-v_4 = u_1\lambda_3 - u_2\lambda_4 + u_3(\lambda_1 - 1) + u_4\lambda_2. \tag{4.19}$$

From these equations we deduce by adding and subtracting two sets of four linear equations each. The first set is

$$u_1 - u_4 = (\lambda_2 + \lambda_3)(v_1 + v_4) + (\lambda_1 - \lambda_4 - 1)(v_2 + v_3), \tag{4.20}$$

$$u_2 - u_3 = -(\lambda_1 + \lambda_4 - 1)(v_1 + v_4) + (\lambda_2 - \lambda_3)(v_2 + v_3), \tag{4.21}$$

$$v_1 + v_4 = (\lambda_2 - \lambda_3)(u_1 - u_4) + (\lambda_1 + \lambda_4 - 1)(u_2 - u_3), \tag{4.22}$$

$$v_2 + v_3 = -(\lambda_1 - \lambda_4 - 1)(u_1 - u_4) + (\lambda_2 + \lambda_3)(u_2 - u_3). \tag{4.23}$$

The second set is

$$u_1 + u_4 = -(\lambda_2 - \lambda_3)(v_1 - v_4) - (\lambda_1 + \lambda_4 + 1)(v_2 - v_3), \tag{4.24}$$

$$u_2 + u_3 = (\lambda_1 - \lambda_4 + 1)(v_1 - v_4) - (\lambda_2 + \lambda_3)(v_2 - v_3), \tag{4.25}$$

$$v_1 - v_4 = (\lambda_2 + \lambda_3)(u_1 + u_4) + (\lambda_1 - \lambda_4 - 1)(u_2 + u_3), \tag{4.26}$$

$$v_2 - v_3 = -(\lambda_1 + \lambda_4 - 1)(u_1 + u_4) + (\lambda_2 - \lambda_3)(u_2 + u_3). \tag{4.27}$$

The Eqs. (4.12)–(4.19) form a homogeneous linear system in the eight variables u_1, u_2, \dots, v_4 . If we choose λ at random, then we have to expect that the above 8×8 system has full rank and thus, λ is not a candidate for an eigenvalue. By adding and subtracting we can recover the solution of the 8×8 system from the solutions of the two 4×4 systems. In short form the two

4×4 systems defined by (4.20)–(4.23) and by (4.24)–(4.27) are the forms given in the lemma. \square

Lemma 4.4. *Let $\lambda \in \mathbb{H}_{\text{coq}}$ be arbitrary and let at least one of the matrices \mathbf{M}, \mathbf{N} have rank four. Then λ is not an eigenvalue.*

Proof. Let both matrices have rank four. Then, $\mathbf{m} = \mathbf{n} = \mathbf{0}$ (for definition see (4.7), (4.8)) which implies $u_1 = u_2, \dots = v_4 = 0$ (see (4.2)) and λ is not a candidate for an eigenvalue. If only one of the two matrices has rank four, then either $\mathbf{m} = \mathbf{0}$ or $\mathbf{n} = \mathbf{0}$, which implies that the two components x_1, x_2 of the eigenvector \mathbf{x} are both noninvertible. Thus, λ is not an eigenvalue. \square

Therefore, only those λ can qualify for an eigenvalue, which render the ranks of both \mathbf{M} and \mathbf{N} less than four.

Lemma 4.5. *Matrix \mathbf{M} has only the ranks two and four, matrix \mathbf{N} all ranks ≥ 2 . Matrix \mathbf{M} has rank two if and only if*

$$\lambda \in [1 + \mathbf{i}]. \tag{4.28}$$

Matrix \mathbf{N} has rank two if and only if

$$\lambda \in [\sqrt{2}\mathbf{j}] \text{ or } \lambda = \pm\sqrt{2}. \tag{4.29}$$

Matrix \mathbf{N} has rank three if and only if $\lambda \neq \pm\sqrt{2}, \lambda \notin [\sqrt{2}\mathbf{j}]$ and

$$\text{abs}_2((\lambda_1 \pm \sqrt{2}, \lambda_2, \lambda_3, \lambda_4)) = 0. \tag{4.30}$$

For the definition of abs_2 see (2.2). Formula (4.30) has the following special cases.

$$\lambda = (\lambda_1, 0, \lambda_3, 0), \quad \lambda_1\lambda_3 \neq 0 : (\lambda_1 \pm \sqrt{2})^2 = \lambda_3^2, \tag{4.31}$$

$$\lambda = (\lambda_1, \lambda_2, \lambda_3, \lambda_4), \quad \lambda_2^2 - \lambda_3^2 = 0, \quad \lambda_1\lambda_4 \neq 0 : (\lambda_1 \pm \sqrt{2})^2 = \lambda_4^2. \tag{4.32}$$

Proof. We put both matrices in block form $\mathbf{M}, \mathbf{N} = \begin{bmatrix} -\mathbf{I} & \mathbf{II} \\ \mathbf{III} & -\mathbf{I} \end{bmatrix}$, where all blocks consist of 2×2 matrices. Because of the presence of \mathbf{I} , which is the 2×2 identity matrix, the rank of both matrices \mathbf{M}, \mathbf{N} is at least two. Because of the similarity of the matrices $\mathbf{II}, \mathbf{III}$ of \mathbf{M} (they have the same trace and determinant), the rank of \mathbf{M} cannot be three. For \mathbf{N} all ranks ≥ 2 are possible. Rank two implies that there are two linear combinations of the first two columns which allow to represent the third and the fourth column. This restricts the set of possible λ and yields eventually (4.28) and (4.29). In more detail, for the case $\text{rank}(\mathbf{M}) = 2$ one has to solve

$$\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \end{bmatrix} = \begin{bmatrix} \lambda p_{2,3} \\ -\lambda p_{1,4} \end{bmatrix}, \quad \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = \begin{bmatrix} \lambda m_{1,4} \\ \lambda m_{2,3} \end{bmatrix}.$$

The solution is

$$a_1 = -\lambda p_{2,3}, \quad a_2 = \lambda p_{1,4}, \quad b_1 = -\lambda m_{1,4}, \quad b_2 = -\lambda m_{2,3}.$$

Let the four columns of \mathbf{M} be c_1, c_2, c_3, c_4 , then, the resulting conditions are

$$a_1c_1 + a_2c_2 = c_3; \quad b_1c_1 + b_2c_2 = c_4.$$

In explicit terms

$$-\lambda p_{2,3} \begin{bmatrix} -1 \\ 0 \\ \lambda m_{2,3} \\ -\lambda m_{1,4} \end{bmatrix} + \lambda p_{1,4} \begin{bmatrix} 0 \\ -1 \\ \lambda p_{1,4} \\ \lambda p_{2,3} \end{bmatrix} = \begin{bmatrix} \lambda p_{2,3} \\ -\lambda p_{1,4} \\ -1 \\ 0 \end{bmatrix},$$

$$-\lambda m_{1,4} \begin{bmatrix} -1 \\ 0 \\ \lambda m_{2,3} \\ -\lambda m_{1,4} \end{bmatrix} - \lambda m_{2,3} \begin{bmatrix} 0 \\ -1 \\ \lambda p_{1,4} \\ \lambda p_{2,3} \end{bmatrix} = \begin{bmatrix} \lambda m_{1,4} \\ \lambda m_{2,3} \\ 0 \\ -1 \end{bmatrix}.$$

By definition, only the last two rows are needed and imply

$$-\lambda p_{2,3} \lambda m_{2,3} + (\lambda p_{1,4})^2 = -1, \tag{4.33}$$

$$\lambda p_{2,3} \lambda m_{1,4} + \lambda p_{1,4} \lambda p_{2,3} = 0, \tag{4.34}$$

$$-\lambda m_{1,4} \lambda m_{2,3} - \lambda m_{2,3} \lambda p_{1,4} = 0, \tag{4.35}$$

$$(\lambda m_{1,4})^2 - \lambda m_{2,3} \lambda p_{2,3} = -1. \tag{4.36}$$

If we subtract the last from the first equation, we obtain

$$\lambda_4(\lambda_1 - 1) = 0.$$

The middle two equations can be written as

$$\lambda p_{2,3}(\lambda m_{1,4} + \lambda p_{1,4}) = 0,$$

$$\lambda m_{2,3}(\lambda m_{1,4} + \lambda p_{1,4}) = 0,$$

which is the same as

$$(\lambda_2 + \lambda_3)(\lambda_1 - 1) = 0,$$

$$(\lambda_2 - \lambda_3)(\lambda_1 - 1) = 0,$$

If we do not put $\lambda = 1$, it follows, that $\lambda_2 = \lambda_3 = \lambda_4 = 0$, however, this λ neither obeys (4.33) nor (4.36). So we have to choose $\lambda_1 = 1$, and $\lambda_2, \lambda_3, \lambda_4$ obeying (4.33), (4.36) results in $\lambda_2^2 - \lambda_3^2 - \lambda_4^2 = 1$, which is (4.28).

Now let us treat the case, where $\text{rank}(\mathbf{N}) = 2$. The technique to find the corresponding conditions is the same as for \mathbf{M} . The equations corresponding to (4.33)–(4.36) are

$$\lambda m_{2,3} \lambda p_{2,3} - (\lambda m_{1,4} + 2) \lambda m_{1,4} = -1, \tag{4.37}$$

$$-\lambda m_{2,3} \lambda p_{1,4} - (\lambda m_{1,4} + 2) \lambda m_{2,3} = 0, \tag{4.38}$$

$$(\lambda p_{1,4} + 2) \lambda p_{2,3} + \lambda p_{2,3} \lambda m_{1,4} = 0, \tag{4.39}$$

$$-(\lambda p_{1,4} + 2) \lambda p_{1,4} + \lambda p_{2,3} \lambda m_{2,3} = -1. \tag{4.40}$$

Subtracting (4.40) from (4.37) yields

$$(\lambda p_{14} - \lambda m_{1,4})(\lambda p_{1,4} + \lambda m_{1,4} + 2) = 0,$$

The middle two equations can be written as

$$\begin{aligned} \lambda m_{2,3}(\lambda p_{1,4} + \lambda m_{1,4} + 2) &= 0, \\ \lambda p_{2,3}(\lambda p_{1,4} + \lambda m_{1,4} + 2) &= 0. \end{aligned}$$

Let $\lambda p_{1,4} + \lambda m_{1,4} + 2 = 2\lambda_1 \neq 0$. Then $\lambda p_{1,4} - \lambda m_{1,4} = 2\lambda_4 = 0$, $\lambda m_{2,3} = \lambda_2 - \lambda_3 = 0$, $\lambda p_{2,3} = \lambda_2 + \lambda_3 = 0$. Hence; $\lambda_2 = \lambda_3 = \lambda_4 = 0$. Putting this in (4.37), (4.40) yields $\lambda_1 = \pm\sqrt{2}$. Now let $\lambda_1 = 0$. Then, (4.37), (4.40) imply $\lambda_2^2 - \lambda_3^2 - \lambda_4^2 = -2$ which is equivalent to $\lambda \in [\sqrt{2}\mathbf{j}]$. Thus, (4.29) is shown.

Let \mathbf{N} have rank three and denote, as before, the four columns of \mathbf{N} by c_1, c_2, c_3, c_4 . In order to find the linear combination to represent the last column by the first three columns we have to find a_1, a_2, a_3 from the first three rows and columns of $a_1c_1 + a_2c_2 + a_3c_3 = c_4$. Explicitly this reads

$$\begin{bmatrix} -1 & 0 & -\lambda m_{2,3} \\ 0 & -1 & \lambda m_{1,4} + 2 \\ \lambda p_{2,3} & \lambda m_{1,4} & -1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} -(\lambda p_{1,4} + 2) \\ -\lambda p_{2,3} \\ 0 \end{bmatrix}. \tag{4.41}$$

If the coefficients a_1, a_2, a_3 are known, then, by using the fourth row of \mathbf{N} , (see (4.6)) there is only one condition to satisfy:

$$-a_1\lambda p_{1,4} + a_2\lambda m_{2,3} = -1. \tag{4.42}$$

It remains, to find the coefficients. We reduce (4.41) to triangular form. Then, the third row becomes

$$[0, 0, (\lambda m_{1,4} + 2)\lambda m_{1,4} - (\lambda m_{2,3}\lambda p_{2,3} + 1)] \cdots = [-\lambda p_{2,3}(\lambda m_{1,4} + \lambda p_{1,4} + 2)].$$

The solution formula, by backward substitution, is, therefore,

$$((\lambda m_{1,4} + 2)\lambda m_{1,4} - (\lambda m_{2,3}\lambda p_{2,3} + 1))a_3 = -\lambda p_{2,3}(\lambda m_{1,4} + \lambda p_{1,4} + 2), \tag{4.43}$$

$$-a_2 + (\lambda m_{1,4} + 2)a_3 = -\lambda p_{2,3}, \tag{4.44}$$

$$-a_1 - \lambda m_{2,3}a_3 = -(\lambda p_{1,4} + 2). \tag{4.45}$$

We insert a_1 , from (4.45) and a_2 from (4.44) into the condition (4.42) and obtain

$$\underbrace{\{\lambda m_{2,3}a_3 - (\lambda p_{1,4} + 2)\}}_{-a_1} \lambda p_{1,4} + \underbrace{\{(\lambda m_{1,4} + 2)a_3 + \lambda p_{2,3}\}}_{a_2} \lambda m_{2,3} = -1.$$

Sorting yields

$$(\lambda m_{2,3}\lambda p_{1,4} + (\lambda m_{1,4} + 2)\lambda m_{2,3})a_3 = -1 + (\lambda p_{1,4} + 2)\lambda p_{1,4} - \lambda p_{2,3}\lambda m_{2,3}.$$

In terms of the components of λ , this condition reads

$$2\lambda_1(\lambda_2 - \lambda_3)a_3 = (\lambda_1 + \lambda_4)^2 - (\lambda_2^2 - \lambda_3^2) - 2.$$

The Eq. (4.44) for a_3 in terms of the λ components reads

$$((\lambda_1 - \lambda_4)^2 - (\lambda_2^2 - \lambda_3^2) - 2)a_3 = -2\lambda_1(\lambda_2 + \lambda_3). \tag{4.46}$$

We put only for this proof

$$A := \lambda_2^2 - \lambda_3^2 + 2$$

and note that the factor $(\lambda_1 - \lambda_4)^2 - A$ in (4.44) at a_3 cannot vanish, because the assumption that \mathbf{N} has rank three implies that the three coefficients a_1, a_2, a_3 are uniquely defined. Inserting a_3 into condition (4.46) yields

$$\begin{aligned} -4\lambda_1^2(\lambda_2^2 - \lambda_3^2) &= ((\lambda_1 + \lambda_4)^2 - A)((\lambda_1 - \lambda_4)^2 - A) \\ \Rightarrow -4\lambda_1^2(A - 2) &= (\lambda_1^2 - \lambda_4^2)^2 - 2A(\lambda_1^2 + \lambda_4^2) + A^2 \\ \Rightarrow 0 &= (\lambda_1^2 - \lambda_4^2 + A)^2 - 8\lambda_1^2 \\ &= (\lambda_1^2 - \lambda_4^2 + A - 2\sqrt{2}\lambda_1)(\lambda_1^2 - \lambda_4^2 + A + 2\sqrt{2}\lambda_1). \end{aligned} \tag{4.47}$$

which proves (4.30) and the special cases follow. □

We summarize our results.

Theorem 4.6. *The matrix $\mathbf{A} \in \mathbb{H}_{\text{coq}}^{2 \times 2}$ defined in (3.5) has no eigenvalues in \mathbb{H}_{coq} , the algebra of coquaternions.*

Proof. The previous Lemma 4.5 says, that λ defined by (4.28), (4.29), (4.30) are the only candidates for eigenvalues of \mathbf{A} over \mathbb{H}_{coq} . However, for all $\lambda \in \mathbb{H}_{\text{coq}}$ at least one of the matrices \mathbf{M}, \mathbf{N} has rank four. Thus, Lemma 4.4 applies and gives the final result. □

Lemma 4.7. *Let \mathbf{D} be a diagonal matrix of order $n \in \mathbb{N}$ over an algebra \mathcal{A} with diagonal elements d_1, d_2, \dots, d_n . Then all diagonal elements $d_j, j = 1, 2, \dots, n$ are eigenvalues of \mathbf{D} . Let \mathbf{T} be a triangular matrix (upper or lower) of order $n \in \mathbb{N}$ over an algebra \mathcal{A} with diagonal elements t_1, t_2, \dots, t_n . Then, at least the first diagonal element t_1 is an eigenvalue of \mathbf{T} .*

Proof. Let $\mathbf{e}_j \in \mathcal{A}^{n \times 1}$ be the j th unit vector, where the j th component consists of the algebraic one, and all other components consist of the algebraic zero. Then, $\mathbf{D}\mathbf{e}_j = d_j\mathbf{e}_j = \mathbf{e}_j d_j$. Therefore, d_j is a candidate for an eigenvalue with respect to \mathbf{e}_j , and since \mathbf{e}_j contains one invertible element, d_j is an eigenvalue for all $j = 1, 2, \dots, n$. Now, $\mathbf{T}\mathbf{e}_1 = t_1\mathbf{e}_1 = \mathbf{e}_1 t_1$, and t_1 is an eigenvalue of \mathbf{T} for the same reasons. □

Corollary 4.8. *The matrix \mathbf{A} , defined in (3.5) is neither diagonalizable nor triangulizable.*

Proof. Since similar matrices have the same set of eigenvalues, and following Lemma 4.7, diagonal and triangular matrices contain eigenvalues, they cannot be similar to a matrix without eigenvalues. □

In general terms that means, we cannot expect, that a matrix of order n over a nondivision algebra has a Schur decomposition.

In the formulas (4.28) to (4.30) we have listed all possible candidates for eigenvalues of \mathbf{A} (for \mathbf{A} see (3.5)), over \mathbb{H}_{coq} , the algebra of coquaternions, which by definition obey the equation $\mathbf{A}\mathbf{x} = \mathbf{x}\lambda, \mathbf{x} \neq \mathbf{0}$. And even if we count only similarity classes, we see from formula (4.30) that this number is infinite since distinct real parts λ_1 define distinct similarity classes. In a paper by Erdoğan and Özdemir [4], only candidates for eigenvalues over \mathbb{H}_{coq}

are considered. More recent information on problems related to quaternions and coquaternions with extensions to other algebras can be found in papers by the current authors in [8–13].

5. Eigenvalues in the Other \mathbb{R}^4 Algebras

In this section we give only a short overview on the problem whether the same matrix \mathbf{A} , defined in (3.5) has eigenvalues in the eight \mathbb{R}^4 algebras or not. If there are eigenvalues, we also present corresponding eigenvectors. Our results are mainly based on numerical computations which produce eigenvalues and eigenvectors. We would like to mention one principal difficulty for the noncommutative algebras. In this computation eigenvalues are computed with four components, but not in *reduced form*. What is reduced form? Theorem 1.3 says that $\lambda \in \sigma(\mathbf{A})$ implies $[\lambda] \subset \sigma(\mathbf{A})$ and because of the similarity identity relation (2.1) arbitrary eigenvalues in the noncommutative cases can be reduced to the short form (called *reduced form*):

- in \mathbb{H} : $\lambda = a + b\mathbf{i}$,
- in \mathbb{H}_{coq} and in \mathbb{H}_{nec} : $\lambda = a + b\mathbf{i}$, or $\lambda = a + b\mathbf{j}$,
- in \mathbb{H}_{con} : $\lambda = a + b\mathbf{i}$, or $\lambda = a + b\mathbf{k}$, with $b \geq 0$ in all cases.

See Lemma 2.2.

5.1. Quaternions \mathbb{H}

Let $\mathcal{A} = \mathbb{H}$. The matrix \mathbf{A} is invertible and

$$\mathbf{A}^{-1} = \frac{1}{2} \begin{bmatrix} 1 & -\mathbf{j} \\ -\mathbf{i} & -\mathbf{k} \end{bmatrix}.$$

There are two similarity classes of eigenvalues, defined by

$$\lambda_1 := -0.5(\sqrt{3} - 1) + 0.5(1 + \sqrt{3})\mathbf{i}, \quad \lambda_2 := 0.5(1 + \sqrt{3}) + 0.5(\sqrt{3} - 1)\mathbf{i}.$$

Eigenvector belonging to (similar to $\lambda_1 := -0.5(\sqrt{3} - 1) + 0.5(1 + \sqrt{3})\mathbf{i}$)

$$\begin{aligned} \lambda_1 &= (-0.366025403784439, -0.589308751210507, \\ &\quad 0.407721950377747, 1.162971801339703), \\ x_1 &= (-0.116136124516394, 0.096493215055665, \\ &\quad 0.431633726722537, -0.047102562471017), \\ x_2 &= (0.457810452774262, -0.094301599467715, \\ &\quad 0.222988193305220, 0.721434818892009), \end{aligned}$$

Eigenvector belonging to (similar to $\lambda_2 := 0.5(1 + \sqrt{3}) + 0.5(\sqrt{3} - 1)\mathbf{i}$)

$$\begin{aligned} \lambda_2 &= (1.366025403784439, 0.292559107612189, \\ &\quad -0.028933142042057, 0.218051915974833), \end{aligned}$$

$$\begin{aligned}
 x_1 &= (-0.429690495367038, 0.355279404544183, \\
 &\quad 0.182197205333501, 0.666799771941649), \\
 x_2 &= (0.063352481853988, 0.401343292837369, \\
 &\quad 0.086788018699258, -0.196730093155170).
 \end{aligned}$$

Since \mathbb{H} is a division algebra, both eigenvalues and all components of the eigenvectors are invertible.

5.2. Coquaternions \mathbb{H}_{coq}

Let $\mathcal{A} = \mathbb{H}_{\text{coq}}$. The matrix \mathbf{A} is invertible and

$$\mathbf{A}^{-1} = \frac{1}{2} \begin{bmatrix} 1 & \mathbf{j} \\ -\mathbf{i} & \mathbf{k} \end{bmatrix}.$$

See (4.1). However, there are no eigenvalues. This is the main result of this paper.

5.3. Tessarines \mathbb{H}_{tes}

Let $\mathcal{A} = \mathbb{H}_{\text{tes}}$. This algebra is commutative. The matrix \mathbf{A} is noninvertible. There are the following eigenvalues and eigenvectors:

$$\begin{aligned}
 \lambda_1 &= 0.5(1, -1, -1, 1), \\
 x_1 &= (-0.210579839771015, 0, 0.281296438799224, -0.613619136436347), \\
 x_2 &= (0.552747707503293, -0.271451268704069, \\
 &\quad 0.060871428933054, 0.342167867732278), \\
 \lambda_2 &= 0.5(1, 1, 1, 1), \\
 x_1 &= (-0.696986695461977, -0.026411017838051, \\
 &\quad 0.008934460636402, 0.115897281671691), \\
 x_2 &= (-0.272871967657917, -0.308217446132369, \\
 &\quad -0.415180267167659, 0.397703709966010).
 \end{aligned}$$

Both eigenvalues are not invertible, but all components of the eigenvectors are invertible in the algebra \mathbb{H}_{tes} .

5.4. Cotessarines $\mathbb{H}_{\text{cotes}}$

Let $\mathcal{A} = \mathbb{H}_{\text{cotes}}$. This algebra is commutative. The matrix \mathbf{A} is not invertible and the search for eigenvalues ended without success.

5.5. Nectarines \mathbb{H}_{nec}

Let $\mathcal{A} = \mathbb{H}_{\text{nec}}$. The matrix \mathbf{A} is invertible and

$$\mathbf{A}^{-1} = \frac{1}{2} \begin{bmatrix} 1 & -\mathbf{j} \\ \mathbf{i} & \mathbf{k} \end{bmatrix}.$$

The search for eigenvalues ended without success.

5.6. Conectarines \mathbb{H}_{con}

Let $\mathcal{A} = \mathbb{H}_{\text{con}}$. The matrix \mathbf{A} is invertible and

$$\mathbf{A}^{-1} = \frac{1}{2} \begin{bmatrix} 1 & \mathbf{j} \\ \mathbf{i} & -\mathbf{k} \end{bmatrix}.$$

The eigenvalues are

$$\lambda_1 := -0.5(\sqrt{3} - 1) + 0.5(1 + \sqrt{3})\mathbf{k}, \quad \lambda_2 := 0.5(1 + \sqrt{3}) + 0.5(\sqrt{3} - 1)\mathbf{k}.$$

The computed results are:

$$\begin{aligned} \lambda_1 &= (-0.366025403784439, 0.109391181348465, \\ &\quad 1.502346361480742, -2.033478896914270), \\ x_1 &= (0.084922455358793, 0.024284516474671, \\ &\quad -0.184640427677686, 0.411620233598415), \\ x_2 &= (0.219050248350320, 0.446277464435259, \\ &\quad -0.678289927178981, 0.285396781196364), \\ \lambda_2 &= (1.366025403784437, 1.565586637796200, \\ &\quad 2.519533443137362, -2.988826674089547), \\ x_1 &= (-0.586632672352341, -0.537466377582145, \\ &\quad 0.392122232068913, -0.043672536102042), \\ x_2 &= (-0.053199649549185, -0.198737203109869, \\ &\quad -0.230707718431950, 0.340253046200944). \end{aligned}$$

The eigenvalues and all components of the eigenvectors are invertible. For comparison, the first 20 digits of $0.5(\sqrt{3} - 1)$, $1 + 0.5(1 + \sqrt{3})$ are

$$0.36602\ 54037\ 84438\ 64676, \quad 1.36602\ 54037\ 84438\ 64676,$$

respectively, which shows, that the computed numerical values are almost precise until the last digit.

5.7. Tangerines \mathbb{H}_{tan}

Let $\mathcal{A} = \mathbb{H}_{\text{tan}}$. This algebra is commutative. The matrix \mathbf{A} is not invertible. There are the following eigenvalues:

$$\begin{aligned} \lambda_1 &= 0.5(1, 1, 1), \quad \lambda_2 = (0, 0, 0), \quad \lambda_3 = (1, 0, 0, 1), \\ \lambda_4 &= 0.5(1, -1, -1, 1), \end{aligned}$$

and the corresponding eigenvectors are:

$$\begin{aligned} x_1 &= (-0.355400905909388, -0.263459947802152, \\ &\quad 0.488848439984882, 0.255550884770960), \\ x_2 &= (-0.418170141431539, -0.326229183324303, \\ &\quad -0.192781649248809, -0.426079204462732), \\ x_3 &= (0.175671087331868, -0.481857471056402, \\ &\quad 0.180972667058841, -0.451887088154085), \end{aligned}$$

$$\begin{aligned}
x_2 &= (0.481857471056402, -0.175671087331868, \\
&\quad 0.451887088154085, -0.180972667058841), \\
x_1 &= (-0.487207160802510, 0.069582586598201, \\
&\quad 0.311355535868199, 0.401055078996997), \\
x_2 &= (-0.311355535868199, -0.401055078996997, \\
&\quad -0.487207160802510, 0.069582586598201), \\
x_1 &= (0.221176703745355, -0.216764176048914, \\
&\quad 0.118351465078905, 0.624569522483968), \\
x_2 &= (0.250902764854311, -0.255315292550752, \\
&\quad -0.152490053884302, -0.590430933678571).
\end{aligned}$$

All eigenvalues are not invertible, whereas all components of the eigenvectors are invertible.

5.8. Cotangerines \mathbb{H}_{\cotan}

Let $\mathcal{A} = \mathbb{H}_{\cotan}$. This algebra is commutative. The matrix \mathbf{A} is not invertible. The search for eigenvalues ended without success.

5.9. Concluding Remarks for this Section

These results, apart from those for coquaternions, are based on numerical computations. The statement *The search for eigenvalues ended without success* means that at least 100 trials with random initial values were carried out, in some cases even several hundreds. In contrast to the cases where there are eigenvalues, which were found in almost all cases within the first ten trials. So there is some likelihood, that not only \mathbf{A} over \mathbb{H}_{coq} does not contain eigenvalues, but also \mathbf{A} over $\mathbb{H}_{\text{cotes}}, \mathbb{H}_{\text{hec}}, \mathbb{H}_{\text{cotan}}$. And it even may be, that in all nondivision algebras \mathcal{A} there are matrices over \mathcal{A} without eigenvalues.

In order to facilitate the computations, we have used the *overloading technique* offered by MATLAB, which allows to use the standard algebra for matrices in all eight \mathbb{R}^4 algebras, and we steered the selection of the algebras by just one global integer parameter with values from one to eight. By this it is very easy to program Newton's method only once for finding eigenvalues and eigenvectors in all eight \mathbb{R}^4 algebras. A technique which is described by Lauterbach and Opfer in [14] was applied.

Acknowledgments

The research of G. Opfer was supported by the German Science Foundation, DFG, GZ: OP 33/19-1.

Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- [1] Brenner, J.L.: Matrices of quaternions. *Pac. J. Math.* **1**, 329–335 (1951)
- [2] Cockle, J.: On systems of algebra involving more than one imaginary; and on equations of the fifth degree. *Phil. Mag.* **35**(3), 434–437 (1849)
- [3] Cockle, J. <http://www.oocities.org/cocklebio/>
- [4] Erdoğan, M., Özdemir, M.: On eigenvalues of split quaternion matrices. *Adv. Appl. Clifford Alg.* **23**, 614–623 (2013)
- [5] Garling, D.J.H.: *Clifford algebras: an introduction*, p. 200. Cambridge University Press, Cambridge (2011)
- [6] Geometric algebra. http://en.wikipedia.org/wiki/Geometric_algebra
- [7] Horn, R.A., Johnson, C.R.: *Matrix analysis*, p. 561. Cambridge University Press, Cambridge (1992)
- [8] Janovská, D., Opfer, G.: Zeros and singular points for one-sided, coquaternionic polynomials with an extension to other \mathbb{R}^4 algebras. *ETNA* **41**, 133–158 (2014)
- [9] Janovská, D., Opfer, G.: Linear equations and the Kronecker product in coquaternions. *Mitt. Math. Ges. Hamburg* **33**, 181–196 (2013)
- [10] Janovská, D., Opfer, G.: The classification and the computation of the zeros of quaternionic, two-sided polynomials. *Numer. Math.* **115**, 81–100 (2010)
- [11] Janovská, D., Opfer, G.: A note on the computation of all zeros of simple quaternionic polynomials. *SIAM J. Numer. Anal.* **48**, 244–256 (2010)
- [12] Janovská, D., Opfer, G.: Linear equations in quaternionic variables. *Mitt. Math. Ges. Hamburg* **27**, 223–234 (2008)
- [13] Janovská, D., Opfer, G.: Computing quaternionic roots by Newton’s method. *Electron. Trans. Numer. Anal.* **26**, 82–102 (2007)
- [14] Lauterbach, R., Opfer, G.: The Jacobi matrix for functions in noncommutative algebras. *Adv. Appl. Clifford Alg.* **24**, 1059–1073 (2014). (**Erratum: Adv. Appl. Clifford Alg.** **24**, 1075 (2014))
- [15] Pogoruy, A.A., Rodríguez-Dagnino, R.M.: Some algebraic and analytic properties of coquaternion algebra. *Adv. Appl. Clifford Alg.* **20**, 79–84 (2010)
- [16] Schmeikal, B.: Tessarinen, Nektarinen und andere Vierheiten. Beweis einer Vermutung von Gerhard Opfer. *Mitt. Math. Ges. Hamburg* **34**, 81–108 (2014)
- [17] van der Waerden, B.L.: *Algebra*, 5th edn, p. 292. Springer, Berlin, Göttingen, Heidelberg (1960)
- [18] Wolf, L.A.: Similarity of matrices in which the elements are real quaternions. *Bull. Am. Math. Soc.* **42**, 737–743 (1936)
- [19] Zhang, F.: Quaternions and matrices of quaternions. *Linear Algebra Appl.* **251**, 21–57 (1997)

Drahoslava Janovská
University of Chemistry and Technology, Prague
Technická 5
166 28 Prague 6
Czech Republic
e-mail: janovskd@vscht.cz

Gerhard Opfer
Faculty for Mathematics, Informatics, and Natural Sciences [MIN]
University of Hamburg
Bundestraße 55
20146 Hamburg
Germany
e-mail: gerhard.opfer@uni-hamburg.de

Received: July 25, 2015.

Accepted: October 9, 2015.