



An outline of necrosome triggers

Tom Vanden Berghe^{1,2} · Behrouz Hassannia^{1,2,3} · Peter Vandenabeele^{1,2}

Received: 14 March 2016 / Accepted: 18 March 2016 / Published online: 6 April 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract Necroptosis was initially identified as a backup cell death program when apoptosis is blocked. However, it is now recognized as a cellular defense mechanism against infections and is presumed to be a detrimental factor in several pathologies driven by cell death. Necroptosis is a prototypic form of regulated necrosis that depends on activation of the necrosome, which is a protein complex in which receptor interacting protein kinase (RIPK) 3 is activated. The RIP homotypic interaction motif (RHIM) is the core domain that regulates activation of the necrosome. To date, three RHIM-containing proteins have been reported to activate the kinase activity of RIPK3 within the necrosome: RIPK1, Toll/IL-1 receptor domain-containing adaptor inducing IFN- β (TRIF), and DNA-dependent activator of interferon regulatory factors (DAI). Here, we review and discuss commonalities and differences of the increasing number of activators of the necrosome. Since the discovery that activation of mixed lineage kinase domain-like (MLKL) by RIPK3 kinase activity is crucial in necroptosis, interest has increased in monitoring and therapeutically targeting their activation. The availability of

new phospho-specific antibodies, pharmacologic inhibitors, and transgenic models will allow us to further document the role of necroptosis in degenerative, inflammatory and infectious diseases.

Keywords Necroptosis · RIPK1 · RIPK3 · MLKL · RHIM · Pathogens

Abbreviations

| | |
|-------|--|
| cIAP | Cellular inhibitor of apoptosis protein |
| cFLIP | Cellular FLICE-like inhibitory protein |
| CypD | Cyclophilin D |
| CYLD | Cylindromatosis |
| DAI | DNA-dependent activator of interferon regulatory factors |
| DR | Death receptor |
| FADD | Fas-associated protein via a death domain |
| FasL | Fas |
| MLKL | Mixed lineage kinase domain-like |
| Nec-1 | Necrostatin-1 |
| NLR | NOD-like receptor |
| NLRP3 | NOD-like receptor family pyrin domain-containing protein 3 |
| PAMP | Pathogen-associated molecular patterns |
| PARP1 | Poly(ADP-ribose) polymerase 1 |
| RHIM | RIP homotypic interaction motif |
| RIG-I | Retinoic acid-inducible gene-I |
| RIPK | Receptor-interacting protein kinase |
| ROS | Reactive oxygen species |
| TAK1 | Transforming growth factor- β -activated kinase 1 |
| TLR | Toll-like receptor |
| TNFR | Tumor necrosis factor receptor |
| TRADD | TNFR-associated death domain protein |

✉ Tom Vanden Berghe
Tom.VandenBerghe@irc.vib-ugent.be

✉ Peter Vandenabeele
Peter.Vandenabeele@irc.vib-ugent.be

¹ Inflammation Research Center, VIB, 9000 Ghent, Belgium

² Department of Biomedical Molecular Biology, Ghent University, 9000 Ghent, Belgium

³ Laboratory of Eukaryotic Gene Expression and Signal Transduction, Department of Physiology, Ghent University, 9000 Ghent, Belgium

| | |
|---------|--|
| TRAF2 | TNFR-associated factor 2 |
| TRAIL-R | TNF-related apoptosis-inducing ligand receptor |
| TRIF | Toll-interleukin-1 receptor domain-containing adaptor inducing interferon- β |
| TWEAKR | TNF-like weak inducer of apoptosis receptor |
| VV | Vaccinia virus |

Introduction

Rudolf Virchow (1821–1902, Prussia), founder of the cell theory (*Omnis cellula e cellula*) and cellular pathology, referred to tissue injury as “parenchymatous inflammation” and introduced the idea that tissue injury is caused by pathological changes within the cells. In 1858, he introduced the notion of cell death as a potential basis for pathology, with ‘necrobiosis’ being a physiological process of spontaneous wearing out of living parts from the body and ‘necrosis’ an accidental process. The term ‘necrosis’ comes from the Greek word ‘nekros,’ which means ‘dead body.’ Virchow’s necrobiosis–necrosis dichotomy resembles to some extent the current apoptosis–necrosis classification [1]. Together with cellular and molecular insights into inflammation came a shift in our understanding of the molecular interplay between cell death and inflammation at the site of tissue injury. This emerging field of research is crucial for understanding organismal homeostasis and how its processes contribute to a growing list of inflammatory and degenerative pathologies. Although cell death during inflammation was initially considered a manifestation of tissue damage, it was later recognized as a mechanism for eliminating pathogens and regulating inflammation by exposing or releasing molecular patterns that attract and alter the functions of other cells [2]. More recently, it became clear that phagocytosis of apoptotic cells can also initiate anti-inflammatory and tissue-regenerative responses [3, 4].

The current notion that not only apoptotic but also necrotic cell death is molecularly controlled by defined signaling mechanisms has increased the interest in studying regulated necrosis in the context of development, homeostasis and inflammation. Although RIPK3 knockout mice and MLKL knockout mice do not show an overt phenotypic abnormality under non-challenged conditions [5–8], it became clear that RIPK3-mediated necroptosis is a highly controlled cell death program that is executed when negative regulators such as caspase-8, IAPs or even RIPK1 are absent [9–17]. This suggests that the default mode during development and homeostasis is strong inhibition of the necroptosis pathway. Regulated necrosis, like passive necrosis due to physico-chemical insult, is

caused by loss of plasma membrane integrity leading to cellular rounding followed by swelling (oncosis). When the immune system recognizes cellular content exposed or released due to loss of membrane integrity, it initiates an inflammatory response. Regulated necrosis can be classified into several cell death modalities, such as necroptosis, parthanatos, ferroptosis, cyclophilin D-dependent necrosis, (n)etosis and pyroptosis [18]. Each type of regulated necrosis has particular biochemical features, yet it is not clear whether the common morphological features of these forms of cell death share or converge on common pathways. Other forms of cellular death are being identified, such as entosis [19], autosis [20, 21] and autoschizis [22].

Necroptosis, the best-characterized form of regulated necrosis, is mediated by the concerted action of receptor interacting protein kinase (RIPK) 3 and mixed lineage kinase domain-like (MLKL). In this review, we provide a snapshot of the activation of RIPK3 within the necrosome, typically by the three RHIM-containing proteins RIPK1, Toll/IL-1 receptor domain-containing adaptor inducing IFN- β (TRIF) and DNA-dependent activator of interferon regulatory factors (DAI). We briefly discuss the pluripotent roles of RIPK1 and RIPK3 in gene regulation and cell death induction.

RIPK1-dependent necroptosis

The mechanism of RIPK1-dependent necroptosis has been discovered mostly from studying tumor necrosis factor (TNF) signaling under conditions favoring cell death. This is because the major function of TNFR1, like that of DR3 (TRAMP/APO-3), is to induce pro-survival and pro-inflammatory genes, in contrast to some other death receptor family members such as CD95 (FAS/APO-1), TRAILR1 (DR4) and TRAILR2 (APO-2/TRICK/DR5/KILLER). The following model is currently proposed for TNFR1 signaling: sensing of trimeric TNF by TNFR1 induces the assembly of a primary receptor-bound complex that triggers activation of signaling pathways leading to gene induction [23–25]. Subsequently, assembly of a secondary TNFR1-unbound cytosolic complex induces cell death. For FAS and TRAILR1/2, the opposite situation is observed [26]. This sequential signaling provides a backup response by the secondary cytosolic complex in case the default pathway activated by the receptor-associated complex fails to resolve the infectious or inflammatory condition. Typically, pathogens and (epi)-genetic factors can interfere with gene activation or cell death induction. Thus, this sequential signaling probably evolved as a host defense against pathogens or conditions that might perturb either pathway.

In this review, we will briefly describe cell death signaling downstream of TNFR1, with a focus on necrosome activation. A cytosolic cell death-inducing complex is formed upon stimulation of TNFR1 only in conditions that sensitize to cell death, for example when cellular inhibitor of apoptosis proteins (c-IAPs) are absent (IAP-antagonist treatment), TAK1 or translation is inhibited, or RIPK1 is deubiquitinated [18]. This cytosolic complex (often referred to as complex II), which is composed of at least RIPK1, the death-fold-containing proteins Fas-associated protein via death domain (FADD), CASP8, and cFLIP (Fig. 1), induces either apoptosis or necroptosis. The formation and/or activity of complex II is tightly regulated by inhibitor κ B kinases (IKKs) through mechanisms that are either dependent or independent of NF- κ B [27].

Necroptosis is typically initiated when caspases are insufficiently activated or their activity is blocked, for example by pharmacological or viral inhibitors. This concept was first proposed based on studies of DR signaling in fibrosarcoma cells [28, 29], and later confirmed in vivo by the rescue of the embryonal lethality of CASP8 or FADD deficiency by RIPK3 depletion [9, 14, 16]. CASP8 cleaves and inactivates RIPK1 [30], RIPK3 [31] and cylindromatosis (CYLD) [32]. This could at least partially explain the protective role of CASP8 against necroptosis [33, 34]. It is thought that CYLD is crucial for translocation of RIPK1 from receptor-bound complex I to the cytosolic death-inducing complex II by removing ubiquitin chains from RIPK1 [35, 36]. However, recently it became clear that complex II is also ubiquitylated, although the E3




Fig. 1 Canonical versus noncanonical necrosome activation by three distinct RHIM-containing host adaptors. RIPK1 is a key pro-necrotic kinase that responds to several DRs, TCR, IFN- γ R and genotoxic stress by forming a RIPK1–RIPK3 complex through RHIM-dependent interactions (referred to as canonical necrosome). Necrostatin-1s (Nec-1s) blocks the cytotoxicity induced by the kinase activity of RIPK1. In TNFR1 signaling, cell death is efficiently induced only upon formation of a cytosolic cell death-inducing complex (complex II) when, for example, cellular inhibitor of apoptosis proteins (c-IAPs) are absent or depleted (IAP-antagonist treatment), TAK1 or translation is inhibited, or RIPK1 is deubiquitinated. In addition, caspases need to be inactivated or depleted, for example by pharmacological or viral inhibitors, to allow necroptosis to occur. TRIF and DAI are two other RHIM-containing adaptors that can activate RIPK3 via RHIM-dependent interactions in response to TLR3/4 and MCMV,

respectively (referred to as noncanonical necrosomes). RIPK3, upon activation by RIPK1, TRIF or DAI, phosphorylates itself and subsequently MLKL to stimulate oligomerization and translocation of MLKL to intracellular and plasma membranes. It is assumed that MLKL induces membrane rupture by binding directly to membrane phospholipids or indirectly with the help of calcium or sodium ion channels. Necrosulfonamide targets human MLKL and blocks necroptosis induction. DAI DNA-dependent activator of interferon regulatory factors, DR death receptor, FADD Fas-associated protein via a death domain, IFN interferon, MCMV murine cytomegalovirus, MLKL mixed lineage kinase domain-like, NSA necrosulfonamide, LPS lipopolysaccharide, RIPK receptor-interacting protein kinase, TCR T cell receptor, TLR Toll like receptor, TNF tumor necrosis factor, TRIF Toll-interleukin-1 receptor domain-containing adaptor inducing interferon- β

ligases have not been identified yet [37–41]. While complex II is formed through interactions that depend on the death effector domain (DED) and the death domain (DD), RIPK1 and RIPK3 interact through RIP homotypic interaction motifs (RHIM) [6, 42–45]. It is assumed that this RHIM-dependent binding of RIPK1 and RIPK3 involves a conformational change that releases the RHIM domain [46]; the conformational change depends on posttranslational modifications, particularly the phosphorylation and ubiquitination status. A series of auto- and cross-phosphorylations between RIPK1 and RIPK3 result in the formation and activation of the canonical necrosome [43, 47], which appears as an amyloid-like structure of RHIM-dependent oligomerized RIPK3 [48]. The phosphorylation of human RIPK3 at Ser227 and mouse RIPK3 at Ser232 is crucial for recruitment of mixed lineage kinase domain-like (MLKL) [49–52]. Subsequent phosphorylation of MLKL at Thr357/Ser358 by human RIPK3 [50] or at Ser345/Ser347/

Ser352/Thr349 by mouse RIPK3 [52] stimulates its oligomerization and translocation to intracellular and plasma membranes. The precise mechanism by which MLKL induces membrane rupture is controversial. Some reports implicate the influx of calcium or sodium through ion channels [53, 54] whereas others show direct binding to membrane phosphatidylinositol phosphates and loss of membrane integrity [55, 56].

In addition to TNF receptor signaling, other receptors induce necroptosis through RIPK1-dependent necrosome activation (Fig. 1; Table 1). These receptors include CD95L (FasL/APO-1L) [57], TRAIL (TNF-related apoptosis-inducing ligand or Apo2L) [57], TWEAK (TNF-like weak inducer of apoptosis) [58], and T cell receptor (TCR) [59]. Also, genotoxic stress [60, 61] and some anti-cancer drugs such as shikonin [62, 63] and obatoclox [64] have been shown to induce RIPK1-dependent necroptosis (Table 2). However, the assumption that many

Table 1 Overview of experimental findings of ligand-induced necroptosis involving MLKL

| Trigger | Major finding | References |
|---|---|------------|
| TNF | Discovery of MLKL downstream of RIPK3 | [50, 125] |
| | Quantitative phosphoproteomic analysis of RIP3-dependent protein phosphorylation | [126, 127] |
| | Trimerized MLKL acts at plasma membrane to allow Ca ²⁺ influx via TRPM7 | [53] |
| | MLKL unleashes four-helical bundle to induce membrane rupture | [128] |
| | Discovery of mouse MLKL targeting compound 1 | |
| | Depletion of RIPK3 or MLKL favors RIPK1 kinase-dependent apoptosis | [92] |
| | RIPK1 can function as an inhibitor rather than an initiator of RIPK3-dependent necroptosis | [129] |
| | Hsp90 and Cdc37 are required for RIPK3 activation | [130] |
| | Necroptosis is preceded by nuclear translocation of RIPK1, RIPK 3 and MLKL | [131] |
| | Radical scavengers attenuate necrosome assembly | [132] |
| | Ponatinib and pazopanib inhibit necroptosis by targeting RIPK1 | [133] |
| | Structure-guided design of potent and selective ponatinib-based hybrid inhibitors for RIPK1 | [134] |
| | Clnk promotes necroptosis downstream of RIPK3 and upstream of MLKL | [135] |
| | Intracellular NAD ⁺ promotes TNF-induced necroptosis in a sirtuin-dependent manner | [136] |
| | TRAF2 suppresses necrosome assembly | [117] |
| | MLKL and cofilin-1 translocation to mitochondria, Bax/Bak oligomerization, CypD and downregulation of Mcl-1 contribute to necroptosis | [137] |
| Ser345 is crucial for MLKL translocation and subsequent necroptosis | [138] | |
| LPS | RIPK3 promotes cell death and NLRP3 inflammasome activation in the absence of MLKL | [139] |
| | MLKL and PGAM5 upstream of NLRP3-mediated IL-1 β maturation in caspase-8-deficient dendritic cells | [94] |
| TLR1/2,3,4,7/8 ligands polyIC/ATP | Caspase blockade induces RIP3-mediated programmed necrosis in Toll-like receptor-activated microglia | [140] |
| | Caspase-8 scaffolding function and MLKL regulate NLRP3 inflammasome activation downstream of TLR3 | [141] |
| α -GalCer | Regulation of NKT cell-mediated immune responses to tumors and liver inflammation by mitochondrial PGAM5-Drp1 signaling | [142] |
| RIPK3 dimerization | Necroptosis is dependent on MLKL but not on DRP1 | [143] |
| MLKL dimerization | MLKL compromises membrane integrity by binding to phosphatidylinositol phosphates | [55] |
| RIPK1 and -3 oligomerization | RIPK3 homo-oligomerization results in amyloid scaffold, RIPK3 autophosphorylation and MLKL docking | [144–146] |
| CD40L | CD40L induces necrosome-dependent necroptosis in low-grade serous carcinomas | [147] |

Table 2 Overview of experimental findings of chemical-, toxin- or virus-induced necroptosis involving MLKL

| Trigger | Major finding | References |
|---------------------------------------|---|------------|
| BV6/MS275/ zVADfmk | Combination of SMAC-mimetics, HDAC inhibitors and caspase inhibition induces RIPK1- and MLKL-dependent necroptosis | [148] |
| NPe6-PDT | Photodynamic therapy using a high dose of the photosensitizer talaporfin sodium induces RIPK1, RIPK3 and MLKL-dependent necroptosis | [149] |
| CCCP | Impaired oxidative phosphorylation regulates MLKL-dependent necroptosis | [150] |
| Cigarette smoke extract | Nex-5 and Nec-1 protect against cell death induced by cigarette smoke extract | [151] |
| Shikonin | RIPK1- and RIPK3-dependent necroptosis | [152] |
| Obatoclox | Necrosome formation at autophagosomal membranes | [64] |
| Staurosporine/ zVADfmk | NSA blocks S/Z-induced necroptosis | [153] |
| Cytarabine/SMAC- mimetic | Nec-1 or NSA block TNF-mediated necroptosis | [154] |
| Thapsigargin tunicamycin | RIPK1/RIPK3/MLKL-mediated necroptosis in mouse fibrosarcoma cells | [155] |
| Edelfosine | RIPK1/RIPK3/MLKL-mediated necroptosis in human glioblastoma cells | [156] |
| Cisplatin | RIPK1-dependent necroptosis in squamous cell carcinoma | [157] |
| Doxorubicine etoposide | RIPK3-/MLKL-dependent cell death | [158] |
| Homoharringtonine | Sensitizes TRAIL-induced necroptosis | [159] |
| Acetylcholinesterase-R peptide | RIPK1/MLKL-mediated necroptosis in human granulosa cells | [160] |
| PMA | Neutrophil extracellular trap formation can involve RIPK1-RIPK3-MLKL signaling | [161] |
| | NET formation can occur independently of RIPK3 and MLKL signaling | [162] |
| CNOT3-deficiency | CNOT3 suppression promotes necroptosis by stabilizing mRNAs for cell death-inducing proteins | [163] |
| Neoalbacanol | Neoalbacanol induces cell death through necroptosis by regulating RIPK-dependent autocrine TNF | [164] |
| Chal-24 | A JNK-mediated autophagy pathway that triggers c-IAP degradation and necroptosis for anticancer chemotherapy | [165] |
| Hypoxia | Glycolytic pyruvate scavenging of mitochondrial superoxide protects against necroptosis | [166] |
| Honokiol | Honokiol induces RIPK3- and CypD-mediated necroptosis | [167] |
| RNA viruses | RIPK1/RIPK3/DRP1-mediated NLRP3 activation | [168] |
| CMV | DAI complexes with RIPK3 to mediate virus-induced programmed necrosis | [74] |
| | Human CMV blocks TNF-induced necroptosis downstream of RIPK3 and MLKL | [169] |
| HSV-1 | Mouse ICP6 triggers RIPK3/MLKL-dependent necroptosis | [170] |
| <i>Staphylococcus aureus</i> toxin | SA toxin triggers RIPK1/RIPK3/MLKL-dependent necroptosis | [107] |
| HSV-1 and -2 | Human, but not mouse, HSV-1 and -2 prevent necrosome-dependent necroptosis | [171–173] |
| HIV | HIV-1 protease cleaves RIPK1 and RIPK2 | [174] |

chemotherapeutics induce RIPK1/3-mediated necroptosis was recently challenged [65].

TRIF-dependent necroptosis

RIPK1 is the central RHIM-containing protein involved in the activation of RIPK3 during TNF-induced necroptosis, which leads to the formation of the so-called canonical necrosome complex. However, in response to some Toll-like receptors (TLRs), the RHIM-containing protein TRIF

somehow activates RIPK3 independently of RIPK1 [44, 45], leading to assembly of the noncanonical necrosome complex [46]. Each member of the TLR family senses particular pathogen-associated molecular patterns [66]. When activated, TLRs recruit adaptors containing the Toll/IL-1R (TIR) domain and initiate NF- κ B and IRF3/7 signaling that trigger the expression of cytokines, chemokines and interferons. All TLRs, with the exception of TLR3 and TLR4, mediate the signal through the adaptor myeloid differentiation primary-response gene 88 (MYD88). On the other hand, TLR3 and TLR4, after binding of the ligand

(dsRNA and LPS, respectively), recruit the RHIM domain containing adaptor TRIF [67]. Both TLR3 and TLR4 directly induce apoptosis or, if caspase activity is compromised, necroptosis [42, 68, 69]. Several other TLRs, such as TLR2, TLR5 and TLR9, also induce cell death but through an endocrine or paracrine TNF-dependent mechanism [70]. TLR3/4-induced necroptosis is critically dependent on RHIM-mediated recruitment of TRIF to activate RIPK3 (Fig. 1) [69, 70]. Like RIPK1 and RIPK3, TRIF is a cleavage substrate of CASP8 that inhibits its ability to stimulate NF- κ B-dependent cytokine expression [71]. Notably, RIPK1 seems to have a cell type specific function in TLR3/4-induced cell death. While fibroblast and endothelial cells undergo TLR3-dependent necroptosis independently of RIPK1, macrophages require RIPK1 to commit to TLR3/4-mediated necroptosis [70].

DAI-dependent necroptosis

In addition to RIPK1 and TRIF, a third RHIM-containing protein, DAI, has been reported to activate the necrosome (Fig. 1). The DAI pathway is typically activated in response to DNA viruses and leads to inhibition of viral replication [72]. Like TLR signaling, the intracellular DNA sensor activates the NF- κ B and IRF3 pathways to promote the synthesis of cytokines and interferons, which is dependent on RHIM-mediated recruitment of RIPK1 [73]. In addition, in response to DNA viruses, DAI induces necroptosis through RHIM-mediated activation of RIPK3 in the noncanonical necrosome [74]. As a virus-encoded countermeasure, the murine cytomegalovirus (CMV) M45-encoded viral inhibitor of RIP activation (vIRA) acts as a RHIM competitor and blocks necroptosis, which explains the virus's successful replication in the host. The potency of this cell autonomous host defense pathway is demonstrated by the remarkable attenuation of M45-deficient viruses in mice. Importantly, as in RIPK3-deficient mice, mCMV lacking M45 has the same pathogenesis in DAI-deficient mice, consistent with the notion of the existence of a DAI-RIPK3 complex as the natural target of M45 [74]. M45 encodes a ribonucleotide reductase (RNR) lacking enzymatic activity. Interestingly, many RNRs from herpesviruses also encode a RHIM [75]. This suggests that viral inhibitors that target the RIPKs via the RHIM represent a common viral evasion strategy.

The Janus faces of RIPK1

The 'two faces' of RIPK1 refers to its dual role. It has a cell death inhibitory role that is shown by the massive cell death observed in RIPK1-deficient models, whereas its

necroptosis-inducing capacity is executed by its kinase activity. In the absence of RIPK1, massive apoptosis is observed in cells [34, 76], in postnatal death knockout mice [77], and in intestinal specific knockout mice [12, 13]. It was initially thought that this was due to the role of RIPK1 in mediating NF- κ B activation, which results in the expression of survival genes such as *Flip_L* [77]. In this respect, cFLIP_L-CASP8 heterodimers have partial enzymatic activity, leading to incomplete cleavage of CASP8 [78, 79], and this consequently prevents apoptosis. Nevertheless, it is thought that CASP8 has some local activity within complex II resulting in cleavage of RIPKs and CYLD [80], which may contribute to the anti-necroptotic role of CASP8. However, mounting evidence questions the necessity or uniqueness of the role of NF- κ B activation in controlling cell death. For example, NF- κ B is still activated in response to TNF stimulation in the absence of RIPK1 in cultured MEF cells [81] and in intestinal organoids [13]. TAB2-deficient mice have a functional NF- κ B pathway, yet they die from massive liver apoptosis like mice deficient in p65, IKK β , TAK1 or NEMO [82]. Moreover, the rescue of mutant RIPK1 kinase-dead knockin mice from TNF-induced shock [10, 11, 83, 84] and from the lethal TNF-induced inflammation in Sharpin mutant mice [83] also calls into question the dominance of NF- κ B activation (that occurs in a RIPK1 kinase independent way). This is underscored by the recent finding that IKK α and IKK β control RIPK1-mediated cell death independently of NF- κ B activation [27].

The dual role of RIPK1 in controlling cell death is also illustrated by the perinatal death of RIPK1 knockout mice due to the aberrant activation of caspase-8 and RIPK3; mice lacking all three enzymes survived to adulthood [10, 14, 85]. Indeed, in addition to its anti-apoptotic role, RIPK1 also prevents RIPK3-driven necroptosis promoted by IFN and the TLR-adaptor TRIF [14]. Since RIPK1 is reported to be essential for RIPK3 activation and subsequent necroptosis induction by TNF, the identification of settings in which RIPK1 actively suppresses RIPK3 was surprising. Moreover, conditional depletion of RIPK1 leads to apoptosis in the intestine and necroptosis in the skin [12, 13]. This dynamic interplay and interdependence of these complex II components confers a crucial host-defense function to limit pathogen spread, especially when any one of these processes is disrupted [72, 86]. This may explain why this complex interrelationship exists and why ablation of specific elements (including RIPK1, FADD, caspase-8 and cFLIP) push the system to lethality [87]. In line with this reasoning, the tissues most affected by disruption of these gene products (intestine, lung, skin, endothelium, hematopoietic cells) represent crucial barriers to infection that are constantly engaged by pathogens [88]. Depending on the tissue, cell type and developmental stage, RIPK1 can certainly either activate or inhibit cell death.

Table 3 Overview of genetic necroptosis studies in experimental mouse pathologies

| Genetic defect | Major finding | References |
|---|--|------------|
| RIPK3 ^{KO} | Protects against pancreatitis | [6, 42] |
| | Sensitizes to vaccinia virus infection | [43] |
| | Protects against photoreceptor loss (+Nec-1) | [102] |
| | Protects against TNF-induced SIRS | [103] |
| | Partially protects from macrophage death and enhances bacterial control upon <i>Salmonella</i> infection | [104] |
| | Rescues the mid-gestation defect of cIap2/1-deficient embryos | [105] |
| | Protects against intracerebroventricular injection of TNF in mouse hippocampus | [106] |
| | Improves clearance of <i>Staphylococcus aureus</i> (+Nec-1s) | [107] |
| | Diminishes cutaneous wound healing | [108] |
| | Protects against ethanol-induced liver injury | [109] |
| | Protects from macrophage necroptosis in atherosclerosis development | [110, 111] |
| | Rescues mild systemic inflammatory disease in macrophage-specific CASP8 knockouts | [112] |
| | Protects against cone cell necrosis in a mouse model of inherited degeneration | [113] |
| | Protects against dsRNA-induced retinal degeneration | [114] |
| | Protects against myocardial infarction | [115] |
| | Protects against non-alcoholic steatohepatitis | [116] |
| | Partial rescue of TRAF-2-deficient mice | [117] |
| Protects against Gaucher's disease | [118] | |
| Protects against Ppm1b-mediated sensitization to TNF-induced SIRS | [119] | |
| Protects RIPK1E-KO from skin inflammation | [12] | |
| MLKL ^{KO} | Deficiency of RIPK3 or MLKL does not protect against shock induced by cecal ligation and puncture | [7] |
| | MLKL deficiency protects against pancreatitis | |
| | Nec-1 treatment and RIPK3 or MLKL deficiency protect against cisplatin-induced acute kidney injury | [120] |
| | Acetaminophen-induced liver toxicity occurs through RIPK1 but independently of RIPK3 and MLKL | [121, 122] |
| | No difference in diabetes onset by RIPK3- or MLKL-deficiency | [123] |
| RIPK3 or MLKL deficiency partially protects from inflammation induced by sharpin deficiency | [124] | |
| RIPK1 ^{IEC-KO} | RIPK1 protects the intestinal epithelium against apoptosis | [12, 13] |
| RIPK1 ^{E-KO} | RIPK1 partially protects the skin from necroptosis | [12] |
| RIPK3 ^{K51A} | Viable and fertile | [89] |
| RIPK3 ^{D161N} | Embryonic lethality rescued by Casp8-deficiency | [11] |
| RIPK1 ^{K45A} | Protect Sharpin-deficient mice from inflammation | [83] |
| RIPK1 ^{D138N} | Protects against TNF-induced shock | [11, 84] |
| | Does not protect against necroptosis of FADD-deficient IEC cells | [12] |
| RIPK1/3/CASP8 ^{KO} | Rescue Ripk1 ^{-/-} neonatal lethality by Ripk3 and Casp8 depletion | [14, 85] |
| RIPK3/CypD ^{KO} | Protects against renal ischemia reperfusion injury | [10] |
| | | [100] |

The pleiotropic role of RIPK3

Whereas RIPK3 knockout mice are viable and fertile [5, 89], RIPK3 D161N kinase dead knockin mice die on embryonic day E10.5 due to massive levels of apoptosis in the embryo and yolk sac vasculature [11]. But this was not observed in RIPK3 D51A kinase dead knockin mice Mandal et al. [89]. The embryonic death of RIPK3 D161N kinase dead knockin mice was rescued by ablation of RIPK1 or caspase-8, indicating that RIPK3 can engage both RIPK1 and caspase-8 [11]. It remains unclear

structurally why the D161N kinase-dead mutation in RIPK3 is proapoptotic, though it is likely that the kinase domain functionally “masks” the RHIM domain to prevent spurious activation [90]. In this scenario, the D161N alters the conformation of RIPK3 so that the RHIM domain is exposed for binding to RIPK1 to initiate apoptosis. This model predicts that the kinase and RHIM domains collaborate to control scaffolding of the necroptotic and apoptotic machineries. Some RIPK3 inhibitors were also found to induce apoptosis in a similar way through RHIM-dependent RIPK1 docking and subsequent FADD/CASP8-

Table 4 Overview of pharmacologic targeting of necroptosis in experimental rodent pathologies

| Pharmacologic agent | Major finding | References |
|-----------------------------|--|------------|
| Nec-1 | Alleviates ischemic brain injury | [47] |
| | Is cardioprotective | [175] |
| | Improves outcome after controlled cortical impact | [176] |
| | Alleviates retinal ischemia reperfusion injury | [177] |
| | Alleviates spinal cord injury pathology in rats | [178] |
| | Alleviates injury in subtotal nephrectomised rats | [179] |
| Nec-1s | Nec-1s treatment reduces disease severity in experimental autoimmune encephalitis model | [180] |
| NSA | Nec-1 or NSA block death in models of both sporadic and familial ALS (ex vivo) | [181] |
| Dabrafenib | Targets RIPK3 and alleviates acetaminophen-induced liver injury | [182] |
| | Upregulated RIPK3 expression potentiates MLKL phosphorylation-mediated programmed necrosis in toxic epidermal necrolysis | [183] |
| Nec-1, SangliferinA, 16-86 | Protects from renal ischemia reperfusion injury | [99] |
| Nec-1, cyclosporine A, 3-AB | Protects from remote lung injury after receiving ischemic renal allografts in rats | [101] |

Table 5 Overview of studies linking necroptosis to human pathologies

| Disease | Major finding | References |
|---------------------------------------|--|------------|
| Ovarian cancer | Low MLKL expression associated with poor prognosis | [184] |
| Inflammatory bowel disease | High RIPK3 and MLKL expression | [185] |
| Drug-induced liver injury | Phosphorylated MLKL in biopsies | [56] |
| Chronic obstructive pulmonary disease | Increased levels of RIPK3 in lung epithelial cells | [151] |
| Multiple sclerosis | Increased levels of RIPK1, RIPK3 and necrosome formation in lesions | [180] |
| Breast cancer | Weak RIPK3 expression | [158] |
| Melanoma | Melanoma cell lines lack RIPK3 expression, whereas primary melanocytes strongly express RIPK3 | [186] |
| HIV | Dysfunctional HIV-specific CD8+ T cell proliferation is associated with increased caspase-8 activity and mediated by necroptosis | [187] |
| Leukemia | RIPK3 is downregulated | [188] |

mediated apoptosis [70, 89]. Note that RIPK3 has also been reported to positively contribute to RIPK1-dependent apoptosis independently of its kinase activity but remarkably also of its RHIM domain [37]. While TNF signaling typically requires RIPK1 to activate RIPK3 in order to induce necroptosis, it has been noted that TNF can trigger RIPK3 activation even in the absence of RIPK1 if RIPK3 levels are high enough [91]. In the absence of RIPK1 and the presence of elevated levels of RIPK3, TNF can activate RIPK3 to induce cell death by both a caspase-8-dependent mechanism and a caspase-independent mechanism [37, 91]. Finally, similar to depletion of RIPK1 [34, 76], blocking TNF-induced necroptosis by suppressing RIPK3 or MLKL toggles the cell death response to apoptosis, albeit with different kinetics [92]. Collectively, these studies indicate that precise control of the complex II

machinery is necessary to prevent a lethal imbalance of necroptotic or apoptotic pathways.

Concluding remarks

There has been a revival of interest in the close interconnection between cell death and inflammation originally recognized by Virchow. This interconnection is emphasized by some recent findings that classical cell death inducers such as caspase-8 and RIPK3 seem to act also upstream of inflammasome activation in a cell autonomous way [93–98]. However, the precise mechanisms of this interaction are unclear. RIPK1 as well as RIPK3 and other cytosolic TNFR complex II components have been implicated in regulating cell death and inflammation, though if

these functions could be uncoupled is not clear. In addition, the potential signal transduction interplay between parenchymal cell necrosis and some forms of necrosis that occur in immune cells, such as pyroptosis and netosis, remains unknown. Considering the central role of RHIM domains in controlling the cell death induced by several stimuli, small molecules that disrupt RHIM signaling might also be therapeutically useful.

The number of genetic (Table 3) and pharmacological studies (Table 4) demonstrating an important role for RIPK1, RIPK3 or MLKL in murine experimental disease models is still increasing, highlighting the therapeutic potential of these necrosome members. In addition, the expression and activation of RIPK1, RIPK3 and MLKL is being increasingly explored in biopsies of patients with particular pathologies driven by cell death and inflammation (Table 5). The availability of new phospho-specific antibodies, pharmacologic inhibitors and transgenic models will allow us to document further the role of necroptosis in degenerative, inflammatory and infectious diseases. It is noteworthy that therapeutic targeting of only necroptosis might be insufficient in some complex pathologies, as exemplified by the additive protective effect of targeting different types of regulated necrosis [99–101]. This observed redundancy of necrosome proteins and interplay between different modalities of necrotic cell death in vivo is an intriguing topic for further research and will generate further insight into how the targeting of these molecules in some cases looks very effective.

Acknowledgments We thank Amin Bredan for excellent editorial help. TV is Assistant Professor at Ghent University. His research is supported by the Methusalem Grant of PV. Research in Vandenaabeele's unit is further supported by Belgian Grants (Interuniversity Attraction Poles, IAP 7/32), Flemish Grants (Research Foundation Flanders: FWO G.0875.11, FWO G.0973.11, FWO G.0A45.12N, FWO G.0787.13N, FWO G.0C31.14N, FWO G0E04.16N, Methusalem Grants—BOF09/01M00709 and BOF16/MET_V/007), Grants from Ghent University (MRP, GROUP-ID consortium), Foundation against Cancer, F94) and VIB.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Virchow R (1860) Cellular pathology: as based upon physiological and pathological histology. Twenty lectures delivered in the pathological Institute of Berlin during the months of February, March and April, 1858
- Wallach D, Kang TB, Kovalenko A (2014) Concepts of tissue injury and cell death in inflammation: a historical perspective. *Nat Rev Immunol* 14(1):51–59. doi:10.1038/nri3561
- Felton JM, Lucas CD, Rossi AG, Dransfield I (2014) Eosinophils in the lung—modulating apoptosis and efferocytosis in airway inflammation. *Front Immunol* 5:302. doi:10.3389/fimmu.2014.00302
- Saas P, Kaminski S, Perruche S (2013) Prospects of apoptotic cell-based therapies for transplantation and inflammatory diseases. *Immunotherapy* 5(10):1055–1073. doi:10.2217/imt.13.103
- Newton K, Sun X, Dixit VM (2004) Kinase RIP3 is dispensable for normal NF-kappa Bs, signaling by the B-cell and T-cell receptors, tumor necrosis factor receptor 1, and Toll-like receptors 2 and 4. *Mol Cell Biol* 24(4):1464–1469
- He S, Wang L, Miao L, Wang T, Du F, Zhao L, Wang X (2009) Receptor interacting protein kinase-3 determines cellular necrotic response to TNF-alpha. *Cell* 137(6):1100–1111. doi:10.1016/j.cell.2009.05.021
- Wu J, Huang Z, Ren J, Zhang Z, He P, Li Y, Ma J, Chen W, Zhang Y, Zhou X, Yang Z, Wu SQ, Chen L, Han J (2013) Mkl1 knockout mice demonstrate the indispensable role of Mkl1 in necroptosis. *Cell Res* 23(8):994–1006. doi:10.1038/cr.2013.91
- Murphy JM, Czabotar PE, Hildebrand JM, Lucet IS, Zhang JG, Alvarez-Diaz S, Lewis R, Lalaoui N, Metcalf D, Webb AI, Young SN, Varghese LN, Tannahill GM, Hatchell EC, Majewski JJ, Okamoto T, Dobson RC, Hilton DJ, Babon JJ, Nicola NA, Strasser A, Silke J, Alexander WS (2013) The pseudokinase MLKL mediates necroptosis via a molecular switch mechanism. *Immunity* 39(3):443–453. doi:10.1016/j.immuni.2013.06.018
- Kaiser WJ, Upton JW, Long AB, Livingston-Rosanoff D, Daley-Bauer LP, Hakem R, Caspari T, Mocarski ES (2011) RIP3 mediates the embryonic lethality of caspase-8-deficient mice. *Nature* 471(7338):368–372. doi:10.1038/nature09857
- Kaiser WJ, Daley-Bauer LP, Thapa RJ, Mandal P, Berger SB, Huang C, Sundararajan A, Guo H, Roback L, Speck SH, Bertin J, Gough PJ, Balachandran S, Mocarski ES (2014) RIP1 suppresses innate immune necrotic as well as apoptotic cell death during mammalian parturition. *Proc Natl Acad Sci USA* 111(21):7753–7758. doi:10.1073/pnas.1401857111
- Newton K, Dugger DL, Wickliffe KE, Kapoor N, de Almagro MC, Vucic D, Komuves L, Ferrando RE, French DM, Webster J, Roose-Girma M, Warming S, Dixit VM (2014) Activity of protein kinase RIPK3 determines whether cells die by necroptosis or apoptosis. *Science* 343(6177):1357–1360. doi:10.1126/science.1249361
- Dannappel M, Vlantis K, Kumari S, Polykratis A, Kim C, Wachsmuth L, Eftychi C, Lin J, Corona T, Hermance N, Zelic M, Kirsch P, Basic M, Bleich A, Kelliher M, Pasparakis M (2014) RIPK1 maintains epithelial homeostasis by inhibiting apoptosis and necroptosis. *Nature* 513(7516):90–94. doi:10.1038/nature13608
- Takahashi N, Vereecke L, Bertrand MJ, Duprez L, Berger SB, Divert T, Goncalves A, Sze M, Gilbert B, Kourula S, Goossens V, Lefebvre S, Gunther C, Becker C, Bertin J, Gough PJ, Declercq W, van Loo G, Vandenaabeele P (2014) RIPK1 ensures intestinal homeostasis by protecting the epithelium against apoptosis. *Nature* 513(7516):95–99. doi:10.1038/nature13706
- Dillon CP, Weinlich R, Rodriguez DA, Cripps JG, Quarato G, Gurung P, Verbist KC, Brewer TL, Llambi F, Gong YN, Janke LJ, Kelliher MA, Kanneganti TD, Green DR (2014) RIPK1 blocks early postnatal lethality mediated by caspase-8 and RIPK3. *Cell* 157(5):1189–1202. doi:10.1016/j.cell.2014.04.018
- Wong WW, Vince JE, Lalaoui N, Lawlor KE, Chau D, Bankovacki A, Anderton H, Metcalf D, O'Reilly L, Jost PJ, Murphy JM, Alexander WS, Strasser A, Vaux DL, Silke J

- (2014) cIAPs and XIAP regulate myelopoiesis through cytokine production in an RIPK1- and RIPK3-dependent manner. *Blood* 123(16):2562–2572. doi:[10.1182/blood-2013-06-510743](https://doi.org/10.1182/blood-2013-06-510743)
16. Oberst A, Dillon CP, Weinlich R, McCormick LL, Fitzgerald P, Pop C, Hakem R, Salvesen GS, Green DR (2011) Catalytic activity of the caspase-8-FLIP(L) complex inhibits RIPK3-dependent necrosis. *Nature* 471(7338):363–367. doi:[10.1038/nature09852](https://doi.org/10.1038/nature09852)
 17. Lu JV, Weist BM, van Raam BJ, Marro BS, Nguyen LV, Srinivas P, Bell BD, Luhrs KA, Lane TE, Salvesen GS, Walsh CM (2011) Complementary roles of Fas-associated death domain (FADD) and receptor interacting protein kinase-3 (RIPK3) in T-cell homeostasis and antiviral immunity. *Proc Natl Acad Sci USA* 108(37):15312–15317. doi:[10.1073/pnas.1102779108](https://doi.org/10.1073/pnas.1102779108)
 18. Vanden Berghe T, Linkermann A, Jouan-Lanhouet S, Walczak H, Vandenabeele P (2014) Regulated necrosis: the expanding network of non-apoptotic cell death pathways. *Nat Rev Mol Cell Biol* 15(2):135–147. doi:[10.1038/nrm3737](https://doi.org/10.1038/nrm3737)
 19. Kroemer G, Perfettini JL (2014) Entosis, a key player in cancer cell competition. *Cell Res* 24(11):1280–1281. doi:[10.1038/cr.2014.133](https://doi.org/10.1038/cr.2014.133)
 20. Liu Y, Levine B (2015) Autosis and autophagic cell death: the dark side of autophagy. *Cell Death Differ* 22(3):367–376. doi:[10.1038/cdd.2014.143](https://doi.org/10.1038/cdd.2014.143)
 21. Munoz-Pinedo C, Martin SJ (2014) Autosis: a new addition to the cell death tower of Babel. *Cell Death Dis* 5:e1319. doi:[10.1038/cddis.2014.246](https://doi.org/10.1038/cddis.2014.246)
 22. Jamison JM, Gilloteaux J, Taper HS, Calderon PB, Summers JL (2002) Autoschizis: a novel cell death. *Biochem Pharmacol* 63(10):1773–1783. doi:[10.1016/S0006-2952\(02\)09004-8](https://doi.org/10.1016/S0006-2952(02)09004-8)
 23. Vandenabeele P, Declercq W, Beyaert R, Fiers W (1995) Two tumour necrosis factor receptors: structure and function. *Trends Cell Biol* 5(10):392–399. doi:[10.1016/S0962-8924\(00\)89088-1](https://doi.org/10.1016/S0962-8924(00)89088-1)
 24. Micheau O, Tschopp J (2003) Induction of TNF receptor I-mediated apoptosis via two sequential signaling complexes. *Cell* 114(2):181–190
 25. Schneider-Brachert W, Tchikov V, Neumeyer J, Jakob M, Winoto-Morbach S, Held-Feindt J, Heinrich M, Merkel O, Ehrenschrwender M, Adam D, Mentlein R, Kabelitz D, Schutze S (2004) Compartmentalization of TNF receptor 1 signaling: internalized TNF receptosomes as death signaling vesicles. *Immunity* 21(3):415–428. doi:[10.1016/j.immuni.2004.08.017](https://doi.org/10.1016/j.immuni.2004.08.017)
 26. Vanden Berghe T, Kaiser WJ, Bertrand MJM, Vandenabeele P (2015) Molecular crosstalk between apoptosis, necroptosis, and survival signaling. *Mol Cell Oncol* 2(4):e975093. doi:[10.4161/23723556.2014.975093](https://doi.org/10.4161/23723556.2014.975093)
 27. Dondelinger Y, Jouan-Lanhouet S, Divert T, Theatre E, Bertin J, Gough PJ, Giansanti P, Heck AJ, Dejardin E, Vandenabeele P, Bertrand MJ (2015) NF- κ B-independent role of IKK α /IKK β in preventing RIPK1 kinase-dependent apoptotic and necroptotic cell death during TNF signaling. *Mol Cell* 60(1):63–76. doi:[10.1016/j.molcel.2015.07.032](https://doi.org/10.1016/j.molcel.2015.07.032)
 28. Vercammen D, Brouckaert G, Denecker G, Van de Craen M, Declercq W, Fiers W, Vandenabeele P (1998) Dual signaling of the Fas receptor: initiation of both apoptotic and necrotic cell death pathways. *J Exp Med* 188(5):919–930
 29. Vercammen D, Beyaert R, Denecker G, Goossens V, Van Loo G, Declercq W, Grooten J, Fiers W, Vandenabeele P (1998) Inhibition of caspases increases the sensitivity of L929 cells to necrosis mediated by tumor necrosis factor. *J Exp Med* 187(9):1477–1485
 30. Lin Y, Devin A, Rodriguez Y, Liu ZG (1999) Cleavage of the death domain kinase RIP by caspase-8 prompts TNF-induced apoptosis. *Genes Dev* 13(19):2514–2526
 31. Feng S, Yang Y, Mei Y, Ma L, Zhu DE, Hoti N, Castanares M, Wu M (2007) Cleavage of RIP3 inactivates its caspase-independent apoptosis pathway by removal of kinase domain. *Cell Signal* 19(10):2056–2067. doi:[10.1016/j.cellsig.2007.05.016](https://doi.org/10.1016/j.cellsig.2007.05.016)
 32. O'Donnell MA, Perez-Jimenez E, Oberst A, Ng A, Massoumi R, Xavier R, Green DR, Ting AT (2011) Caspase 8 inhibits programmed necrosis by processing CYLD. *Nat Cell Biol* 13(12):1437–1442. doi:[10.1038/ncb2362](https://doi.org/10.1038/ncb2362)
 33. Salvesen GS, Walsh CM (2014) Functions of caspase 8: the identified and the mysterious. *Semin Immunol* 26(3):246–252. doi:[10.1016/j.smim.2014.03.005](https://doi.org/10.1016/j.smim.2014.03.005)
 34. Vanlangenakker N, Bertrand MJ, Bogaert P, Vandenabeele P, Vanden Berghe T (2011) TNF-induced necroptosis in L929 cells is tightly regulated by multiple TNFR1 complex I and II members. *Cell Death Dis* 2:e230. doi:[10.1038/cddis.2011.111](https://doi.org/10.1038/cddis.2011.111)
 35. O'Donnell MA, Legarda-Addison D, Skountzou P, Yeh WC, Ting AT (2007) Ubiquitination of RIP1 regulates an NF- κ B-independent cell-death switch in TNF signaling. *Curr Biol* 17(5):418–424. doi:[10.1016/j.cub.2007.01.027](https://doi.org/10.1016/j.cub.2007.01.027)
 36. Wright A, Reiley WW, Chang M, Jin W, Lee AJ, Zhang M, Sun SC (2007) Regulation of early wave of germ cell apoptosis and spermatogenesis by deubiquitinating enzyme CYLD. *Dev Cell* 13(5):705–716. doi:[10.1016/j.devcel.2007.09.007](https://doi.org/10.1016/j.devcel.2007.09.007)
 37. Dondelinger Y, Aguilera MA, Goossens V, Dubuisson C, Grootjans S, Dejardin E, Vandenabeele P, Bertrand MJ (2013) RIPK3 contributes to TNFR1-mediated RIPK1 kinase-dependent apoptosis in conditions of cIAP1/2 depletion or TAK1 kinase inhibition. *Cell Death Differ* 20(10):1381–1392. doi:[10.1038/cdd.2013.94](https://doi.org/10.1038/cdd.2013.94)
 38. Moquin DM, McQuade T, Chan FK (2013) CYLD deubiquitinates RIP1 in the TNF α -induced necrosome to facilitate kinase activation and programmed necrosis. *PLoS One* 8(10):e76841. doi:[10.1371/journal.pone.0076841](https://doi.org/10.1371/journal.pone.0076841)
 39. Onizawa M, Oshima S, Schulze-Topphoff U, Osés-Prieto JA, Lu T, Tavares R, Prodhomme T, Duong B, Whang MI, Advincula R, Agelidis A, Barrera J, Wu H, Burlingame A, Malynn BA, Zamvil SS, Ma A (2015) The ubiquitin-modifying enzyme A20 restricts ubiquitination of the kinase RIPK3 and protects cells from necroptosis. *Nat Immunol* 16(6):618–627. doi:[10.1038/ni.3172](https://doi.org/10.1038/ni.3172)
 40. Biton S, Ashkenazi A (2011) NEMO and RIP1 control cell fate in response to extensive DNA damage via TNF- α feedforward signaling. *Cell* 145(1):92–103. doi:[10.1016/j.cell.2011.02.023](https://doi.org/10.1016/j.cell.2011.02.023)
 41. de Almagro MC, Goncharov T, Newton K, Vucic D (2015) Cellular IAP proteins and LUBAC differentially regulate necrosome-associated RIP1 ubiquitination. *Cell Death Dis* 6:e1800. doi:[10.1038/cddis.2015.158](https://doi.org/10.1038/cddis.2015.158)
 42. Zhang DW, Shao J, Lin J, Zhang N, Lu BJ, Lin SC, Dong MQ, Han J (2009) RIP3, an energy metabolism regulator that switches TNF-induced cell death from apoptosis to necrosis. *Science* 325(5938):332–336. doi:[10.1126/science.1172308](https://doi.org/10.1126/science.1172308)
 43. Cho YS, Challa S, Moquin D, Genga R, Ray TD, Guildford M, Chan FK (2009) Phosphorylation-driven assembly of the RIP1-RIP3 complex regulates programmed necrosis and virus-induced inflammation. *Cell* 137(6):1112–1123. doi:[10.1016/j.cell.2009.05.037](https://doi.org/10.1016/j.cell.2009.05.037)
 44. Guo H, Kaiser WJ, Mocarski ES (2015) Manipulation of apoptosis and necroptosis signaling by herpesviruses. *Med Microbiol Immunol* 204(3):439–448. doi:[10.1007/s00430-015-0410-5](https://doi.org/10.1007/s00430-015-0410-5)
 45. Mocarski ES, Guo H, Kaiser WJ (2015) Necroptosis: the Trojan horse in cell autonomous antiviral host defense. *Virology* 479–480:160–166. doi:[10.1016/j.virol.2015.03.016](https://doi.org/10.1016/j.virol.2015.03.016)
 46. Chan FK, Luz NF, Moriwaki K (2015) Programmed necrosis in the cross talk of cell death and inflammation. *Annu Rev*

- Immunol 33:79–106. doi:[10.1146/annurev-immunol-032414-112248](https://doi.org/10.1146/annurev-immunol-032414-112248)
47. Degtarev A, Huang Z, Boyce M, Li Y, Jagtap P, Mizushima N, Cuny GD, Mitchison TJ, Moskowitz MA, Yuan J (2005) Chemical inhibitor of nonapoptotic cell death with therapeutic potential for ischemic brain injury. *Nat Chem Biol* 1(2):112–119. doi:[10.1038/nchembio711](https://doi.org/10.1038/nchembio711)
 48. Li J, McQuade T, Siemer AB, Napetschnig J, Moriwaki K, Hsiao YS, Damko E, Moquin D, Walz T, McDermott A, Chan FK, Wu H (2012) The RIP1/RIP3 necrosome forms a functional amyloid signaling complex required for programmed necrosis. *Cell* 150(2):339–350. doi:[10.1016/j.cell.2012.06.019](https://doi.org/10.1016/j.cell.2012.06.019)
 49. Chen W, Zhou Z, Li L, Zhong CQ, Zheng X, Wu X, Zhang Y, Ma H, Huang D, Li W, Xia Z, Han J (2013) Diverse sequence determinants control human and mouse receptor interacting protein 3 (RIP3) and mixed lineage kinase domain-like (MLKL) interaction in necroptotic signaling. *J Biol Chem* 288(23):16247–16261. doi:[10.1074/jbc.M112.435545](https://doi.org/10.1074/jbc.M112.435545)
 50. Sun L, Wang H, Wang Z, He S, Chen S, Liao D, Wang L, Yan J, Liu W, Lei X, Wang X (2012) Mixed lineage kinase domain-like protein mediates necrosis signaling downstream of RIP3 kinase. *Cell* 148(1–2):213–227. doi:[10.1016/j.cell.2011.11.031](https://doi.org/10.1016/j.cell.2011.11.031)
 51. McQuade T, Cho Y, Chan FK (2013) Positive and negative phosphorylation regulates RIP1- and RIP3-induced programmed necrosis. *Biochem J* 456(3):409–415. doi:[10.1042/BJ20130860](https://doi.org/10.1042/BJ20130860)
 52. Xie T, Peng W, Yan C, Wu J, Gong X, Shi Y (2013) Structural insights into RIP3-mediated necroptotic signaling. *Cell Rep* 5(1):70–78. doi:[10.1016/j.celrep.2013.08.044](https://doi.org/10.1016/j.celrep.2013.08.044)
 53. Cai Z, Jitkaew S, Zhao J, Chiang HC, Choksi S, Liu J, Ward Y, Wu LG, Liu ZG (2014) Plasma membrane translocation of trimerized MLKL protein is required for TNF-induced necroptosis. *Nat Cell Biol* 16(1):55–65. doi:[10.1038/ncb2883](https://doi.org/10.1038/ncb2883)
 54. Chen X, Li W, Ren J, Huang D, He WT, Song Y, Yang C, Li W, Zheng X, Chen P, Han J (2014) Translocation of mixed lineage kinase domain-like protein to plasma membrane leads to necrotic cell death. *Cell Res* 24(1):105–121. doi:[10.1038/cr.2013.171](https://doi.org/10.1038/cr.2013.171)
 55. Dondelinger Y, Declercq W, Montessuit S, Roelandt R, Gonçalves A, Bruggeman I, Hulpiau P, Weber K, Sehon CA, Marquis RW, Bertin J, Gough PJ, Savvides S, Martinou JC, Bertrand MJ, Vandenabeele P (2014) MLKL compromises plasma membrane integrity by binding to phosphatidylinositol phosphates. *Cell Rep* 7(4):971–981. doi:[10.1016/j.celrep.2014.04.026](https://doi.org/10.1016/j.celrep.2014.04.026)
 56. Wang H, Sun L, Su L, Rizo J, Liu L, Wang LF, Wang FS, Wang X (2014) Mixed lineage kinase domain-like protein MLKL causes necrotic membrane disruption upon phosphorylation by RIP3. *Mol Cell* 54(1):133–146. doi:[10.1016/j.molcel.2014.03.003](https://doi.org/10.1016/j.molcel.2014.03.003)
 57. Holler N, Zaru R, Micheau O, Thome M, Attinger A, Valitutti S, Bodmer JL, Schneider P, Seed B, Tschopp J (2000) Fas triggers an alternative, caspase-8-independent cell death pathway using the kinase RIP as effector molecule. *Nat Immunol* 1(6):489–495. doi:[10.1038/82732](https://doi.org/10.1038/82732)
 58. Wilson CA, Browning JL (2002) Death of HT29 adenocarcinoma cells induced by TNF family receptor activation is caspase-independent and displays features of both apoptosis and necrosis. *Cell Death Differ* 9(12):1321–1333. doi:[10.1038/sj.cdd.4401107](https://doi.org/10.1038/sj.cdd.4401107)
 59. Ch'en IL, Beisner DR, Degtarev A, Lynch C, Yuan J, Hoffmann A, Hedrick SM (2008) Antigen-mediated T cell expansion regulated by parallel pathways of death. *Proc Natl Acad Sci USA* 105(45):17463–17468. doi:[10.1073/pnas.0808043105](https://doi.org/10.1073/pnas.0808043105)
 60. Tenev T, Bianchi K, Darding M, Broemer M, Langlais C, Wallberg F, Zachariou A, Lopez J, MacFarlane M, Cain K, Meier P (2011) The Ripoptosome, a signaling platform that assembles in response to genotoxic stress and loss of IAPs. *Mol Cell* 43(3):432–448. doi:[10.1016/j.molcel.2011.06.006](https://doi.org/10.1016/j.molcel.2011.06.006)
 61. Feoktistova M, Geserick P, Kellert B, Dimitrova DP, Langlais C, Hupe M, Cain K, MacFarlane M, Hacker G, Leverkus M (2011) cIAPs block Ripoptosome formation, a RIP1/caspase-8 containing intracellular cell death complex differentially regulated by cFLIP isoforms. *Mol Cell* 43(3):449–463. doi:[10.1016/j.molcel.2011.06.011](https://doi.org/10.1016/j.molcel.2011.06.011)
 62. Han W, Li L, Qiu S, Lu Q, Pan Q, Gu Y, Luo J, Hu X (2007) Shikonin circumvents cancer drug resistance by induction of a necroptotic death. *Mol Cancer Ther* 6(5):1641–1649. doi:[10.1158/1535-7163.MCT-06-0511](https://doi.org/10.1158/1535-7163.MCT-06-0511)
 63. Huang C, Luo Y, Zhao J, Yang F, Zhao H, Fan W, Ge P (2013) Shikonin kills glioma cells through necroptosis mediated by RIP-1. *PLoS One* 8(6):e66326. doi:[10.1371/journal.pone.0066326](https://doi.org/10.1371/journal.pone.0066326)
 64. Basit F, Cristofanon S, Fulda S (2013) Obatoclox (GX15-070) triggers necroptosis by promoting the assembly of the necrosome on autophagosomal membranes. *Cell Death Differ* 20(9):1161–1173. doi:[10.1038/cdd.2013.45](https://doi.org/10.1038/cdd.2013.45)
 65. Moriwaki K, Bertin J, Gough PJ, Orłowski GM, Chan FK (2015) Differential roles of RIPK1 and RIPK3 in TNF-induced necroptosis and chemotherapeutic agent-induced cell death. *Cell Death Dis* 6:e1636. doi:[10.1038/cddis.2015.16](https://doi.org/10.1038/cddis.2015.16)
 66. Gay NJ, Symmons MF, Gangloff M, Bryant CE (2014) Assembly and localization of Toll-like receptor signalling complexes. *Nat Rev Immunol* 14(8):546–558. doi:[10.1038/nri3713](https://doi.org/10.1038/nri3713)
 67. Kaiser WJ, Offermann MK (2005) Apoptosis induced by the Toll-like receptor adaptor TRIF is dependent on its receptor interacting protein homotypic interaction motif. *J Immunol* 174(8):4942–4952
 68. Kalai M, Van Loo G, Vanden Berghe T, Meeus A, Burm W, Saelens X, Vandenabeele P (2002) Tipping the balance between necrosis and apoptosis in human and murine cells treated with interferon and dsRNA. *Cell Death Differ* 9(9):981–994. doi:[10.1038/sj.cdd.4401051](https://doi.org/10.1038/sj.cdd.4401051)
 69. He S, Liang Y, Shao F, Wang X (2011) Toll-like receptors activate programmed necrosis in macrophages through a receptor-interacting kinase-3-mediated pathway. *Proc Natl Acad Sci USA* 108(50):20054–20059. doi:[10.1073/pnas.1116302108](https://doi.org/10.1073/pnas.1116302108)
 70. Kaiser WJ, Sridharan H, Huang C, Mandal P, Upton JW, Gough PJ, Sehon CA, Marquis RW, Bertin J, Mocarski ES (2013) Toll-like receptor 3-mediated necrosis via TRIF, RIP3, and MLKL. *J Biol Chem* 288(43):31268–31279. doi:[10.1074/jbc.M113.462341](https://doi.org/10.1074/jbc.M113.462341)
 71. Rebsamen M, Meylan E, Curran J, Tschopp J (2008) The antiviral adaptor proteins Cardif and Trif are processed and inactivated by caspases. *Cell Death Differ* 15(11):1804–1811. doi:[10.1038/cdd.2008.119](https://doi.org/10.1038/cdd.2008.119)
 72. Upton JW, Chan FK (2014) Staying alive: cell death in antiviral immunity. *Mol Cell* 54(2):273–280. doi:[10.1016/j.molcel.2014.01.027](https://doi.org/10.1016/j.molcel.2014.01.027)
 73. Kaiser WJ, Upton JW, Mocarski ES (2008) Receptor-interacting protein homotypic interaction motif-dependent control of NF- κ B activation via the DNA-dependent activator of IFN regulatory factors. *J Immunol* 181(9):6427–6434
 74. Upton JW, Kaiser WJ, Mocarski ES (2012) DAI/ZBP1/DLM-1 complexes with RIP3 to mediate virus-induced programmed necrosis that is targeted by murine cytomegalovirus vIRA. *Cell Host Microbe* 11(3):290–297. doi:[10.1016/j.chom.2012.01.016](https://doi.org/10.1016/j.chom.2012.01.016)
 75. Lembo D, Brune W (2009) Tinkering with a viral ribonucleotide reductase. *Trends Biochem Sci* 34(1):25–32. doi:[10.1016/j.tibs.2008.09.008](https://doi.org/10.1016/j.tibs.2008.09.008)
 76. Vanden Berghe T, Kalai M, van Loo G, Declercq W, Vandenabeele P (2003) Disruption of HSP90 function reverts tumor necrosis factor-induced necrosis to apoptosis. *J Biol Chem* 278(8):5622–5629. doi:[10.1074/jbc.M208925200](https://doi.org/10.1074/jbc.M208925200)

77. Kelliher MA, Grimm S, Ishida Y, Kuo F, Stanger BZ, Leder P (1998) The death domain kinase RIP mediates the TNF-induced NF- κ B signal. *Immunity* 8(3):297–303
78. Wachter T, Sprick M, Hausmann D, Kerstan A, McPherson K, Stassi G, Brocker EB, Walczak H, Leverkus M (2004) cFLIPL inhibits tumor necrosis factor-related apoptosis-inducing ligand-mediated NF- κ B activation at the death-inducing signaling complex in human keratinocytes. *J Biol Chem* 279(51):52824–52834. doi:10.1074/jbc.M409554200
79. Scaffidi C, Schmitz I, Kramer PH, Peter ME (1999) The role of c-FLIP in modulation of CD95-induced apoptosis. *J Biol Chem* 274(3):1541–1548
80. Feoktistova M, Leverkus M (2015) Programmed necrosis and necroptosis signalling. *FEBS J* 282(1):19–31. doi:10.1111/febs.13120
81. Wong WW, Gentle IE, Nachbur U, Anderton H, Vaux DL, Silke J (2010) RIPK1 is not essential for TNFR1-induced activation of NF- κ B. *Cell Death Differ* 17(3):482–487. doi:10.1038/cdd.2009.178
82. Sanjo H, Takeda K, Tsujimura T, Ninomiya-Tsuji J, Matsumoto K, Akira S (2003) TAB2 is essential for prevention of apoptosis in fetal liver but not for interleukin-1 signaling. *Mol Cell Biol* 23(4):1231–1238. doi:10.1128/mcb.23.4.1231-1238.2003
83. Berger SB, Kasparcova V, Hoffman S, Swift B, Dare L, Schaeffer M, Capriotti C, Cook M, Finger J, Hughes-Earle A, Harris PA, Kaiser WJ, Mocarski ES, Bertin J, Gough PJ (2014) Cutting edge: RIP1 kinase activity is dispensable for normal development but is a key regulator of inflammation in SHARPIN-deficient mice. *J Immunol* 192(12):5476–5480. doi:10.4049/jimmunol.1400499
84. Polykratis A, Hermance N, Zelic M, Roderick J, Kim C, Van TM, Lee TH, Chan FK, Pasparakis M, Kelliher MA (2014) Cutting edge: RIPK1 Kinase inactive mice are viable and protected from TNF-induced necroptosis in vivo. *J Immunol* 193(4):1539–1543. doi:10.4049/jimmunol.1400590
85. Rickard JA, O'Donnell JA, Evans JM, Lalaoui N, Poh AR, Rogers T, Vince JE, Lawlor KE, Ninnis RL, Anderton H, Hall C, Spall SK, Pesses TJ, Abud HE, Cengia LH, Corbin J, Mifsud S, Di Rago L, Metcalf D, Ernst M, Dewson G, Roberts AW, Alexander WS, Murphy JM, Ekert PG, Masters SL, Vaux DL, Croker BA, Gerlic M, Silke J (2014) RIPK1 regulates RIPK3-MLKL-driven systemic inflammation and emergency hematopoiesis. *Cell* 157(5):1175–1188. doi:10.1016/j.cell.2014.04.019
86. Kaiser WJ, Upton JW, Mocarski ES (2013) Viral modulation of programmed necrosis. *Curr Opin Virol* 3(3):296–306. doi:10.1016/j.coviro.2013.05.019
87. Silke J, Rickard JA, Gerlic M (2015) The diverse role of RIP kinases in necroptosis and inflammation. *Nat Immunol* 16(7):689–697. doi:10.1038/ni.3206
88. Khan N, Lawlor KE, Murphy JM, Vince JE (2014) More to life than death: molecular determinants of necroptotic and non-necroptotic RIP3 kinase signaling. *Curr Opin Immunol* 26:76–89. doi:10.1016/j.coi.2013.10.017
89. Mandal P, Berger SB, Pillay S, Moriwaki K, Huang C, Guo H, Lich JD, Finger J, Kasparcova V, Votta B, Ouellette M, King BW, Wisnoski D, Lakdawala AS, DeMartino MP, Casillas LN, Haile PA, Sehon CA, Marquis RW, Upton J, Daley-Bauer LP, Roback L, Ramia N, Dovey CM, Carette JE, Chan FK, Bertin J, Gough PJ, Mocarski ES, Kaiser WJ (2014) RIP3 induces apoptosis independent of pronecrotic kinase activity. *Mol Cell* 56(4):481–495. doi:10.1016/j.molcel.2014.10.021
90. Zhang J, Chan FK (2014) Cell biology. RIPK3 takes another deadly turn. *Science* 343(6177):1322–1323. doi:10.1126/science.1252526
91. Moujalled DM, Cook WD, Okamoto T, Murphy J, Lawlor KE, Vince JE, Vaux DL (2013) TNF can activate RIPK3 and cause programmed necrosis in the absence of RIPK1. *Cell Death Dis* 4:e465. doi:10.1038/cddis.2012.201
92. Remijsen Q, Goossens V, Grootjans S, Van den Haute C, Vanlangenakker N, Dondelinger Y, Roelandt R, Bruggeman I, Goncalves A, Bertrand MJ, Baekelandt V, Takahashi N, Berghe TV, Vandenabeele P (2014) Depletion of RIPK3 or MLKL blocks TNF-driven necroptosis and switches towards a delayed RIPK1 kinase-dependent apoptosis. *Cell Death Dis* 5:e1004. doi:10.1038/cddis.2013.531
93. Vince JE, Wong WW, Gentle I, Lawlor KE, Allam R, O'Reilly L, Mason K, Gross O, Ma S, Guarda G, Anderton H, Castillo R, Hacker G, Silke J, Tschopp J (2012) Inhibitor of apoptosis proteins limit RIP3 kinase-dependent interleukin-1 activation. *Immunity* 36(2):215–227. doi:10.1016/j.immuni.2012.01.012
94. Kang TB, Yang SH, Toth B, Kovalenko A, Wallach D (2013) Caspase-8 blocks kinase RIPK3-mediated activation of the NLRP3 inflammasome. *Immunity* 38(1):27–40. doi:10.1016/j.immuni.2012.09.015
95. Weng D, Marty-Roix R, Ganesan S, Proulx MK, Vladimer GI, Kaiser WJ, Mocarski ES, Pouliot K, Chan FK, Kelliher MA, Harris PA, Bertin J, Gough PJ, Shayakhmetov DM, Goguen JD, Fitzgerald KA, Silverman N, Lien E (2014) Caspase-8 and RIP kinases regulate bacteria-induced innate immune responses and cell death. *Proc Natl Acad Sci USA* 111(20):7391–7396. doi:10.1073/pnas.1403477111
96. Man SM, Tourlomousis P, Hopkins L, Monie TP, Fitzgerald KA, Bryant CE (2013) Salmonella infection induces recruitment of Caspase-8 to the inflammasome to modulate IL-1 β production. *J Immunol* 191(10):5239–5246. doi:10.4049/jimmunol.1301581
97. Gurung P, Anand PK, Malireddi RK, Vande Walle L, Van Opdenbosch N, Dillon CP, Weinlich R, Green DR, Lamkanfi M, Kanneganti TD (2014) FADD and caspase-8 mediate priming and activation of the canonical and noncanonical Nlrp3 inflammasomes. *J Immunol* 192(4):1835–1846. doi:10.4049/jimmunol.1302839
98. Philip NH, Dillon CP, Snyder AG, Fitzgerald P, Wynosky-Dolfi MA, Zwack EE, Hu B, Fitzgerald L, Mauldin EA, Copenhaver AM, Shin S, Wei L, Parker M, Zhang J, Oberst A, Green DR, Brodsky IE (2014) Caspase-8 mediates caspase-1 processing and innate immune defense in response to bacterial blockade of NF- κ B and MAPK signaling. *Proc Natl Acad Sci USA* 111(20):7385–7390. doi:10.1073/pnas.1403252111
99. Linkermann A, Skouta R, Himmerkus N, Mulay SR, Dewitz C, De Zen F, Prokai A, Zuchtriegel G, Krombach F, Welz PS, Weinlich R, Vanden Berghe T, Vandenabeele P, Pasparakis M, Bleich M, Weinberg JM, Reichel CA, Brasen JH, Kunzendorf U, Anders HJ, Stockwell BR, Green DR, Krautwald S (2014) Synchronized renal tubular cell death involves ferroptosis. *Proc Natl Acad Sci USA* 111(47):16836–16841. doi:10.1073/pnas.1415518111
100. Linkermann A, Brasen JH, Darding M, Jin MK, Sanz AB, Heller JO, De Zen F, Weinlich R, Ortiz A, Walczak H, Weinberg JM, Green DR, Kunzendorf U, Krautwald S (2013) Two independent pathways of regulated necrosis mediate ischemia-reperfusion injury. *Proc Natl Acad Sci USA* 110(29):12024–12029. doi:10.1073/pnas.1305538110
101. Zhao H, Ning J, Lemaire A, Koumpa FS, Sun JJ, Fung A, Gu J, Yi B, Lu K, Ma D (2015) Necroptosis and parthanatos are involved in remote lung injury after receiving ischemic renal allografts in rats. *Kidney Int* 87(4):738–748. doi:10.1038/ki.2014.388
102. Trichonas G, Murakami Y, Thanos A, Morizane Y, Kayama M, Deboucq CM, Hisatomi T, Miller JW, Vavvas DG (2010) Receptor interacting protein kinases mediate retinal detachment-induced photoreceptor necrosis and compensate for inhibition of apoptosis. *Proc Natl Acad Sci USA* 107(50):21695–21700. doi:10.1073/pnas.1009179107

103. Duprez L, Takahashi N, Van Hauwermeiren F, Vandendriessche B, Goossens V, Vanden Berghe T, Declercq W, Libert C, Cauwels A, Vandanaabeele P (2011) RIP kinase-dependent necrosis drives lethal systemic inflammatory response syndrome. *Immunity* 35(6):908–918. doi:10.1016/j.immuni.2011.09.020
104. Robinson N, McComb S, Mulligan R, Dudani R, Krishnan L, Sad S (2012) Type I interferon induces necroptosis in macrophages during infection with *Salmonella enterica* serovar Typhimurium. *Nat Immunol* 13(10):954–962. doi:10.1038/ni.2397
105. Moulin M, Anderton H, Voss AK, Thomas T, Wong WW, Bankovacki A, Feltham R, Chau D, Cook WD, Silke J, Vaux DL (2012) IAPs limit activation of RIP kinases by TNF receptor 1 during development. *EMBO J* 31(7):1679–1691. doi:10.1038/emboj.2012.18
106. Liu S, Wang X, Li Y, Xu L, Yu X, Ge L, Li J, Zhu Y, He S (2014) Necroptosis mediates TNF-induced toxicity of hippocampal neurons. *Biomed Res Int* 2014:290182. doi:10.1155/2014/290182
107. Kitur K, Parker D, Nieto P, Ahn DS, Cohen TS, Chung S, Wachtel S, Bueno S, Prince A (2015) Toxin-induced necroptosis is a major mechanism of *Staphylococcus aureus* lung damage. *PLoS Pathog* 11(4):e1004820. doi:10.1371/journal.ppat.1004820
108. Godwin A, Sharma A, Yang WL, Wang Z, Nicastrò J, Coppa GF, Wang P (2015) Receptor-interacting protein kinase 3 deficiency delays cutaneous wound healing. *PLoS One* 10(10):e0140514. doi:10.1371/journal.pone.0140514
109. Roychowdhury S, McMullen MR, Pisano SG, Liu X, Nagy LE (2013) Absence of receptor interacting protein kinase 3 prevents ethanol-induced liver injury. *Hepatology* 57(5):1773–1783. doi:10.1002/hep.26200
110. Meng L, Jin W, Wang X (2015) RIP3-mediated necrotic cell death accelerates systematic inflammation and mortality. *Proc Natl Acad Sci USA* 112(35):11007–11012. doi:10.1073/pnas.1514730112
111. Lin J, Li H, Yang M, Ren J, Huang Z, Han F, Huang J, Ma J, Zhang D, Zhang Z, Wu J, Huang D, Qiao M, Jin G, Wu Q, Huang Y, Du J, Han J (2013) A role of RIP3-mediated macrophage necrosis in atherosclerosis development. *Cell Rep* 3(1):200–210. doi:10.1016/j.celrep.2012.12.012
112. Cuda CM, Misharin AV, Khare S, Saber R, Tsai F, Archer AM, Homan PJ, Haines GK 3rd, Hutcheson J, Dorfleutner A, Budinger GR, Stehlik C, Perlman H (2015) Conditional deletion of caspase-8 in macrophages alters macrophage activation in a RIPK-dependent manner. *Arthr Res Ther* 17:291. doi:10.1186/s13075-015-0794-z
113. Murakami Y, Matsumoto H, Roh M, Suzuki J, Hisatomi T, Ikeda Y, Miller JW, Vavvas DG (2012) Receptor interacting protein kinase mediates necrotic cone but not rod cell death in a mouse model of inherited degeneration. *Proc Natl Acad Sci USA* 109(36):14598–14603. doi:10.1073/pnas.1206937109
114. Murakami Y, Matsumoto H, Roh M, Giani A, Kataoka K, Morizane Y, Kayama M, Thanos A, Nakatake S, Notomi S, Hisatomi T, Ikeda Y, Ishibashi T, Connor KM, Miller JW, Vavvas DG (2014) Programmed necrosis, not apoptosis, is a key mediator of cell loss and DAMP-mediated inflammation in dsRNA-induced retinal degeneration. *Cell Death Differ* 21(2):270–277. doi:10.1038/cdd.2013.109
115. Luedde M, Lutz M, Carter N, Sosna J, Jacoby C, Vucur M, Gautheron J, Roderburg C, Borg N, Reisinger F, Hippe HJ, Linkermann A, Wolf MJ, Rose-John S, Lullmann-Rauch R, Adam D, Fogel U, Heikenwalder M, Luedde T, Frey N (2014) RIP3, a kinase promoting necroptotic cell death, mediates adverse remodelling after myocardial infarction. *Cardiovasc Res* 103(2):206–216. doi:10.1093/cvr/cvu146
116. Gautheron J, Vucur M, Reisinger F, Cardenas DV, Roderburg C, Koppe C, Kreggenwinkel K, Schneider AT, Bartneck M, Neumann UP, Canbay A, Reeves HL, Luedde M, Tacke F, Trautwein C, Heikenwalder M, Luedde T (2014) A positive feedback loop between RIP3 and JNK controls non-alcoholic steatohepatitis. *EMBO Mol Med* 6(8):1062–1074. doi:10.15252/emmm.201403856
117. Petersen SL, Chen TT, Lawrence DA, Marsters SA, Gonzalez F, Ashkenazi A (2015) TRAF2 is a biologically important necroptosis suppressor. *Cell Death Differ* 22(11):1846–1857. doi:10.1038/cdd.2015.35
118. Vitner EB, Salomon R, Farfel-Becker T, Meshcheriakova A, Ali M, Klein AD, Platt FM, Cox TM, Futerman AH (2014) RIPK3 as a potential therapeutic target for Gaucher's disease. *Nat Med* 20(2):204–208. doi:10.1038/nm.3449
119. Chen W, Wu J, Li L, Zhang Z, Ren J, Liang Y, Chen F, Yang C, Zhou Z, Su SS, Zheng X, Zhang Z, Zhong CQ, Wan H, Xiao M, Lin X, Feng XH, Han J (2015) Ppm1b negatively regulates necroptosis through dephosphorylating Rip3. *Nat Cell Biol* 17(4):434–444. doi:10.1038/ncb3120
120. Xu Y, Ma H, Shao J, Wu J, Zhou L, Zhang Z, Wang Y, Huang Z, Ren J, Liu S, Chen X, Han J (2015) A role for tubular necroptosis in cisplatin-induced AKI. *J Am Soc Nephrol* 26(11):2647–2658. doi:10.1681/ASN.2014080741
121. Dara L, Johnson H, Suda J, Win S, Gaarde W, Han D, Kaplowitz N (2015) Receptor interacting protein kinase 1 mediates murine acetaminophen toxicity independent of the necrosome and not through necroptosis. *Hepatology* 62(6):1847–1857. doi:10.1002/hep.27939
122. Yang X, Chao X, Wang ZT, Ding WX (2015) The end of RIPK1-RIPK3-MLKL-mediated necroptosis in acetaminophen-induced hepatotoxicity? *Hepatology*. doi:10.1002/hep.28263
123. Zhao Y, Scott NA, Fynch S, Elkerbout L, Wong WW, Mason KD, Strasser A, Huang DC, Kay TW, Thomas HE (2015) Autoreactive T cells induce necrosis and not BCL-2-regulated or death receptor-mediated apoptosis or RIPK3-dependent necroptosis of transplanted islets in a mouse model of type 1 diabetes. *Diabetologia* 58(1):140
124. Rickard JA, Anderton H, Etemadi N, Nachbur U, Darding M, Peltzer N, Lalaoui N, Lawlor KE, Vanyai H, Hall C, Bankovacki A, Gangoda L, Wong WW, Corbin J, Huang C, Mocarski ES, Murphy JM, Alexander WS, Voss AK, Vaux DL, Kaiser WJ, Walczak H, Silke J (2014) TNFR1-dependent cell death drives inflammation in Sharpin-deficient mice. *Elife* 3. doi:10.7554/eLife.03464
125. Zhao J, Jitkaew S, Cai Z, Choksi S, Li Q, Luo J, Liu ZG (2012) Mixed lineage kinase domain-like is a key receptor interacting protein 3 downstream component of TNF-induced necrosis. *Proc Natl Acad Sci USA* 109(14):5322–5327. doi:10.1073/pnas.1200012109
126. Wu X, Tian L, Li J, Zhang Y, Han V, Li Y, Xu X, Li H, Chen X, Chen J, Jin W, Xie Y, Han J, Zhong CQ (2012) Investigation of receptor interacting protein (RIP3)-dependent protein phosphorylation by quantitative phosphoproteomics. *Mol Cell Proteomics* 11(12):1640–1651. doi:10.1074/mcp.M112.019091
127. Zhong CQ, Li Y, Yang D, Zhang N, Xu X, Wu Y, Chen J, Han J (2014) Quantitative phosphoproteomic analysis of RIP3-dependent protein phosphorylation in the course of TNF-induced necroptosis. *Proteomics* 14(6):713–724. doi:10.1002/pmic.201300326
128. Hildebrand JM, Tanzer MC, Lucet IS, Young SN, Spall SK, Sharma P, Pierotti C, Garnier JM, Dobson RC, Webb AI, Tripathydonis A, Babon JJ, Mulcair MD, Scanlon MJ, Alexander WS, Wilks AF, Czabotar PE, Lessene G, Murphy JM, Silke J (2014) Activation of the pseudokinase MLKL unleashes the four-helix bundle domain to induce membrane localization and necroptotic cell death. *Proc Natl Acad Sci USA* 111(42):15072–15077. doi:10.1073/pnas.1408987111

129. Kearney CJ, Cullen SP, Clancy D, Martin SJ (2014) RIPK1 can function as an inhibitor rather than an initiator of RIPK3-dependent necroptosis. *FEBS J* 281(21):4921–4934. doi:10.1111/febs.13034
130. Li D, Xu T, Cao Y, Wang H, Li L, Chen S, Wang X, Shen Z (2015) A cytosolic heat shock protein 90 and cochaperone CDC37 complex is required for RIP3 activation during necroptosis. *Proc Natl Acad Sci USA* 112(16):5017–5022. doi:10.1073/pnas.1505244112
131. Yoon S, Bogdanov K, Kovalenko A, Wallach D (2015) Necroptosis is preceded by nuclear translocation of the signaling proteins that induce it. *Cell Death Differ*. doi:10.1038/cdd.2015.92
132. Schenk B, Fulda S (2015) Reactive oxygen species regulate Smac mimetic/TNF α -induced necroptotic signaling and cell death. *Oncogene*. doi:10.1038/onc.2015.35
133. Fauster A, Rebsamen M, Huber KV, Bigenzahn JW, Stukalov A, Lardeau CH, Scorzoni S, Bruckner M, Gridling M, Parapatics K, Colinge J, Bennett KL, Kubicek S, Krautwald S, Linkermann A, Superti-Furga G (2015) A cellular screen identifies ponatinib and pazopanib as inhibitors of necroptosis. *Cell Death Dis* 6:e1767. doi:10.1038/cddis.2015.130
134. Najjar M, Suebsuwong C, Ray SS, Thapa RJ, Maki JL, Nogusa S, Shah S, Saleh D, Gough PJ, Bertin J, Yuan J, Balachandran S, Cuny GD, Degterev A (2015) Structure guided design of potent and selective ponatinib-based hybrid inhibitors for RIPK1. *Cell Rep* 10(11):1850–1860. doi:10.1016/j.celrep.2015.02.052
135. Xu M, Cai C, Sun X, Chen W, Li Q, Zhou H (2015) Clnk plays a role in TNF- α -induced cell death in murine fibrosarcoma cell line L929. *Biochem Biophys Res Commun* 463(3):275–279. doi:10.1016/j.bbrc.2015.05.046
136. Preyat N, Rossi M, Kers J, Chen L, Bertin J, Gough PJ, Le Moine A, Rongvaux A, Van Gool F, Leo O (2015) Intracellular nicotinamide adenine dinucleotide promotes TNF-induced necroptosis in a sirtuin-dependent manner. *Cell Death Differ*. doi:10.1038/cdd.2015.60
137. Karch J, Kanisicak O, Brody MJ, Sargent MA, Michael DM, Molkentin JD (2015) Necroptosis interfaces with MOMP and the MPTP in mediating cell death. *PLoS One* 10(6):e0130520. doi:10.1371/journal.pone.0130520
138. Rodriguez DA, Weinlich R, Brown S, Guy C, Fitzgerald P, Dillon CP, Oberst A, Quarato G, Low J, Cripps JG, Chen T, Green DR (2015) Characterization of RIPK3-mediated phosphorylation of the activation loop of MLKL during necroptosis. *Cell Death Differ*. doi:10.1038/cdd.2015.70
139. Lawlor KE, Khan N, Mildenhall A, Gerlic M, Croker BA, D'Cruz AA, Hall C, Kaur Spall S, Anderton H, Masters SL, Rashidi M, Wicks IP, Alexander WS, Mitsuuchi Y, Benetatos CA, Condon SM, Wong WW, Silke J, Vaux DL, Vince JE (2015) RIPK3 promotes cell death and NLRP3 inflammasome activation in the absence of MLKL. *Nat Commun* 6:6282. doi:10.1038/ncomms7282
140. Kim SJ, Li J (2013) Caspase blockade induces RIP3-mediated programmed necrosis in Toll-like receptor-activated microglia. *Cell Death Dis* 4:e716. doi:10.1038/cddis.2013.238
141. Kang S, Fernandes-Alnemri T, Rogers C, Mayes L, Wang Y, Dillon C, Roback L, Kaiser W, Oberst A, Sagar J, Fitzgerald KA, Green DR, Zhang J, Mocarski ES, Alnemri ES (2015) Caspase-8 scaffolding function and MLKL regulate NLRP3 inflammasome activation downstream of TLR3. *Nat Commun* 6:7515. doi:10.1038/ncomms8515
142. Kang YJ, Bang BR, Han KH, Hong L, Shim EJ, Ma J, Lerner RA, Otsuka M (2015) Regulation of NKT cell-mediated immune responses to tumours and liver inflammation by mitochondrial PGAM5-Drp1 signalling. *Nat Commun* 6:8371. doi:10.1038/ncomms9371
143. Moujalled DM, Cook WD, Murphy JM, Vaux DL (2014) Necroptosis induced by RIPK3 requires MLKL but not Drp1. *Cell Death Dis* 5:e1086. doi:10.1038/cddis.2014.18
144. Cook WD, Moujalled DM, Ralph TJ, Lock P, Young SN, Murphy JM, Vaux DL (2014) RIPK1- and RIPK3-induced cell death mode is determined by target availability. *Cell Death Differ* 21(10):1600–1612. doi:10.1038/cdd.2014.70
145. Wu XN, Yang ZH, Wang XK, Zhang Y, Wan H, Song Y, Chen X, Shao J, Han J (2014) Distinct roles of RIP1-RIP3 hetero- and RIP3-RIP3 homo-interaction in mediating necroptosis. *Cell Death Differ* 21(11):1709–1720. doi:10.1038/cdd.2014.77
146. Orozco S, Yatim N, Werner MR, Tran H, Gunja SY, Tait SW, Albert ML, Green DR, Oberst A (2014) RIPK1 both positively and negatively regulates RIPK3 oligomerization and necroptosis. *Cell Death Differ* 21(10):1511–1521. doi:10.1038/cdd.2014.76
147. Qiu X, Klausen C, Cheng JC, Leung PC (2015) CD40 ligand induces RIP1-dependent, necroptosis-like cell death in low-grade serous but not serous borderline ovarian tumor cells. *Cell Death Dis* 6:e1864. doi:10.1038/cddis.2015.229
148. Steinwascher S, Nuges AL, Schoeneberger H, Fulda S (2015) Identification of a novel synergistic induction of cell death by Smac mimetic and HDAC inhibitors in acute myeloid leukemia cells. *Cancer Lett* 366(1):32–43. doi:10.1016/j.canlet.2015.05.020
149. Miki Y, Akimoto J, Moritake K, Hironaka C, Fujiwara Y (2015) Photodynamic therapy using talaporfin sodium induces concentration-dependent programmed necroptosis in human glioblastoma T98G cells. *Lasers Med Sci* 30(6):1739–1745. doi:10.1007/s10103-015-1783-9
150. Koo MJ, Rooney KT, Choi ME, Ryter SW, Choi AM, Moon JS (2015) Impaired oxidative phosphorylation regulates necroptosis in human lung epithelial cells. *Biochem Biophys Res Commun* 464(3):875–880. doi:10.1016/j.bbrc.2015.07.054
151. Mizumura K, Cloonan SM, Nakahira K, Bhashyam AR, Cervo M, Kitada T, Glass K, Owen CA, Mahmood A, Washko GR, Hashimoto S, Ryter SW, Choi AM (2014) Mitophagy-dependent necroptosis contributes to the pathogenesis of COPD. *J Clin Invest* 124(9):3987–4003. doi:10.1172/JCI74985
152. Fu Z, Deng B, Liao Y, Shan L, Yin F, Wang Z, Zeng H, Zuo D, Hua Y, Cai Z (2013) The anti-tumor effect of shikonin on osteosarcoma by inducing RIP1 and RIP3 dependent necroptosis. *BMC Cancer* 13:580. doi:10.1186/1471-2407-13-580
153. Dunai ZA, Imre G, Barna G, Korcsmaros T, Petak I, Bauer PI, Mihalik R (2012) Staurosporine induces necroptotic cell death under caspase-compromised conditions in U937 cells. *PLoS One* 7(7). doi:10.1371/journal.pone.0041945
154. Chromik J, Safferthal C, Serve H, Fulda S (2014) Smac mimetic primes apoptosis-resistant acute myeloid leukaemia cells for cytarabine-induced cell death by triggering necroptosis. *Cancer Lett* 344(1):101–109. doi:10.1016/j.canlet.2013.10.018
155. Saveljeva S, Mc Laughlin SL, Vandenabeele P, Samali A, Bertrand MJ (2015) Endoplasmic reticulum stress induces ligand-independent TNFR1-mediated necroptosis in L929 cells. *Cell Death Disease* 6. doi:10.1038/cddis.2014.548
156. Melo-Lima S, Celeste Lopes M, Mollinedo F (2014) Necroptosis is associated with low procaspase-8 and active RIPK1 and -3 in human glioma cells. *Oncoscience* 1(10):649–664
157. Ratovitski EA (2015) Phospho- Δ Np63 α -responsive microRNAs contribute to the regulation of necroptosis in squamous cell carcinoma upon cisplatin exposure. *FEBS Lett* 589(12):1352–1358. doi:10.1016/j.febslet.2015.04.020
158. Koo GB, Morgan MJ, Lee DG, Kim WJ, Yoon JH, Koo JS, Kim SI, Kim SJ, Son MK, Hong SS, Levy JM, Pollyea DA, Jordan CT, Yan P, Frankhouser D, Nicolet D, Maharry K, Marcucci G,

- Choi KS, Cho H, Thorburn A, Kim YS (2015) Methylation-dependent loss of RIP3 expression in cancer represses programmed necrosis in response to chemotherapeutics. *Cell Res* 25(6):707–725. doi:[10.1038/cr.2015.56](https://doi.org/10.1038/cr.2015.56)
159. Philipp S, Sosna J, Plenge J, Kalthoff H, Adam D (2015) Homoharringtonine, a clinically approved anti-leukemia drug, sensitizes tumor cells for TRAIL-induced necroptosis. *Cell Commun Signal* 13:25. doi:[10.1186/s12964-015-0103-0](https://doi.org/10.1186/s12964-015-0103-0)
 160. Blohberger J, Kunz L, Einwang D, Berg U, Berg D, Ojeda SR, Dissen GA, Frohlich T, Arnold GJ, Soreq H, Lara H, Mayerhofer A (2015) Readthrough acetylcholinesterase (AChE-R) and regulated necrosis: pharmacological targets for the regulation of ovarian functions? *Cell Death Dis* 6:e1685. doi:[10.1038/cddis.2015.51](https://doi.org/10.1038/cddis.2015.51)
 161. Desai J, Vr SK, Mulay SR, Konrad L, Romoli S, Schauer C, Herrmann M, Bilyy R, Muller S, Popper B, Nakazawa D, Weidenbusch M, Thomasova D, Krautwald S, Linkermann A, Anders HJ (2015) Neutrophil extracellular trap formation can involve RIPK1-RIPK3-MLKL signalling. *Eur J Immunol*. doi:[10.1002/eji.201545605](https://doi.org/10.1002/eji.201545605)
 162. Amini P, Stojkov D, Wang X, Wicki S, Kaufmann T, Wong WW, Simon H-UU, Yousefi S (2015) NET formation can occur independently of RIPK3 and MLKL signaling. *Eur J Immunol*. doi:[10.1002/eji.201545615](https://doi.org/10.1002/eji.201545615)
 163. Suzuki T, Kikuguchi C, Sharma S, Sasaki T, Tokumasu M, Adachi S, Natsume T, Kanegae Y, Yamamoto T (2015) CNOT3 suppression promotes necroptosis by stabilizing mRNAs for cell death-inducing proteins. *Sci Rep* 5:14779. doi:[10.1038/srep14779](https://doi.org/10.1038/srep14779)
 164. Yu X, Deng Q, Li W, Xiao L, Luo X, Liu X, Yang L, Peng S, Ding Z, Feng T, Zhou J, Fan J, Bode AM, Dong Z, Liu J, Cao Y (2015) Neolbalconol induces cell death through necroptosis by regulating RIPK-dependent autocrine TNF α and ROS production. *Oncotarget* 6(4):1995–2008
 165. He W, Wang Q, Srinivasan B, Xu J, Padilla MT, Li Z, Wang X, Liu Y, Gou X, Shen HM, Xing C, Lin Y (2014) A JNK-mediated autophagy pathway that triggers c-IAP degradation and necroptosis for anticancer chemotherapy. *Oncogene* 33(23):3004–3013. doi:[10.1038/onc.2013.256](https://doi.org/10.1038/onc.2013.256)
 166. Huang CY, Kuo WT, Huang YC, Lee TC, Yu LC (2013) Resistance to hypoxia-induced necroptosis is conferred by glycolytic pyruvate scavenging of mitochondrial superoxide in colorectal cancer cells. *Cell Death Dis* 4:e622. doi:[10.1038/cddis.2013.149](https://doi.org/10.1038/cddis.2013.149)
 167. Tian W, Xu D, Han W, He H, Cai H, Chen H, Zhou M, Chen J, Deng YC (2013) Cyclophilin D modulates cell death transition from early apoptosis to programmed necrosis induced by honokiol. *Int J Oncol* 42(5):1654–1663. doi:[10.3892/ijo.2013.1863](https://doi.org/10.3892/ijo.2013.1863)
 168. Wang XQ, Jiang W, Yan YQ, Gong T, Han JH, Tian ZG, Zhou RB (2014) RNA viruses promote activation of the NLRP3 inflammasome through a RIP1-RIP3-DRP1 signaling pathway. *Nat Immunol* 15(12):1126–1133. doi:[10.1038/ni.3015](https://doi.org/10.1038/ni.3015)
 169. Omoto S, Guo H, Talekar GR, Roback L, Kaiser WJ, Mocarski ES (2015) Suppression of RIP3-dependent necroptosis by human cytomegalovirus. *J Biol Chem* 290(18):11635–11648. doi:[10.1074/jbc.M115.646042](https://doi.org/10.1074/jbc.M115.646042)
 170. Wang X, Li Y, Liu S, Yu X, Li L, Shi C, He W, Li J, Xu L, Hu Z, Yu L, Yang Z, Chen Q, Ge L, Zhang Z, Zhou B, Jiang X, Chen S, He S (2014) Direct activation of RIP3/MLKL-dependent necrosis by herpes simplex virus 1 (HSV-1) protein ICP6 triggers host antiviral defense. *Proc Natl Acad Sci USA* 111(43):15438–15443. doi:[10.1073/pnas.1412767111](https://doi.org/10.1073/pnas.1412767111)
 171. Yu X, Li Y, Chen Q, Su C, Zhang Z, Yang C, Hu Z, Hou J, Zhou J, Gong L, Jiang X, Zheng C, He S (2015) Herpes simplex virus type 1 (HSV-1) and HSV-2 mediate species specific modulations of programmed necrosis through the viral ribonucleotide reductase large subunit R1. *J Virol*. doi:[10.1128/JVI.02446-15](https://doi.org/10.1128/JVI.02446-15)
 172. Guo H, Omoto S, Harris PA, Finger JN, Bertin J, Gough PJ, Kaiser WJ, Mocarski ES (2015) Herpes simplex virus suppresses necroptosis in human cells. *Cell Host Microbe* 17(2):243–251. doi:[10.1016/j.chom.2015.01.003](https://doi.org/10.1016/j.chom.2015.01.003)
 173. Huang Z, Wu SQ, Liang Y, Zhou X, Chen W, Li L, Wu J, Zhuang Q, Ca Chen, Li J, Zhong CQQ, Xia W, Zhou R, Zheng C, Han J (2015) RIP1/RIP3 binding to HSV-1 ICP6 initiates necroptosis to restrict virus propagation in mice. *Cell Host Microbe* 17(2):229–242. doi:[10.1016/j.chom.2015.01.002](https://doi.org/10.1016/j.chom.2015.01.002)
 174. Wagner RN, Reed JC, Chanda SK (2015) HIV-1 protease cleaves the serine-threonine kinases RIPK1 and RIPK2. *Retrovirology* 12:74. doi:[10.1186/s12977-015-0200-6](https://doi.org/10.1186/s12977-015-0200-6)
 175. Lim SY, Davidson SM, Mocanu MM, Yellon DM, Smith CC (2007) The cardioprotective effect of necrostatin requires the cyclophilin-D component of the mitochondrial permeability transition pore. *Cardiovasc Drugs Ther* 21(6):467–469. doi:[10.1007/s10557-007-6067-6](https://doi.org/10.1007/s10557-007-6067-6)
 176. You Z, Savitz SI, Yang J, Degterev A, Yuan J, Cuny GD, Moskowitz MA, Whalen MJ (2008) Necrostatin-1 reduces histopathology and improves functional outcome after controlled cortical impact in mice. *J Cereb Blood Flow Metab* 28(9):1564–1573. doi:[10.1038/jcbfm.2008.44](https://doi.org/10.1038/jcbfm.2008.44)
 177. Rosenbaum DM, Degterev A, David J, Rosenbaum PS, Roth S, Grotta JC, Cuny GD, Yuan J, Savitz SI (2010) Necroptosis, a novel form of caspase-independent cell death, contributes to neuronal damage in a retinal ischemia-reperfusion injury model. *J Neurosci Res* 88(7):1569–1576. doi:[10.1002/jnr.22314](https://doi.org/10.1002/jnr.22314)
 178. Wang Y, Wang H, Tao Y, Zhang S, Wang J, Feng X (2014) Necroptosis inhibitor necrostatin-1 promotes cell protection and physiological function in traumatic spinal cord injury. *Neuroscience* 266:91–101. doi:[10.1016/j.neuroscience.2014.02.007](https://doi.org/10.1016/j.neuroscience.2014.02.007)
 179. Zhu Y, Cui H, Gan H, Xia Y, Wang L, Wang Y, Sun Y (2015) Necroptosis mediated by receptor interaction protein kinase 1 and 3 aggravates chronic kidney injury of subtotal nephrectomized rats. *Biochem Biophys Res Commun* 461(4):575–581. doi:[10.1016/j.bbrc.2015.03.164](https://doi.org/10.1016/j.bbrc.2015.03.164)
 180. Ofengeim D, Ito Y, Najafav A, Zhang Y, Shan B, DeWitt JP, Ye J, Zhang X, Chang A, Vakifahmetoglu-Norberg H, Geng J, Py B, Zhou W, Amin P, Berlink Lima J, Qi C, Yu Q, Trapp B, Yuan J (2015) Activation of necroptosis in multiple sclerosis. *Cell Rep* 10(11):1836–1849. doi:[10.1016/j.celrep.2015.02.051](https://doi.org/10.1016/j.celrep.2015.02.051)
 181. Re DB, Le Verche V, Yu C, Amoroso MW, Politi KA, Phani S, Ikiz B, Hoffmann L, Koolen M, Nagata T, Papadimitriou D, Nagy P, Mitsumoto H, Kariya S, Wichterle H, Henderson CE, Przedborski S (2014) Necroptosis drives motor neuron death in models of both sporadic and familial ALS. *Neuron* 81(5):1001–1008. doi:[10.1016/j.neuron.2014.01.011](https://doi.org/10.1016/j.neuron.2014.01.011)
 182. Li JX, Feng JM, Wang Y, Li XH, Chen XX, Su Y, Shen YY, Chen Y, Xiong B, Yang CH, Ding J, Miao ZH (2014) The B-Raf(V600E) inhibitor dabrafenib selectively inhibits RIP3 and alleviates acetaminophen-induced liver injury. *Cell Death Dis* 5:e1278. doi:[10.1038/cddis.2014.241](https://doi.org/10.1038/cddis.2014.241)
 183. Kim SK, Kim WJ, Yoon JH, Ji JH, Morgan MJ, Cho H, Kim YC, Kim YS (2015) Upregulated RIP3 expression potentiates MLKL phosphorylation-mediated programmed necrosis in toxic epidermal necrolysis. *J Invest Dermatol* 135(8):2021–2030. doi:[10.1038/jid.2015.90](https://doi.org/10.1038/jid.2015.90)
 184. He L, Peng K, Liu Y, Xiong J, Zhu FF (2013) Low expression of mixed lineage kinase domain-like protein is associated with poor prognosis in ovarian cancer patients. *Oncotargets Ther* 6:1539–1543. doi:[10.2147/OTT.S52805](https://doi.org/10.2147/OTT.S52805)
 185. Pierdomenico M, Negroni A, Stronati L, Vitali R, Prete E, Bertin J, Gough PJ, Aloisi M, Cucchiara S (2014) Necroptosis is

- active in children with inflammatory bowel disease and contributes to heighten intestinal inflammation. *Am J Gastroenterol* 109(2):279–287. doi:[10.1038/ajg.2013.403](https://doi.org/10.1038/ajg.2013.403)
186. Geserick P, Wang J, Schilling R, Horn S, Harris PA, Bertin J, Gough PJ, Feoktistova M, Leverkus M (2015) Absence of RIPK3 predicts necroptosis resistance in malignant melanoma. *Cell Death Dis* 6:e1884. doi:[10.1038/cddis.2015.240](https://doi.org/10.1038/cddis.2015.240)
187. Gaiha GD, McKim KJ, Woods M, Pertel T, Rohrbach J, Barteneva N, Chin CR, Liu D, Soghoian DZ, Cesa K, Wilton S, Waring MT, Chicoine A, Doering T, Wherry EJ, Kaufmann DE, Lichterfeld M, Brass AL, Walker BD (2014) Dysfunctional HIV-specific CD8+ T cell proliferation is associated with increased caspase-8 activity and mediated by necroptosis. *Immunity* 41(6):1001–1012. doi:[10.1016/j.immuni.2014.12.011](https://doi.org/10.1016/j.immuni.2014.12.011)
188. Nugues AL, El Bouazzati H, Hetuin D, Berthon C, Loyens A, Bertrand E, Jouy N, Idziorek T, Quesnel B (2014) RIP3 is downregulated in human myeloid leukemia cells and modulates apoptosis and caspase-mediated p65/RelA cleavage. *Cell Death Dis* 5:e1384. doi:[10.1038/cddis.2014.347](https://doi.org/10.1038/cddis.2014.347)