



ERRATUM

## Erratum to: Synthesis and evaluation of mefway analogs as ligands for serotonin 5HT<sub>1A</sub> receptors

Joanne P. Thio<sup>1</sup> · Christopher Liang<sup>1</sup> · Alisha K. Bajwa<sup>1</sup> ·  
Dustin W. Wooten<sup>2</sup> · Bradley T. Christian<sup>2</sup> · Jogeshwar Mukherjee<sup>1</sup>

Published online: 29 April 2015  
© Springer Science+Business Media New York 2015

### Erratum to: Med Chem Res (2015) 24:1480–1486 DOI 10.1007/s00044-014-1238-z

The authors would like to correct an error in Fig. 4 (cyclohexyl ketone shown instead of cyclohexyl ester, in structures **11**, **12**, **13**) of the original publication. The corrected Fig. 4 is published with this erratum.

---

The online version of the original article can be found under  
doi:[10.1007/s00044-014-1238-z](https://doi.org/10.1007/s00044-014-1238-z).

---

✉ Jogeshwar Mukherjee  
j.mukherjee@uci.edu

<sup>1</sup> Preclinical Imaging, B138 Medical Sciences, Department of Radiological Sciences, University of California, Irvine, CA 92697-5000, USA

<sup>2</sup> Department of Medical Physics, University of Wisconsin-Madison, Madison, WI 53705, USA

**Fig. 4** Synthesis scheme of Mefway. Starting with the conversion of the carboxylic acids (**11**) to the corresponding cyclohexanoyl chlorides (**12**), the amine WAY-100634 (**7**) was converted to the corresponding carboxamides (**13**). Reduction of the carboxylic acid esters with lithium aluminum hydride ( $\text{LiAlH}_4$ ) gave the corresponding alcohols (**14**) which were subsequently fluorinated with DAST to provide 4-Mefway (**9**) and 3-Mefway (**10**)

