



# Evaluating monitoring options for conservation: comparing traditional and environmental DNA tools for a critically endangered mammal

Chanjuan Qu<sup>1</sup> · Kathryn A. Stewart<sup>1,2</sup>

Received: 3 August 2018 / Revised: 6 December 2018 / Accepted: 30 January 2019 / Published online: 18 February 2019  
© The Author(s) 2019

## Abstract

While conservation management has made tremendous strides to date, deciding where, when and how to invest limited monitoring budgets is a central concern for impactful decision-making. New analytical tools, such as environmental DNA (eDNA), are now facilitating broader biodiversity monitoring at unprecedented scales, in part, due to time, and presumably cost, of methodological efficiency. Genetic approaches vary from conventional PCR (cPCR; species presence), to metabarcoding (community structure), and qPCR (relative DNA abundance, detection sensitivity). Knowing when to employ these techniques over traditional protocols could enable practitioners to make more informed choices concerning data collection. Using 12 species-specific primers designed for cPCR, eDNA analysis of the Yangtze finless porpoise (YFP; *Neophocaena asiaeorientalis asiaeorientalis*), a critically endangered aquatic mammal within the Yangtze River, we validated and optimized these primers for use in qPCR. We tested repeatability and sensitivity to detect YFP eDNA and subsequently compared the cost of traditional (visual and capture) sampling to eDNA tools. Our results suggest cPCR as the least expensive sampling option but the lack of PCR sensitivity suggests it may not be the most robust method for this taxon, predominately useful as a supplementary tool or with large expected populations. Alternatively, qPCR remained less expensive than traditional surveys, representing a highly repeatable and sensitive method for this behaviorally elusive species. Cost comparisons of surveying practices have scarcely been discussed; however, given budgetary constraints particularly for developing countries with limited local oversight but high endemism, we encourage managers to carefully consider the trade-offs among accuracy, cost, coverage, and speed for biodiversity monitoring.

**Keywords** Yangtze finless porpoise · Environmental DNA · Systematic conservation planning

## Introduction

As a discipline, systematic conservation planning (e.g., Margules and Pressey 2000) has now surpassed three decades, and its achievements around the world have been remarkable. Still, the

social and political environment in which conservation issues are addressed is highly complex, often forming a nexus between the cost, speed, and accuracy of collecting necessary biodiversity data. Indeed, so much of conservation fundamentally involves choices about *where*, *when* and *how* to make investments, thus requiring organismal information at scales commensurate with both goals and monetary availability. Therefore, there remains a vital need to continually highlight new analytical frameworks, innovations, and advances in our collective understanding and approaches to conservation's core objectives.

Despite the prerequisite for comprehensive monitoring initiatives in conservation programs, a thorough knowledge of organismal distribution and abundance is often prohibitive due largely to difficulties in data collection for hard-to-study taxa (e.g., cryptic, behaviourally elusive, low site fidelity, or rare), difficult to sample locales (e.g., aquatic environments), and affiliated costs, particularly in developing countries (Danielsen et al. 2003). Advancements in biodiversity data monitoring via DNA sampled straight from the environment

---

Communicated by: Matthias Waltert

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s00114-019-1605-1>) contains supplementary material, which is available to authorized users.

✉ Kathryn A. Stewart  
stewart.a.kat@gmail.com

<sup>1</sup> State Key Laboratory for Pollution Control and Resource Reuse, College of Environmental Science and Engineering, Tongji University, Shanghai, People's Republic of China

<sup>2</sup> Institute for Biodiversity and Ecosystem Dynamics, Department of Evolutionary and Population Biology, University of Amsterdam, Amsterdam, Netherlands

(without invasively targeting taxa) known as eDNA has recently revolutionized conservation biology.

Advocated as a time and cost-effective alternative to traditional methods of biodiversity data monitoring, eDNA is a highly sensitive technology that has been successfully employed for myriad species and goals (e.g., Ficetola et al. 2008; Lodge et al. 2012; Fukumoto et al. 2015; Ma et al. 2015; Dougherty et al. 2016). However, just as surveying methods are varied, so too are eDNA applications, and few studies to date have assessed and compared the cost of these protocols. Applications of eDNA methodology can range from detecting the presence/absence of species with conventional PCR (cPCR) (e.g., Jerde et al. 2011; Dejean et al. 2012; Thomsen et al. 2012; Mahon et al. 2013; Piaggio et al. 2014; Fukumoto et al. 2015) or community constituents via metabarcoding (e.g., Evans et al. 2015; Valentini et al. 2016; Hänfling et al. 2016; Shaw et al. 2016), to quantifying the relative abundance of DNA sequences (proxies for relative species abundance) or increasing detection sensitivity via quantitative PCR (qPCR) (e.g., Takahara et al. 2012; Goldberg et al. 2013; Pilliod et al. 2013; Klymus et al. 2015; Laramie et al. 2015; Balasingham et al. 2017). Still, it is difficult to discern when and how to match appropriate sampling protocols with conservation goals in the most cost-effective manner; this understanding could propel the entire field of conservation biology forward and galvanize eDNA as a tractable tool for managers.

At present, the Yangtze finless porpoise, *Neophocaena asiaeorientalis asiaeorientalis* (henceforth YFP), is the only cetacean species to be found within the Yangtze River and its tributaries (China), the world's only freshwater porpoise, and is currently classified as a critically endangered species by the IUCN (estimated to encompass less than 1050 individuals in the wild; Wan et al. 2016; Mei et al. 2012). The YFP is a far-ranging aquatic species distributed across the middle-to-lower Yangtze River, many lake systems, and an oxbow lake reserve hydrologically linked to the Yangtze River, totally approximately 1890 km in length (Li 2011). Although a conservation priority for China, YFP natural history make it difficult to survey: i) they remain behaviorally elusive, ii) are benthic feeders infrequently surfacing to breathe, and iii) their dark coloration and lack of dorsal fin to break the water surface makes visual detection difficult (Stewart et al. 2017). Thus, current traditional population monitoring for this taxon is primarily restricted to presence/absence information via visual or acoustic surveying unless individuals are physically captured for demographic information (although the invasiveness of the capture method remains a concern). YFP's extensive distribution across the Yangtze River and its tributaries also makes approaches to population management and data collection costly, an important consideration for this developing country.

Previous research has successfully developed 12 species-specific YFP primer pairs used in cPCR for eDNA sampling

(Ma et al. 2016). However, this eDNA approach only allows for the identification of presence/absence data. While more informative (e.g., abundance information for target taxa), qPCR is further suggested to be more sensitive for eDNA detection (Balasingham et al. 2017). Thus, for this study we optimize and validate species-specific primer pairs for qPCR amplification (quantifying relative DNA abundance), simultaneously testing and comparing the sensitivity (theoretical limits of detection) of each primer pair for eDNA applications. We additionally conduct a cost-assessment of YFP traditional visual monitoring to conventional and quantitative PCR data collections. We ultimately aim to generate management recommendations for conservation practitioners commensurate with the scale, goal, and financial feasibility of each sampling application currently available (i.e., compare different eDNA detection methods to that of traditional surveying).

## Methods

### Primer optimization and validation

Using 12 published YFP primers pairs validated for use in cPCR (Ma et al. 2016), we optimized qPCR protocols for use in eDNA surveying. Species-specificity was previously determined for cPCR *in silico* (bioinformatically) via the comparison of 20 complete YFP cytochrome *b* (*cyt b*) sequences (range of haplotypes) to other geographically overlapping species; *in vitro* (blood and tissue samples) from YFP and sister species, and *in situ* (within environments where YFP are known to be present, positive control, and where they are absent, negative control) (for details see Ma et al. 2016). Briefly for cPCR, DNA was extracted from both blood and environmental samples using a DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany) in a final elution volume of 200  $\mu$ L. DNA concentrations were measured with a NanoDrop Lite spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA) and proceeding reaction volumes were 25  $\mu$ L, consisting of 1  $\mu$ L of 1–20 ng/ $\mu$ L template DNA, 12.5  $\mu$ L of 1  $\times$  Taq PCR Master Mix (BIO BASIC CANADA INC.), 1  $\mu$ L each of 0.2  $\mu$ M each of forward and reverse primer, and 9.5  $\mu$ L of ddH<sub>2</sub>O. cPCR amplification included the following thermal cycling profile: 94  $^{\circ}$ C initial denaturation for 4 min, followed by 35 cycles of 94  $^{\circ}$ C for 30 s, 52  $^{\circ}$ C for 30 s, 72  $^{\circ}$ C for 20 s, and one final extension step at 72  $^{\circ}$ C for 5 min. cPCR products were visualized using electrophoresis on 2% agarose gel, stained with 7  $\mu$ L of 4S Red Plus Nucleic Acid Stain (BBI) per gel, and all amplifications were repeated in triplicate (following Ficetola et al. 2015). Finally, cPCR products of the target sequences were sent to Sangon Biotech (Shanghai, China) utilizing an ABI 3730XL sequencer to validate PCR products as indeed being from YFP. Sequences were visualized using MEGA6 (Tamura

et al. 2013) and then compared to all available sequences in GenBank using Primer-BLAST (Ye et al. 2012). For sensitivity analyses, concentration assays involved diluting the DNA extracted from blood samples with sterilized ddH<sub>2</sub>O to 4 ratios: 1:2, 1:10, 1:100, 1:1000, replicating PCR amplification in triplicate, and visualizing the electrophoresis results to note maximum detection sensitivity of each primer pair. Detection using cPCR was denoted as positive if at least 2 of the 3 replicates amplified YFP DNA. If only 1 out of 3 demonstrated positive amplification, we re-ran another triplicate test to eliminate possible contamination.

For optimization of qPCR, we first acquired a synthetic gene solution as template DNA. This synthetic gene was then used to optimize amplification and explore the sensitivity/limit of detection (LOD; the minimum level of target DNA detected in a sample) of these primer pairs. The synthetic *cyt b* gene sequence (1140 bp) was retrieved from GenBank (accession number KJ472902), and encompassed the entire region in which all 12 primer pairs are located for YFP (Ma et al. 2016). Each qPCR reaction contained 10 µL of SYBR Premix Ex Taq II (Tli RNaseH Plus) (2×) (TaKaRa.), 0.4 µL of ROX Reference Dye II (50×) (TaKaRa.), 6.4 µL of RNase-free water, 0.6 µL of forward primer (10 µM), 0.6 µL of reverse primer (10 µM), and 2 µL of template DNA, for a total volume of 20 µL. All amplifications were performed on an Applied Biosystems 7500 Real-Time PCR System (Life Technologies) with qPCR thermal profile as follows: holding stage (95 °C for 30s), followed by cycling stage (40 cycles of 95 °C for 5 s and 64 °C for 34 s), then followed to a melt-curve stage (95 °C for 15 s, 60 °C for 1 min, 95 °C for 30 s, and 60 °C for 15 s).

We judged qPCR amplification to be successful if the results met the following requirements: 1) qPCR standards (or samples) displayed the expected amplification curve; 2) amplifications displayed an expected melting curve peak; and 3) negative controls showed no amplification prior to critical threshold. qPCR reactions were performed in triplicate at concentrations of 500 and 5000 copies/µL due to high repeatability (near 100% positive amplification), but to increase our confidence in the amplification results at lower concentrations (5–50 copies/µL), qPCR reactions were tested in 13 technical replicates including negative controls (RNase-free water). Amplifications were recorded as zero detection when reactions did not cross the fluorescent threshold prior to the 40th cycle (Ct; 40), and quantified as a positive detection when samples demonstrated amplification prior to Ct.

Amplification efficiency (%) for qPCR reactions was additionally calculated for each primer pair based on the slope of the standard curve, assuming a slope of  $-3.322$  with an amplification factor of 2, which equates to 100% efficiency.

To test the sensitivity of each primer pair for use in eDNA analysis, we amplified the synthetic gene at concentrations of 5, 25, 50, 500, and 5000 copies/µL. The synthetic gene solution concentrations in ng/µL (as measured via NanoDrop Lite

spectrophotometer; Thermo Fisher Scientific, Wilmington, DE, USA), were converted from  $x$  ng/µL to  $y$  copies/µL using the standard formula:

$$y = \frac{x \times 10^{-9}}{650 \times (\text{length})} \times 6.02 \times 10^{23}$$

For this calculation, we assumed 650 Da the weight of each base pair and  $6.022 \times 10^{23}$  for the conversion to molecules/mol (Avogadro's number). For use in qPCR calculations, we used the combined synthetic gene length of 1140 bp and 2710 bp for the synthetic gene clone vector length (pUC57), and for PCR calculations we used the length of each primer amplicon (bp). Maximum cPCR sensitivity data was extracted from Ma et al. (2016) in which extracted DNA from YFP blood samples were diluted to a minimum level of detection (ng/µL converted to copies/µL).

### Validation via eDNA collection

To verify primer efficiency, we tested whether these primer pairs could amplify YFP eDNA from a location where we knew this species to be present. Thus for a positive control, we collected water from the “netted cage” (approximately a 15 m × 15 m enclosure) within Tian-e-Zhou Baiji National Nature Reserve (29° 51' 11 N, 112° 35' 15E), a natural protected area for the YFP which runs parallel, and is hydrologically linked to, the Yangtze River (similar in protocol to Foote et al. 2012), and acts as a lake due to limited water flow (Stewart et al. 2017). This reserve is hydrologically similar to other natural lake systems within which over half the YFP species populations are found in China (e.g., Dongting and Poyang lake). Within the netted cage area, 2 YFP individuals were temporarily housed for breeding research purposes before being released into the reserve. The water within the netted cage was connected, and freely flowed to the rest of the reserve which further housed approximately 58 other YFP individuals at the time of sampling (April 2015). Water was sampled 0.4 m below water surface and acquired with a VanDorn water sampler (GRASP CG-00, GRASP Science & Technology Co., Ltd.) which had been sterilized with 20% diluted bleach, rinsed with sterilized water, and then dried. The water sample was then filtered using a portable field peristaltic pump (Spectra Scientific Inc. Spectra Filled-Pro Professional Grade) with 47 mm diameter mixed cellulose esters (MCE) filter paper, 0.45 µm pore size, utilizing a sterilized, reusable filter holder (Sterifil 47 mm Filter Holder, MILLIPORE). To prevent clogging the filter paper with large suspended particles (e.g., algae), we wrapped the inlet of the filter holder with sterilized, disposable, medical-grade gauze. Based on collection procedures and sensitivity measures from cPCR protocols (Ma et al. 2016), we collected and filtered approximately 3–1 L samples of water, calculating

final extracted DNA concentration from each sample as a proportion of their original sample volume. After filtration, we folded the filter paper using sterilized tweezers and placed the filter paper into an Eppendorf tube with 95% ethanol. Samples were immediately stored in a  $-20^{\circ}\text{C}$  freezer until DNA extraction. For our negative control, we chose Gonghuwan ( $31^{\circ} 27' 21'' \text{N}$ ,  $120^{\circ} 20' \text{E}$ ), an artificial lake in Wuxi, China, where porpoises are absent and water supply is not linked with the Yangtze River or any other waterway containing YFP.

DNA was subsequently extracted from stored, dried filter paper with the PowerWater® DNA Isolation Kit (Mo Bio Laboratories, Inc) via placing filter paper into a 5-mL PowerWater® Bead Tube and following manufacturer's protocol. After DNA extraction, the mean total eDNA concentration was measured with a NanoDrop Lite Spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA) and tested with each primer pair using both cPCR and qPCR protocols. Collected eDNA was also diluted 1:10, 1:20, and 1:30 to check for PCR inhibition (Davy et al. 2015) such as those caused by accidental humic acid collection in samples (Thomsen and Willerslev 2015).

To minimize contamination among samples, we utilized filtered pipette tips, separate clean rooms for DNA extraction and PCR amplification, and all equipment was thoroughly sterilized with 20% diluted bleach, rinsed with sterilized water, and then dried under UV light for 30 min. Latex gloves were also used for each sample collected. After each filtering session, medical-grade sterilized gauze covering the inlet of the filter holder was replaced, and the filter holder and tweezers were washed in a 20% bleach dilution, rinsed with sterilized water, and allowed to be fully dry (Davy et al. 2015).

All collections were ethically conducted with permission from conservation managers and WWF officers affiliated with the Tian-e-Zhou Nature Reserve.

### Cost assessment

To make a simplified cost comparison between traditional visual, capture, and eDNA survey methods (both using cPCR and qPCR) to detect and quantify YFP populations, we used 2015 and 2016 data on visual monitoring efforts by WWF officers at the Tian-e-Zhou Nature Reserve (presence/absence). In reality, the full cost of such sampling would require the transportation of samples and/or sampling personnel to and from the reserve. However, this will vary among practitioners employing these techniques (i.e., whether or not they filter on site, and/or transit distance for visual observers/eDNA collectors) and our purpose is merely to create a streamlined evaluation among monitoring methods. Thus, we have restricted our comparison to average labor costs for this particular region (government recommendations), laboratory consumables, and boat rentals. We also

assume eDNA collections will be conducted in laboratories already fully equipped for extraction and analysis, that (species-specific) optimization of genetic analysis is already complete (e.g., this study; Ma et al. 2016), and that amplification failure is negligible given recommendations for detection cutoffs and the proven methodological sensitivity shown here. Cost assessments for eDNA surveys utilizing both cPCR and qPCR were based on 3–1 L water sample across 15 sites (1 site per 1 km of habitable reserve length to reduce potential overlap in the dispersion of eDNA detection signals; Goldberg et al. 2016), for a total of 45 samples. cPCR costs were calculated based on data presented in Ma et al. (2016), including consumables and extraction kits. Therefore, both visual and genetic methodological comparisons overlap in both year of data collection and geographical distribution.

### Data analysis

We derived equations of the line and slopes for each primer pair's ability to predict known concentrations of synthetic gene DNA (log copies/ $\mu\text{L}$ ) based on its qPCR critical threshold (Ct). From these correlations, we acquired Pearson's coefficient  $R^2$  to calculate the efficiency of each primer pair and direct managers towards the best markers for use in YFP eDNA analysis. To analyze whether DNA quantity (log copies/ $\mu\text{L}$ ) and primer pair influenced the critical threshold, we also performed a two-way ANOVA. All analyses were performed using JMP v.12.0 (SAS Institute Inc., Cary, NC, 1989–2007).

### Results

For our assays, we did not detect any evidence of inhibition. In general, amplification demonstrated 8/12 and 12/12 primer pairs successfully amplified for cPCR and qPCR respectively, suggesting good species detection *in situ* for the positive control (Tian-e-Zhou) and no amplification for the negative control (Gonghuwan). For cPCR analysis, primer pairs showed low sensitivity, ranging from the lowest at  $2.96 \times 10^{10}$  copies/ $\mu\text{L}$  for FP97, and the highest at  $1.95 \times 10^8$  copies/ $\mu\text{L}$  for FP147 required for positive detection (Table 1).

Our qPCR amplification demonstrated  $R^2$  values ranging between 0.993–0.999 for all primer pairs, and efficiency ranging from 67.02%–128.97%, with only six primer pairs falling within the standard acceptable range (MIQE guidelines) of 90–110% (Table 1). qPCR amplification also demonstrated a significant difference in slope among the 12 primer pairs ( $F_{23,301} = 25.459$ ,  $P < 0.001$ ), with primer ( $F_{11} = 2.72$ ,  $p = 0.002$ ) and DNA quantity (copies/ $\mu\text{L}$ ) ( $F_1 = 549.227$ ,  $P < 0.001$ ), but not their interaction ( $F_{11} = 0.835$ ,  $P = 0.61$ ), showing an influence on the quantification slope. Accordingly, *post hoc* analysis (Tukey's HSD test) demonstrated primer FP133 to statistically have the steepest slope ( $t$  ratio =

**Table 1** The amplification sensitivity (LOD; copies/ $\mu$ L) of 12 primer pairs using either cPCR (via DNA tissue dilution) or qPCR (synthetic gene dilutions), including the intercept, slope, and efficiency (%) of qPCR amplifications

Primer	qPCR					cPCR					
	LOD 5 copies/ $\mu$ L	LOD 25 copies/ $\mu$ L	LOD 50 copies/ $\mu$ L	LOD 500 copies/ $\mu$ L	LOD 5000 copies/ $\mu$ L	Intercept	Slope	Efficiency (%)	<i>In situ</i> Detection	Max. Sensitivity (copies/ $\mu$ L)	<i>In situ</i> Detection
FP76	31%	46%	77%	66.6%	100%	38.10	-3.27	97.18	+	$1.89 \times 10^{10}$	+
FP90	15%	46%	85%	100%	100%	40.26	-3.92	<i>128.95</i>	+	$3.19 \times 10^9$	+
FP97	23%	23%	62%	100%	100%	38.69	-3.33	99.85	+	$2.96 \times 10^{10}$	-
FP104	23%	23%	46%	100%	100%	37.61	-2.67	72.85	+	$2.76 \times 10^{10}$	-
FP133	31%	62%	85%	100%	100%	39.15	-3.00	<i>85.71</i>	+	$1.08 \times 10^{10}$	+
FP147	15%	46%	54%	100%	100%	37.83	-2.51	<i>67.02</i>	+	$1.95 \times 10^8$	+
FP161	54%	39%	85%	100%	100%	38.73	-3.37	101.66	+	$1.78 \times 10^9$	+
FP162	23%	23%	46%	100%	100%	37.69	-2.96	<i>84.09</i>	+	$1.77 \times 10^9$	-
FP171	23%	23%	23%	100%	100%	36.80	-3.13	91.12	+	$1.68 \times 10^9$	+
FP221	15%	15%	31%	100%	100%	38.39	-3.42	103.95	+	$6.5 \times 10^9$	-
FP233	39%	39%	70%	100%	100%	37.79	-2.65	<i>72.11</i>	+	$6.16 \times 10^9$	+
FP249	39%	39%	77%	33.3%	100%	38.42	-3.33	99.85	+	$5.77 \times 10^9$	+

Amplification efficiencies outside the optimal range (90–110%) are italicized. LOD percentage is represented as the number of positive amplifications out of 13 replicates. Primer pair eDNA detection for *in situ* analysis is presented as ‘+’ for positive detection and ‘-’ for no detection

3.16,  $P = 0.002$ ) and primer FP171 to statistically have the shallowest slope ( $t$  ratio = -3.63,  $P < 0.001$ ) of all primers tested (Supplementary Information). Indeed, for qPCR, primers FP171 and FP221 showed little increase in sample detection with an increase to LOD (5 to 50 copies/ $\mu$ L). All primer pairs however did show 100% detection rates at an LOD of 500 copies/ $\mu$ L with the exception of FP76 and FP249 (Table 1).

### Cost analysis

Visual surveys in Tian-e-Zhou Nature Reserve are conducted once per month, lasting 7 days in total. Costs accrued for visual surveys in Tian-e-Zhou, including boat rental fees, a three-person-team, and 7 days of labor, approximated to 8000 CNY per month—accordingly, 24,000 CNY per season or 96,000 CNY per year from 2015 to 2016. Additionally, once a year, WWF China officers also complete a capture monitoring protocol in which all individuals within the Tian-e-Zhou reserve are corralled for data collection (e.g., demographic, morphological, health inspection). For this process to be achieved, a minimum of 300,000 CNY was spent on a 10 day, 15 boat, 40 personnel endeavor (Table 2).

Environmental DNA sampling, however, including labor, filtering water collections, extractions, amplifications, and sequencing, ranged from 4257 CNY for cPCR to 5686 CNY for qPCR per sampling event (45 samples tested in triplicate) (Table 2). Given eDNA for this species can persist in the aquatic environment for 30 days or more (Ma et al. 2016), sampling events with the aim to demonstrate current population distribution (or relative abundance) could conceivably take place at most once a month, to approximately once per season.

Visual surveying on a monthly basis thus costs 1.88X that of eDNA collections utilizing cPCR (species detection) at the same temporal schedule. If, however, eDNA sampling using cPCR occurred only once per season, visual surveys would approximate 5.64X more expensive. Similarly, visual surveys compared to eDNA sampling utilizing qPCR would equate to 1.41X on a monthly sampling schedule, and 4.22X on a seasonal sampling schedule.

### Discussion

Efforts towards species conservation are only as good as the data used to measure a change in distribution and abundance. Yet, the high cost and substantial workforce required for species detection surveys, especially in developing countries, has hindered our ability to assess the status of species and populations globally. In part, this is reflected in depauperate information accrued in areas of low-governance but high biodiversity (including gaps in data collection), and due to difficult-to-survey habitats (Frazier et al. 2016) causing selection biases towards developed countries in temperate regions (McGeoch et al. 2010; Martin et al. 2012; Hudson et al. 2014) and terrestrial ecosystems (Mittermeier et al. 2011). Although rapidly emerging as a salient and valuable technology for biodiversity data collection, systematic appraisals for when to apply eDNA over traditional methods are scarce (but see Biggs et al. 2015; Davy et al. 2015; Sigsgaard et al. 2015; Smart et al. 2016), and comparisons between different eDNA approaches, to our knowledge, are non-existent to date. Given recent demonstration that traditional survey methods are not always more

**Table 2** Comparison of Yangtze finless porpoise (*Neophocaena asiaorientalis asiaorientalis*) survey cost estimates (Chinese Yuan; CNY) in the Tian-e-Zhou Baiji National Nature Reserve as of 2015–2016 (Chinese Yuan; CNY) between eDNA protocols (45 samples in triplicate) utilizing conventional PCR (cPCR) or quantitative PCR (qPCR), and traditional methods, such as visual or capture (WWF China; *personal communication*)

Survey method	Details	Cost (CNY)
cPCR	eDNA collection labor	495
	Filter papers+consumables	180
	Extraction QIAGEN DNEasy blood and tissue kit	1701
	Amplification	405
	Confirmation (visualization)	675
	PCR labor	486
	Total	4257
qPCR	eDNA collection labor	495
	Filter papers+consumables	180
	Extraction MOBIO DNEasy PowerWater kit	3701
	Amplification and quantification	230
	qPCR labor	1080
Total	5686	
Visual monitoring	$X = 7$ days, 1 boat, 3 personnel	
	Per month ( $1X$ )	8000
	Per season ( $3X$ )	24,000
	Per year ( $12X$ )	96,000
Capture monitoring	$X = 10$ days, 15 boats, 40 personnel	
	Per year ( $1X$ )	300,000

Traditional surveying (visual and capture) cost calculations are based on sampling per event ( $X$ ), currently employed once a month for visual monitoring, and once per year for capture monitoring. Sampling events are then multiplied based on time period

expensive than eDNA approaches, particularly for smaller census budgets and small passive sampling tactics (e.g., Smart et al. 2016), providing conservation managers with the most practical tools for their specific goals is imperative to speed data assembly.

For YFP monitoring, our analysis demonstrated cPCR to be the least expensive option for species-specific surveying compared to both qPCR and traditional methods. Nevertheless, as with other studies (e.g., Amberg et al. 2015), cPCR remained a less-sensitive tool compared to qPCR, and thus more prone to false-negatives, making it more useful for data collection in areas where larger populations are expected, or as a supplementary test for when false-negatives can be negated (e.g., when combined with visual surveying efforts). On the other hand, qPCR, a slightly more expensive option than cPCR, but comparatively cheaper than traditional visual surveys for YFP, demonstrated relatively high sensitivity and reproducibility for managers, suggesting high detection accuracy. Typically, thresholds for positive (or negative) detection scoring of samples in qPCR eDNA analyses can vary from study to study, spanning

one of three, one of eight (Jerde et al. 2011, 2013; Mahon et al. 2013; Piaggio et al. 2014; Rees et al. 2014), or two of twelve (Schneider et al. 2016) replicates. For our study, we present all scoring data ranging from two in thirteen (LOD of 15% at 5 copies/ $\mu$ L) to all thirteen replicates (LOD 100% at 5000 copies/ $\mu$ L) showing positive amplification, thereby allowing practitioners to establish their own level of confidence for eDNA detection (Table 1). In our study, the primer pairs representing the best efficiency and sensitivity via qPCR (FP76, FP97, FP161, FP171, FP221, and FP249) also demonstrated a minimal detection capability of > 23% positive scoring at an LOD of 25 DNA copies/ $\mu$ L (with the exception of FP221). This may suggest a high likelihood of picking up dilute eDNA signals either as a result of habitat nuances (i.e., large aquatic environments), or relatively small population numbers, with relatively low sampling replicates. Given the multitude of information qPCR could afford managers (distribution combined with relative abundance) while simultaneously increasing eDNA sensitivity, qPCR may offer the best option for species detection data collection for YFP. However, if managers are faced with financial constraints, either eDNA method may still represent a better alternative than visual monitoring for YFP presence/absence population counts. Despite its ability to accumulate myriad ecological observations (e.g., species demography, co-occurring species, and abiotic parameters such as weather or hydrological patterns), traditional visual surveys within Tian-e-Zhou remain a costly, invasive, and potentially inaccurate (e.g., due to organismal behavior) method for presence/absence data. Indeed, because the visual survey takes place within a reserve, its focus remains on YFP detection, with limited additional information. Whether the presented eDNA monitoring method is applicable to a fast-flowing environment such as the Yangtze River, however, still necessitates testing, though using these techniques in tributaries and lakes where nearly half the species population resides, remains promising.

Although eDNA analysis may require high initial costs (laboratory equipment, primer development, primer validation, and optimization of sampling protocols), our assessment corroborates the few other studies available (e.g., Biggs et al. 2015; Davy et al. 2015; Sigsgaard et al. 2015; Smart et al. 2016) suggesting eDNA methods may still represent the most cost-efficient means of acquiring biodiversity data, especially given budgetary constraints. Moreover, numerous companies are now available for outsourcing eDNA analyses, promising vast reductions to these initial expenses. Costs associated with eDNA protocols are also likely to mirror the observed decrease in DNA sequencing costs in general (Metzker 2010), causing eDNA sampling to become more cost-efficient over time.

Admittedly, our study only analyzed data from a single enclosed site (Tian-e-Zhou Nature Reserve) and because the hydrological dynamics of the Yangtze River are vastly different to that of enclosed lake, the costs required to eDNA

sample the entirety of the Yangtze River (and its tributaries) may exceed the cost of traditional visual methods currently employed. Still, managers are encouraged to consider the likelihood of detecting this taxon across its range with traditional (e.g., visual) protocols given known organismal ecology and behavior. For example, due to their preference for benthic prey (Park et al. 2011; Shirakihara et al. 2009), YFP have been observed to spend approximately 60% of their time on long dives (Beasley and Jefferson 2002) reducing opportunities for visual encounters. Visual surveys also inadvertently increase the probability of accidental harm or death due to boat traffic, either through a disruption in the species' sonar navigation ability or through fatal contact with ship propellers (Chen et al. 1997; Wang 2013). Moreover, collecting water samples for eDNA analysis requires less technical expertise than accurate taxonomic proficiency in morphological identification, a skill suggested to currently be in rapid decline (Wheeler et al. 2004). As the ecology of YFP (spring reproductive bouts, feeding on benthic invertebrates, and seasonal dispersal) may influence detection sensitivities and the amount of eDNA found within the water column (Goldberg et al. 2016), careful consideration of the sinks/sources of this information should remain paramount to managers before applying this methodology. However, this information could also give invaluable insights into the cryptic ecology of the species that may be otherwise difficult to attain. We thus encourage managers to consider the trade-offs between cost and data accuracy when considering which sampling method to employ when collecting valuable biodiversity data.

**Acknowledgments** The authors would like to thank the WWF China officers and Dr. Jinsong Zheng for assistance in data collection and access to the Tian-e-Zhou Nature Reserve.

**Funding information** The authors would like to thank the National Natural Sciences Foundation of China (KAS; 31400468) and the Tongji University State Key Laboratory for Pollution Control and Resource Reuse for financially supporting this research.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## References

- Amberg JJ, McCalla SG, Monroe E, Lance R, Baerwaldt K, Gaikowski MP (2015) Improving efficiency and reliability of environmental DNA analysis for silver carp. *J Great Lakes Res* 41:367–373. <https://doi.org/10.1016/j.jglr.2015.02.009>
- Balasingham KD, Walter RP, Heath DD (2017) Residual eDNA detection sensitivity assessed by quantitative real-time PCR in a river system. *Mol Ecol Resour* 17:523–532
- Beasley I, Jefferson TA (2002) Surface and dive times of finless porpoises in Hong Kong's coastal waters. *Raffles B Zool* 10:125–129
- Biggs J, Ewald N, Valentini A, Gaboriaud C, Dejean T, Griffiths RA, Foster J, Wilkinson JW, Amell A, Brotherton P, Williams P, Dunn F (2015) Using eDNA to develop a national citizen science-based monitoring programme for the great crested newt (*Triturus cristatus*). *Biol Conserv* 183:19–28. <https://doi.org/10.1016/j.biocon.2014.11.029>
- Chen P, Liu R, Wang D, Zhang X (1997) Biology, rearing and protection of Baiji. Academic Press, Beijing. (in Chinese)
- Danielsen F, Mendoza MM, Alviola P, Balette DS, Enghoff M, Poulsen MK, Jensen AE (2003) Biodiversity monitoring in developing countries: what are we trying to achieve? *Oryx* 37:407–409. <https://doi.org/10.1017/S0030605303000735>
- Davy CM, Kidd AG, Wilson CC (2015) Development and validation of environmental DNA (eDNA) markers for detection of freshwater turtles. *PLoS One* 10:e0130965. <https://doi.org/10.1371/journal.pone.0130965>
- Dejean T, Valentini A, Miquel C, Taberlet P, Bellemain E, Miaud C (2012) Improved detection of an alien invasive species through environmental DNA barcoding, the example of the American bullfrog *Lithobates catesbeianus*. *J Appl Ecol* 49:953–959. <https://doi.org/10.1111/j.1365-2664.2012.02171.x>
- Dougherty MM, Larson ER, Renshaw MA, Gantz CA, Egan SP, Erickson DM, Lodge DM (2016) Environmental DNA (eDNA) detects the invasive rusty crayfish *Orconectes rusticus* at low abundances. *J Appl Ecol* 53:722–732. <https://doi.org/10.1111/1365-2664.12621>
- Evans NT, Olds BP, Renshaw MA, Turner CR, Li Y, Jerde CL, Mahon AR, Pfrender ME, Lamberti GA, Lodge DM (2015) Quantification of mesocosm fish and amphibian species diversity via environmental DNA metabarcoding. *Mol Ecol Resour* 24:315–321. <https://doi.org/10.1111/1755-0998.12433>
- Ficetola GF, Miaud C, Pompanon F, Taberlet P (2008) Species detection using environmental DNA from water samples. *Biol Lett* 4:423–425. <https://doi.org/10.1098/rsbl.2008.0118>
- Ficetola GF, Pansu J, Bonin A, Coissac E, Giguët-Covex C, Barba MD et al (2015) Replication levels, false presences and the estimation of the presence/absence from eDNA metabarcoding data. *Mol Ecol Resour* 15:543–556. <https://doi.org/10.1111/1755-0998.12338>
- Foote AD, Thomsen PF, Sveegaard S, Wahlberg M, Kielgast J, Kyhn LA, Salling AB, Galatius A, Orlando L, Gilbert MTP (2012) Investigating the potential use of environmental DNA (eDNA) for genetic monitoring of marine mammals. *PLoS One* 7:e41781. <https://doi.org/10.1371/journal.pone.0041781>
- Frazier M, Longo C, Halpern BS (2016) Mapping uncertainty due to missing data in the global ocean health index. *PLoS One* 11:e0160377. <https://doi.org/10.1371/journal.pone.0160377>
- Fukumoto S, Ushimaru A, Minamoto T (2015) A basin-scale application of environmental DNA assessment for rare endemic species and closely related exotic species in rivers: a case study of giant salamanders in Japan. *J Appl Ecol* 52:358–365. <https://doi.org/10.1111/1365-2664.12392>
- Goldberg CS, Sepulveda A, Ray A, Baumgardt J, Waits LP (2013) Environmental DNA as a new method for early detection of New Zealand mudsnails (*Potamopyrgus antipodarum*). *Freshw Sci* 32:792–800. <https://doi.org/10.1899/13-046.1>
- Goldberg CS, Turner CR, Deiner K, Klymus KE, Thomsen PF, Murphy MA, Spear SF, McKee A, Oyler-McCance SJ, Cornman RS, Laramie MB, Mahon AR, Lance RF, Pilliod DS, Strickler KM, Waits LP, Fremier AK, Takahara T, Herder JE, Taberlet P (2016) Critical considerations for the application of environmental DNA

- methods to detect aquatic species. *Methods Ecol Evol* 7:1299–1307. <https://doi.org/10.1111/2041-210X.12595>
- Hänfling B, Lawson Handley L, Read DS, Hahn C, Li J, Nichols P, Blackman RC, Oliver A, Winfield IJ (2016) Environmental DNA metabarcoding of lake fish communities reflects long-term data from established survey methods. *Mol Ecol* 25:3101–3119. <https://doi.org/10.1111/mec.13660>
- Hudson LN, Newbold T, Contu S, Hill SL, Lysenko I, De Palma A, Phillips HR, Senior RA et al (2014) The PREDICTS database: a global database of how local terrestrial biodiversity responds to human impacts. *Ecol Evol* 4:4701–4735. <https://doi.org/10.1002/ece3.1303>
- Jerde CL, Mahon AR, Chadderton WL, Lodge DM (2011) “Sight-unseen” detection of rare aquatic species using environmental DNA. *Conserv Lett* 4:150–157. <https://doi.org/10.1111/j.1755-263X.2010.00158.x>
- Jerde CL, Chadderton WL, Mahon AR, Renshaw MA, Corush J, Budny ML, Mysorekar S, Lodge DM (2013) Detection of Asian carp DNA as part of a Great Lakes basin-wide surveillance program. *Can J Fish Aquat Sci* 70:522–526. <https://doi.org/10.1139/cjfas-2012-0478>
- Klymus KE, Richter CA, Chapman DC, Paukert C (2015) Quantification of eDNA shedding rates from invasive bighead carp *Hypophthalmichthys nobilis* and silver carp *Hypophthalmichthys molitrix*. *Biol Conserv* 183:77–84. <https://doi.org/10.1016/j.biocon.2014.11.020>
- Laramie MB, Pilliod DS, Goldberg CS (2015) Characterizing the distribution of an endangered salmonid using environmental DNA analysis. *Biol Conserv* 183:29–37. <https://doi.org/10.1016/j.biocon.2014.11.025>
- Li H (2011) Construction status and ecological protection of the Yangtze River basin nature reserve. *Resour Environ Yangtze Basin* 20:150–155 (in Chinese)
- Lodge DM, Turner CR, Jerde CL, Barnes MA, Chadderton L, Egan SP, Feder JL, Mahon AR, Pfreder ME (2012) Conservation in a cup of water: estimating biodiversity and population abundance from environmental DNA. *Mol Ecol* 21:2555–2558. <https://doi.org/10.1111/j.1365-294X.2012.05600.x>
- Ma H, Stewart KA, Ma L, Ren W, Zhao J (2015) Environmental DNA and its application in protecting aquatic ecosystems. *Chin J Ecol* 35(2):516–523
- Ma H, Stewart KA, Lougheed SC, Zheng J, Wang Y, Zhao J (2016) Characterization, optimization, and validation of environmental DNA (eDNA) markers to detect an endangered aquatic mammal. *Conserv Genet Resour* 8:1–8. <https://doi.org/10.1007/s12686-016-0597-9>
- Mahon AR, Jerde CL, Galaska M, Bergner JL, Chadderton WL, Lodge DM, Hunter ME, Nico LG (2013) Validation of eDNA surveillance sensitivity for detection of Asian carps in controlled and field experiments. *PLoS One* 8:e58316. <https://doi.org/10.1371/journal.pone.0058316>
- Margules CR, Pressey RL (2000) Systematic conservation planning. *Nature* 405:243–253
- Martin LJ, Blossey B, Ellis EC (2012) Mapping where ecologists work: biases in the global distribution of terrestrial ecological observations. *Front Ecol Environ* 10:195–201. <https://doi.org/10.2307/41480052>
- McGeoch MA, Butchart SHM, Spear D, Marais E, Kleynhans EJ, Symes A, Chanson J, Hoffmann M (2010) Global indicators of biological invasion: species numbers, biodiversity impact and policy responses. *Divers Distrib* 16:95–108. <https://doi.org/10.1111/j.1472-4642.2009.00633.x>
- Mei Z, Huang SL, Hao Y, Turvey ST, Gong W, Wang D (2012) Accelerating population decline of Yangtze finless porpoise (*Neophocaena asiaeorientalis asiaeorientalis*). *Biol Conserv* 153:192–200. <https://doi.org/10.1016/j.biocon.2012.04.029>
- Metzker ML (2010) Sequencing technologies—the next generation. *Nat Rev Genet* 11:31–46. <https://doi.org/10.1038/nrg2626>
- Mittermeier RA, Turner WR, Larsen FW, Brooks TM, Gascon C (2011) Global biodiversity conservation: the critical role of hotspots. In: Zachos FE, Habel JC (eds) *Biodiversity hotspots*. Springer, Berlin, pp 3–22
- Park KJ, An YR, Lee YR, Park JE, Moon DY, Choi SG (2011) Feeding habits and consumption by finless porpoises (*Neophocaena asiaeorientalis*) in the Yellow Sea. *Kor J Fish Aquat Sci* 44:78–84. <https://doi.org/10.5657/kfas.2011.44.1.078> (in Korean)
- Piaggio AJ, Engeman RM, Hopken MW, Humphrey JS, Keacher KL, Bruce WE, Avery ML (2014) Detecting an elusive invasive species: a diagnostic PCR to detect Burmese python in Florida waters and an assessment of persistence of environmental DNA. *Mol Ecol Resour* 14:374–380. <https://doi.org/10.1111/1755-0998.12180>
- Pilliod DS, Goldberg CS, Arkle RS, Waits LP (2013) Estimating occupancy and abundance of stream amphibians using environmental DNA from filtered water samples. *Can J Fish Aquat Sci* 70:1123–1130. <https://doi.org/10.1139/cjfas-2013-0047>
- Rees HC, Maddison BC, Middleditch DJ, Patmore JR, Gough KC (2014) The detection of aquatic animal species using environmental DNA—a review of eDNA as a survey tool in ecology. *J Appl Ecol* 51:1450–1459. <https://doi.org/10.1111/1365-2664.12306>
- Schneider J, Valentini A, Dejean T, Montarsi F, Taberlet P, Glairot O, Fumagalli L (2016) Detection of invasive mosquito vectors using environmental DNA (eDNA) from water samples. *PLoS One* 11:e0162493. <https://doi.org/10.1371/journal.pone.0162493>
- Shaw JLA, Clarke LJ, Wedderburn SD, Barnes TC, Weyrich LS, Cooper A (2016) Comparison of environmental DNA metabarcoding and conventional fish survey methods in a river system. *Biol Conserv* 197:131–138. <https://doi.org/10.1016/j.biocon.2016.03.010>
- Shirakihara M, Seki K, Takemura A, Shirakihara K, Yoshida H, Yamazaki T (2009) Food habits of the finless porpoise, *Neophocaena phocaenoides* in Western Kyushu, Japan. *J Mammal* 98:1248–1256. <https://doi.org/10.1644/07-MAMM-A-264.1>
- Sigsgaard EE, Carl H, Møller PR, Thomsen PF (2015) Monitoring the near-extinct European weather loach in Denmark based on environmental DNA from water samples. *Biol Conserv* 183:46–52. <https://doi.org/10.1016/j.biocon.2014.11.023>
- Smart AS, Weeks AR, van Rooyen AR, Moore A, McCarthy MA, Tingley R (2016) Assessing the cost-efficiency of environmental DNA sampling. *Methods Ecol Evol* 7:1291–1298. <https://doi.org/10.1111/2041-210X.12598>
- Stewart KA, Ma H, Zheng J, Zhao J (2017) Using environmental DNA to assess population-wide spatiotemporal reserve use. *Conserv Biol* 31(5):1173–1182
- Takahara T, Minamoto T, Yamanaka H, Doi H, Kawabata Z (2012) Estimation of fish biomass using environmental DNA. *PLoS One* 7:e35868. <https://doi.org/10.1371/journal.pone.0035868>
- Tamura K, Stecher G, Peterson D, Filipksi A, Kumar S (2013) MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30:2725–2729
- Thomsen PF, Willerslev E (2015) Environmental DNA—an emerging tool in conservation for monitoring past and present biodiversity. *Biol Conserv* 183:4–18
- Thomsen PF, Kielgast J, Iversen LL, Wiuf C, Rasmussen M, Gilbert MTP, Orlando L, Willerslev E (2012) Monitoring endangered freshwater biodiversity using environmental DNA. *Mol Ecol* 21:2565–2573. <https://doi.org/10.1111/j.1365-294X.2011.05418.x>
- Valentini A, Taberlet P, Miaud C, Civade R, Herder J, Thomsen PF, Bellemain E, Besnard A, Coissac E, Boyer F, Gaboriaud C, Jean P, Poulet N, Roset N, Copp GH, Geniez P, Pont D, Argillier C, Baudoin JM, Peroux T, Crivelli AJ, Olivier A, Acqueberge M, le Brun M, Møller PR, Willerslev E, Dejean T (2016) Next-generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. *Mol Ecol* 25:929–942. <https://doi.org/10.1111/mec.13428>



- Wan X, Ruan R, McLaughlin RW, Hao Y, Zheng J, Wang D (2016) Fecal bacterial composition of the endangered Yangtze finless porpoises living under captive and semi-natural conditions. *Curr Microbiol* 72: 306–314. <https://doi.org/10.1007/s00284-015-0954-z>
- Wang D (2013) Population status, threats, and conservation of the Yangtze finless porpoise. *Chin Sci Bull* 54:3473–3484. <https://doi.org/10.1007/s11434-009-0522-7>
- Wheeler QD, Raven PH, Wilson EO (2004) Taxonomy: impediment or expedient? *Science* 303:285. <https://doi.org/10.1126/science.303.5656.285>
- Ye J, Coulouris G, Zaretskaya L, Cutcutache L, Rozen S, Madden TL (2012) Primer-BLAST: a tool to design target-specific primers for polymerase chain reaction. *BMC Bioinform* 13: 134