

The *GCKR* rs780094 polymorphism is associated with elevated fasting serum triacylglycerol, reduced fasting and OGTT-related insulinaemia, and reduced risk of type 2 diabetes

T. Sparsø · G. Andersen · T. Nielsen · K. S. Burgdorf ·
A. P. Gjesing · A. L. Nielsen · A. Albrechtsen ·
S. S. Rasmussen · T. Jørgensen · K. Borch-Johnsen ·
A. Sandbæk · T. Lauritzen · S. Madsbad · T. Hansen ·
O. Pedersen

Published online: 7 December 2007
© Springer-Verlag 2007

Unfortunately, there was a mistake in the last sentence of the Abstract. The Conclusions/Interpretation should have read as follows:

‘The *GCKR* rs780094 polymorphism, or another variant with which it is in tight linkage disequilibrium, is likely to

increase glucokinase regulatory protein activity to induce improved glycaemic regulation at the expense of hypertriacylglycerolaemia as reflected in the present study of 16,853 Danes. We also suggest an additive effect of *GCK* and *GCKR* risk alleles on serum insulin release.’

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-007-0865-z>.

T. Sparsø (✉) · G. Andersen · T. Nielsen · K. S. Burgdorf ·
A. P. Gjesing · A. L. Nielsen · S. S. Rasmussen ·
K. Borch-Johnsen · T. Hansen · O. Pedersen
Steno Diabetes Center,
Niels Steensens Vej 1, NLC2.13,
DK-2820 Gentofte, Denmark
e-mail: tspr@steno.dk

A. Albrechtsen
Department of Biostatistics, University of Copenhagen,
Copenhagen, Denmark

T. Jørgensen · K. Borch-Johnsen
Research Centre for Prevention and Health,
Glostrup University Hospital,
Glostrup, Denmark

K. Borch-Johnsen · O. Pedersen
Faculty of Health Science, University of Aarhus,
Aarhus, Denmark

A. Sandbæk · T. Lauritzen
Department of General Practice, University of Aarhus,
Aarhus, Denmark

S. Madsbad
Department of Epidemiology, Hvidovre University Hospital,
Hvidovre, Denmark