

## **Erratum to: Physical activity attenuates the mid-adolescent peak in insulin resistance but by late adolescence the effect is lost: a longitudinal study with annual measures from 9–16 years (EarlyBird 66)**

**Brad S. Metcalf<sup>1,2</sup> · Joanne Hosking<sup>3</sup> · William E. Henley<sup>1</sup> · Alison N. Jeffery<sup>3</sup> ·  
Mohammad Mostazir<sup>4</sup> · Linda D. Voss<sup>3</sup> · Terence J. Wilkin<sup>1</sup>**

Published online: 10 September 2015  
© Springer-Verlag Berlin Heidelberg 2015

**Erratum to: Diabetologia**  
**DOI 10.1007/s00125-015-3714-5**

Unfortunately, the affiliation for Mohammad Mostazir was incorrect in this paper. His correct affiliation is listed below.

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00125-015-3714-5>.

---

✉ Brad S. Metcalf  
[b.metcalf@exeter.ac.uk](mailto:b.metcalf@exeter.ac.uk)

- <sup>1</sup> Institute of Health Research, University of Exeter Medical School, Exeter, UK
- <sup>2</sup> Sport and Health Sciences, College of Life and Environmental Sciences, University of Exeter, St Luke's Campus, Heavitree Road, Exeter EX1 2LU, UK
- <sup>3</sup> Department of Endocrinology and Metabolism, Plymouth University Peninsula Schools of Medicine and Dentistry, Plymouth, UK
- <sup>4</sup> Wellcome Trust Biomedical Informatics Hub, College of Life and Environmental Sciences (CLES), University of Exeter, Exeter, UK