

Erratum to: The factor-of-risk biomechanical approach predicts hip fracture in men and women: the Framingham Study

A. B. Dufour · B. Roberts · K. E. Broe · D. P. Kiel ·
M. L. Bouxsein · M. T. Hannan

Published online: 20 March 2014

© International Osteoporosis Foundation and National Osteoporosis Foundation 2014

Erratum to: Osteoporos Int DOI 10.1007/s00198-011-1569-2

This article contains an error on page 515, right column, where the equations given for calculating trochanteric soft tissue thickness in men and women separately are mislabeled. The sentences in question should read as follows:

“For women, trochanteric soft tissue thickness (millimeters) is equal to $3.4795 \times (\text{BMI (kilograms per square meter)}) - 38.015$. For men, trochanteric soft tissue thickness (millimeters) is equal to $2.3415 \times (\text{BMI (kilograms per square meter)}) - 33.444$.”

The authors apologise for any inconvenience caused.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00198-011-1569-2>.

A. B. Dufour
Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA

A. B. Dufour (✉) · K. E. Broe · D. P. Kiel · M. T. Hannan
Institute for Aging Research, Hebrew SeniorLife, 1200 Centre St., Boston, MA 02131, USA
e-mail: alysadufour@hsl.harvard.edu

K. E. Broe
e-mail: broe@hsl.harvard.edu

D. P. Kiel
e-mail: kiel@hsl.harvard.edu

M. T. Hannan
e-mail: hannan@hsl.harvard.edu

D. P. Kiel · M. L. Bouxsein · M. T. Hannan
Harvard Medical School, Boston, MA, USA

M. L. Bouxsein
e-mail: mbouxsei@bidmc.harvard.edu

B. Roberts · M. L. Bouxsein
Orthopedic Biomechanics Laboratory, Beth Israel Deaconess Medical Center, Boston, MA, USA

B. Roberts
e-mail: bjrobert@bidmc.harvard.edu

D. P. Kiel · M. T. Hannan
Division of Gerontology, Department of Medicine, Beth Israel Deaconess Medical Center, Boston, MA, USA