

## Erratum to: Determination of $\omega$ -6 and $\omega$ -3 PUFA metabolites in human urine samples using UPLC/MS/MS

Ai Sasaki · Hayato Fukuda · Narumi Shiida ·  
Nobuaki Tanaka · Ayako Furugen · Jiro Ogura ·  
Satoshi Shuto · Nariyasu Mano · Hiroaki Yamaguchi

Received: 23 January 2015 / Accepted: 23 January 2015 / Published online: 24 February 2015  
© Springer-Verlag Berlin Heidelberg 2015

**Erratum to: Analytical and Bioanalytical Chemistry**  
**DOI: 10.1007/s00216-014-8412-5**

The original publication of this article unfortunately contains errors. “PD1 (Protectin D1)” should be “PDX (Protectin DX)”. Figure S1(b) is also incorrect and the corrected figure is attached below. These errors do not affect the conclusion in the paper.

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00216-014-8412-5>.


---

A. Sasaki · N. Shiida · A. Furugen · J. Ogura  
Laboratory of Clinical Pharmaceutics and Therapeutics, Faculty of  
Pharmaceutical Sciences, Hokkaido University, Sapporo, Japan

H. Fukuda · S. Shuto  
Laboratory of Organic Chemistry for Drug Development, Faculty of  
Pharmaceutical Sciences, Hokkaido University, Sapporo, Japan

N. Tanaka · N. Mano · H. Yamaguchi  
Laboratory of Clinical Pharmacy, Graduate School of Pharmaceutical  
Sciences, Tohoku University, Sendai, Japan

N. Mano · H. Yamaguchi (✉)  
Department of Pharmaceutical Sciences, Tohoku University  
Hospital, 1-1 Seiryō-machi, Aoba-ku, Sendai 980-8574, Japan  
e-mail: yamaguchi@hosp.tohoku.ac.jp



**Fig. S1** Metabolic pathway of  $\omega$ -3 PUFAs (b) including urinary metabolites. The compounds indicated by bold face are the target analytes in our method. HpDHA, hydroperoxy docosahexaenoic acid; HpETE, hydroperoxy eicosatetraenoic acid