

Calvarial Osteoclasts Express a Higher Level of Tartrate-Resistant Acid Phosphatase Than Long Bone Osteoclasts and Activation Does not Depend on Cathepsin K or L Activity

S. Perez-Amadio · D. C. Jansen · T. Schoenmaker ·
I. M. C. Vogels · T. Reinheckel · A. R. Hayman ·
T. M. Cox · P. Saftig · W. Beertsen · V. Everts

Published online: 14 February 2007
© Springer Science+Business Media, Inc. 2007

Erratum to: Calcif Tissue Int (2006) 79:245–254
DOI 10.1007/s00223-005-0289-z

Due to a production error in the text, but not in the figures,
TRAP should be read as TRACP.

The online version of the original article can be found under
doi:[10.1007/s00223-005-0289-z](https://doi.org/10.1007/s00223-005-0289-z).

S. Perez-Amadio · D. C. Jansen · T. Schoenmaker ·
W. Beertsen
Experimental Periodontology, Academic Center for Dentistry
Amsterdam, Universiteit van Amsterdam and Vrije Universiteit,
Louwesweg 1, 1066 EA Amsterdam, The Netherlands

I. M. C. Vogels
Department of Cell Biology and Histology, Academic Medical
Center, Universiteit van Amsterdam, P.O. Box 22700,
1100 DE Amsterdam, The Netherlands

T. Reinheckel
Department of Molecular Medicine and Cell Research,
Albert-Ludwigs-Universität Freiburg, Freiburg 79104,
Germany

A. R. Hayman
School of Clinical Veterinary Science, University of Bristol,
Langford, Bristol BS40 5DU, UK

T. M. Cox
Department of Medicine, University of Cambridge,
Addenbrooke's Hospital, Hills Road, Cambridge
CB2 2QQ, UK

P. Saftig
Biochemical Institute, Christian-Albrechts University,
Olshausenstr. 40, 24098 Kiel, Germany

V. Everts (✉)
Department of Oral Cell Biology, Academic Centre for Dentistry
Amsterdam, Universiteit van Amsterdam and Vrije Universiteit,
Van der Boechorststraat 7, 1081 BT Amsterdam,
The Netherlands
e-mail: V.Everts@vumc.nl