

Erratum to: Changes in ecosystem engineers: the effects of kelp forest type on currents and benthic assemblages in Kachemak Bay, Alaska

Dominic Hondolero¹  · Matthew S. Edwards²

Published online: 12 April 2017
© Springer-Verlag Berlin Heidelberg 2017

Erratum to: Mar Biol (2017) 164:81
DOI [10.1007/s00227-017-3111-3](https://doi.org/10.1007/s00227-017-3111-3)

Unfortunately, Fig. 4 was incorrectly published in the original version. The correct Fig. 4 is updated here and in the original publication as well.

The online version of the original article can be found under doi:[10.1007/s00227-017-3111-3](https://doi.org/10.1007/s00227-017-3111-3).

The original article was corrected.

✉ Dominic Hondolero
Dominic.Hondolero@noaa.gov

¹ Kasitsna Bay Laboratory, NOAA/NOS/CCFHR, Seldovia, AK 99663, USA

² Department of Biology, San Diego State University, San Diego, CA 92182, USA




Fig. 4 nMDS plots showing relative similarities in invertebrate settlement on Tuffys placed at different locations (inside versus outside) in the two forest types (*Nereocystis luetkeana* versus *Eualaria fistulosa*) measured each week (Date) during the experiment. Symbols that are nearer to one another represent samples that are more similar in spe-

cies composition. Triangles refer to samples from within the *Eualaria fistulosa* beds and squares refer to samples from within the *Nereocystis leutkeana* beds. Filled (black) symbols refer to samples from inside the kelp beds and open (white) symbols refer to samples from outside the kelp beds