



# Correction to: Comparing the horizontal and vertical approaches used to identify foraging areas of two diving marine predators

Yann Planque<sup>1</sup> · Mathilde Huon<sup>1</sup> · Florence Caurant<sup>1</sup> · David Pinaud<sup>2</sup> · Cécile Vincent<sup>1</sup>

Published online: 5 August 2020  
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

**Correction to: Marine Biology (2020) 167:25**  
<https://doi.org/10.1007/s00227-019-3636-8>

The author would like to correct the Fig. 4 of the published article. The correct Fig. 4 is shown below:

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---


The original article can be found online at <https://doi.org/10.1007/s00227-019-3636-8>.

---

✉ Yann Planque  
yann.planque@univ-lr.fr

<sup>1</sup> Centre d'Etudes Biologiques de Chizé, UMR 7372, CNRS - La Rochelle Université, 5 allée de l'Océan, 17000 La Rochelle, France

<sup>2</sup> Centre d'Etudes Biologiques de Chizé, UMR 7372, CNRS - La Rochelle Université, 405 route de La Canauderie, 79360 Villiers-en-Bois, France



**Fig. 4** Grey and harbour seal foraging areas determined using the horizontal (left) and vertical (right) approaches. The foraging areas are represented on grids of 2.5 × 2.5 km cells and characterized by the spatial density of dives in ARS selected from the hori-

zontal approach and faster U-shaped dives selected from the vertical approach. Cores to extended foraging areas are represented by 90% kernel densities of foraging locations (black lines). *BDS* Baie de Somme, the capture site