

Erratum to: Ultrasound-guided joint injections for MR arthrography in pediatric patients: how we do it

Jeffrey Otjen¹ · Shawn E. Parnell¹ · Sarah Menashe¹ ·
Mahesh M. Thapa¹

Published online: 2 June 2015
© Springer-Verlag Berlin Heidelberg 2015

Erratum to: Pediatr Radiol
DOI 10.1007/s00247-014-3212-9

A formatting error occurred in the production of Table 1 Summary of ultrasound guided joint injection techniques. As a result, incorrect information is given for the line “Volume estimate.” The correct entry should read as follows:

	Shoulder	Elbow	Hip	Knee	Ankle	Subtalar
Volume estimate*	15–20 cc	3–7 cc	15–20 cc	30–40 cc	3–7 cc	3–7 cc

*Volume varies from patient to patient and with age and size. This estimate reflects an average adolescent volume to generate adequate joint capsule distention

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00247-014-3212-9>.

✉ Mahesh M. Thapa
thapamd@u.washington.edu

¹ Department of Radiology, Seattle Children’s Hospital,
University of Washington School of Medicine,
4800 Sand Point Way NE M/S-5417,
Seattle, WA 98105, USA