

Erratum to: Development of SCAR markers for rapid and specific detection of *Pseudomonas syringae* pv. *morsprunorum* races 1 and 2, using conventional and real-time PCR

Monika Kałużna¹ · Pedro Albuquerque^{2,3} · Fernando Tavares^{2,3} · Piotr Sobiczewski¹ · Joanna Puławska¹

Published online: 20 December 2016
© Springer-Verlag Berlin Heidelberg 2016

Erratum to: This complements the above paper in Appl Microbiol Biotechnol, volume 100 on page 3705
DOI 10.1007/s00253-016-7295-0

In the original version of this article inadvertently there is a lack of the information concerning the annealing temperature used for primers designed for conventional PCR.

It should be written:

For the primers designed for conventional PCR, among the tested annealing temperatures, 59°C for primers Psm1-6F and Psm1-6R (for detection of *Psm1* strains) and 54°C for primers Psm2-8F and Psm2-8R (for detection of *Psm2* strains) were selected as the optimal ones.

The online version of the original article can be found at doi:10.1007/s00253-016-7295-0.

✉ Monika Kałużna
monika.kaluzna@inhort.pl

¹ Research Institute of Horticulture, Konstytucji 3 Maja 1/3, 96-100 Skierniewice, Poland

² Centro de Investigação em Biodiversidade e Recursos Genéticos (CIBIO), InBIO, Laboratório Associado, Universidade do Porto, Campus Agrário de Vairão, 4485-661 Vairão, Portugal

³ Faculdade de Ciências, Departamento de Biologia, Universidade do Porto, Edifício FC4, Via Panorâmica No. 36, 4150-564 Porto, Portugal