

## Male mating effort in a polygynous ungulate

Fanie Pelletier · John T. Hogg · Marco Festa-Bianchet

Published online: 8 August 2006  
© Springer-Verlag 2006

Unfortunately, in Fig. 1, an incorrect *x*-axis title appeared. Here is the correct figure:


The online version of the original article can be found at: <http://dx.doi.org/10.1007/s00265-006-0208-y>.

F. Pelletier · M. Festa-Bianchet  
Département de Biologie, Université de Sherbrooke,  
2500 Boulevard de l'Université,  
Sherbrooke, Quebec J1K 2R1, Canada

M. Festa-Bianchet  
e-mail: M.Festa@Usherbrooke.ca

F. Pelletier (✉)  
Division of Biology, Faculty of Natural Sciences,  
Imperial College London,  
Silwood Park,  
Ascot, Berkshire SL5 7PY, UK  
e-mail: fanie.pelletier@imperial.ac.uk

J. T. Hogg  
Montana Conservation Science Institute, Limited,  
5200 Upper Miller Creek Road,  
Missoula, MT 59803, USA  
e-mail: jthogg@montana.com



**Fig. 1** Rank-specific male mating behavior ( $\pm$ SE) for bighorn rams during 14 breeding seasons (1989–2002) at Sheep River, Alberta. **a** Searching index (in meters per day), **b** courting index, and **c** tending index