

ERRATUM

Erratum to: Cone Beam Computed Tomography (CBCT) in the Field of Interventional Oncology of the Liver

Blanche Bapst¹ · Matthieu Lagadec¹ · Romain Breguet⁴ ·

Valérie Vilgrain^{1,2,3} · Maxime Ronot^{1,2,3}

Published online: 1 August 2015

© Springer Science+Business Media New York and the Cardiovascular and Interventional Radiological Society of Europe (CIRSE) 2015

Erratum to: Cardiovasc Intervent Radiol DOI: 10.1007/s00270-015-1180-6

In the published article, it is indicated that the rotation duration and the X-ray time delay of the CBCT scanning are 8 and 6–9 s, respectively (Table 1) in the study by Paul et al. The correct values are 2.72 and 6 or 9 s (for global and segmental injection), respectively.

The online version of the original article can be found under
doi:[10.1007/s00270-015-1180-6](https://doi.org/10.1007/s00270-015-1180-6).

✉ Maxime Ronot
maxime.ronot@bjn.aphp.fr

Blanche Bapst
blanchebapst@hotmail.com

Matthieu Lagadec
matthieu.lagadec@bjn.aphp.fr

Romain Breguet
romain.breguet@hcuge.ch

Valérie Vilgrain
Valerie.vilgrain@bjn.aphp.fr

¹ Department of Radiology, Beaujon Hospital, University Hospitals Paris Nord Val de Seine, Beaujon, 100 bd general Leclerc, 92118 Clichy, Hauts-de-Seine, France

² University Paris Diderot, Sorbonne Paris Cité, Paris, France

³ INSERM U1149, Centre de Recherche Biomédicale Bichat-Beaujon, CRB3, Paris, France

⁴ University Hospital of Geneva, Geneva, Switzerland