

Erratum to: Shifts in Antarctic megabenthic structure after ice-shelf disintegration in the Larsen area east of the Antarctic Peninsula

Julian Gutt · Mattias Cape · Werner Dimmler ·
Laura Fillinger · Enrique Isla · Verena Lieb ·
Tomas Lundälv · Christian Pulcher

Published online: 24 January 2014
© Springer-Verlag Berlin Heidelberg 2014

Erratum to: Polar Biol (2013) 36:895–906 DOI 10.1007/s00300-013-1315-7

In the original paper the labels of the abscissae of the plots shown in Fig. 6 (running years) were erroneously displaced. Here the corrected Fig. 6 is provided. The one-year shift in the year labels does not affect the general conclusions of the paper.

The online version of the original article can be found under
doi:10.1007/s00300-013-1315-7.

J. Gutt (✉) · L. Fillinger · V. Lieb · C. Pulcher
Alfred Wegener Institute Helmholtz Centre for Polar and Marine
Research, Columbusstr., 27568 Bremerhaven, Germany
e-mail: julian.gutt@awi.de

M. Cape
Scripps Institution of Oceanography, University of California,
8615 Discovery Way, La Jolla, San Diego, CA 92037, USA

W. Dimmler
Fielax Gesellschaft für wissenschaftliche Datenverarbeitung
mbH, Schleusenstr. 14, 27568 Bremerhaven, Germany

E. Isla
Institut de Ciències del Mar-CSIC, Passeig Marítim de la
Barceloneta, 37-49, 08003 Barcelona, Spain

V. Lieb
Christian-Albrechts-Universität Kiel, Christian-Albrechts-Platz 4,
24118 Kiel, Germany

T. Lundälv
Sven Lovén Centre for Marine Sciences, University
of Gothenburg, 45296 Strömstad, Sweden

C. Pulcher
Institut für Biologie und Umweltwissenschaften, Carl von
Ossietzky Universität Oldenburg, Carl-von-Ossietzky-Straße
9-11, 26129 Oldenburg, Germany

Fig. 6 Mean sea-ice concentration for a region of the Larsen A (**a**) and Larsen B (**b**) embayments. Mean values appear as *open circles*, while the *line* represents a 7-day moving average. A threshold of 15 % sea-ice concentration generally indicates open-water conditions. Areas of analysis are depicted in Fig. 1

