



## Correction to: Three-dimensional black-blood contrast-enhanced MRI improves detection of intraluminal thrombi in patients with acute ischaemic stroke

Won Jang<sup>1</sup> · Hyo Sung Kwak<sup>1</sup> · Gyung Ho Chung<sup>1</sup> · Seung Bae Hwang<sup>1</sup>

Published online: 19 November 2018  
© European Society of Radiology 2018

**Correction to: European Radiology (2018) 28:3840–3847**  
<https://doi.org/10.1007/s00330-018-5323-4>

The original version of this article, published on 19 March 2018, unfortunately contained a mistake. In the section “MR examination,” the contrast medium Gadoterate meglumine was incorrectly named Gadodiamide. The correct wording is “Gadoterate meglumine 0.1 mmol/kg (Dotarem; Guerbet, Aulnay-sous-Bois, France) was injected as a bolus intravenously in all patients.”

---

The online version of the original article can be found at <https://doi.org/10.1007/s00330-018-5323-4>

---

✉ Hyo Sung Kwak  
[kwak8140@jbnu.ac.kr](mailto:kwak8140@jbnu.ac.kr)

<sup>1</sup> Radiology and Research Institute of Clinical Medicine of Chonbuk National University-Biomedical Research Institute of Chonbuk National University Hospital, 567 Baekje-daero, Deokjin-gu, Jeonju-si, Jeollabuk-do 561-756, Republic of Korea