



Correction to: Effectiveness of percutaneous vacuum-assisted excision (VAE) of breast lesions of uncertain malignant potential (B3 lesions) as an alternative to open surgical biopsy

Elisabetta Giannotti¹ · Jonathan J. James¹ · Yan Chen² · Rachel Sun¹ · Amanjot Karuppiah¹ · Julia Yemm³ · Andrew H. S. Lee⁴

Published online: 5 July 2021
© European Society of Radiology 2021

Correction to: European Radiology

<https://doi.org/10.1007/s00330-021-08060-z>

The original version of this article, published on 08 June 2021, unfortunately contained a mistake. The following correction has therefore been made in the original: The spelling of Julia Yemm's name was incorrect. The corrected author list is given above. The original article has been corrected.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00330-021-08060-z>

✉ Elisabetta Giannotti
Elisabetta.Giannotti@nuh.nhs.uk

¹ Nottingham Breast Institute, Nottingham University Hospitals NHS Trust, Nottingham City Hospital Hucknall Road, Nottingham NG51PB, UK

² Division of Cancer and Stem Cells, School of Medicine, University of Nottingham, Nottingham City Hospital Hucknall Road, Nottingham NG51PB, UK

³ Department of Radiology, King's Mill Hospital, Mansfield Road, Sutton-In-Ashfield, Mansfield NG174JL, UK

⁴ Department of Histopathology, Nottingham University Hospitals NHS Trust, Nottingham City Hospital Hucknall Road, Nottingham NG51PB, UK