

# Different ways in which CO<sub>2</sub> can be released during the turnover of roots in soil

Venkata S. S. R. Marella<sup>1</sup> · Paula Roberts<sup>1</sup> · Paul W. Hill<sup>1</sup> · Davey L. Jones<sup>1</sup>

Received: 22 September 2016 / Revised: 20 February 2017 / Accepted: 7 March 2017 / Published online: 23 March 2017  
© The Author(s) 2017. This article is published with open access at Springerlink.com

**Abstract** Here, we investigated how root age and mode of death influenced their subsequent turnover and rate of C loss from soil. Young white-coloured and older pigmented roots of *Cistus monspeliensis* were excised (to simulate death by mechanical severance) or frozen (to simulate death by cell rupture) and immediately buried in soil. CO<sub>2</sub> loss from soil was then measured over time. In a parallel experiment, the rate of CO<sub>2</sub> loss from severed or ruptured roots in the absence of soil was determined. Our results revealed large differences in root chemistry related to age, with young roots having a lower C:N ratio and a greater nutrient content (soluble C, N, P and K). Both root age and mode of death resulted in very different temporal patterns of C release from soil. The amount of C lost from soil followed the series: severed white roots ( $42.6 \pm 3.3$  mg C) > ruptured pigmented roots ( $27.7 \pm 0.4$  mg C) = ruptured white roots ( $27.1 \pm 0.5$  mg C) > severed pigmented roots ( $10.1 \pm 1.0$  mg C) > soil only ( $3.0 \pm 0.2$  mg C). Therefore, depending on the treatment, 7 to 41% of the total root-derived C was lost as CO<sub>2</sub> over the duration of the experiment. Comparison with soil-free treatments revealed that the CO<sub>2</sub> release from the severed roots buried in soil was not associated with microbial breakdown but caused by root-induced autophagy in an attempt to keep themselves metabolically active. Ruptured roots also induced a rapid loss of CO<sub>2</sub> which we ascribe to the diffusive loss of root solutes into the soil and subsequent microbial mineralization. Surprisingly, the rate of C loss from soil was

greater from the severed root tips than those that were ruptured. Our results imply two distinct routes of C loss dependent on how roots die, one which bypasses the microbial community and one which flows through it.

**Keywords** Apoptosis · Carbon cycling · Heterotrophic respiration · Necromass · Rhizosphere · Rhizodeposition · Carbon sequestration

## Introduction

Fine roots contribute approximately 33% of the annual net primary production in terrestrial ecosystems (Jackson et al. 1997). Upon senescence, their turnover is responsible for the formation of stable humic material, the release of nutrients that contribute to soil fertility, whilst also releasing substantial amounts of CO<sub>2</sub> back to the atmosphere. Fine root decomposition is plant species dependent and is regulated by a range of plant factors (e.g. presence of mycorrhizal symbionts, age and physiological state) and edaphic factors (Ghidey and Alberts 1993; Silver and Miya 2001; Redin et al. 2014; Goebel et al. 2011; Han et al. 2015; Tahir et al. 2016). The quantity, quality and lability of root C inputs to soil are also dependent on the way in which roots enter the necromass pool (Jones et al. 2004; van Doorn et al. 2011). At least three distinct pathways for fine root death can be identified:

- (1) Type I death: Programmed (vacuolar) root cell death where C and nutrients are re-translocated to other growing areas of the plant prior to metabolic activity ceasing leaving just a nutrient-poor root corpse behind (e.g. occurs when a root has exhausted a patch of soil and is no longer needed or occurs during senescence and seed setting of annual crops; Jones et al. 2004).

**Electronic supplementary material** The online version of this article (doi:10.1007/s00374-017-1193-1) contains supplementary material, which is available to authorized users.

✉ Davey L. Jones  
d.jones@bangor.ac.uk

<sup>1</sup> School of Environment, Natural Resources & Geography, Bangor University, Gwynedd LL57 2UW, UK

- (2) Type II death: Necrotic root death, where live roots become excised from the host plant (e.g. severing of during tillage, by meso/macrofauna, during tree windthrow) but still remain structurally intact and metabolically active. As the supply of C from the host plant is cut off, the roots maintain their supply of energy by breaking down all available internal C reserves. This autophagy progressively exhausts all internal soluble and structural C reserves before the roots ultimately die many days or weeks later (Saglio and Pradet 1980). There is no nutrient re-translocation, and the cell corpse (largely insoluble and structural C) remains largely unprocessed.
- (3) Type III death: Catastrophic root death where metabolic activity in the root ceases immediately, and all C and nutrients from the roots are free to enter soil (e.g. occurs upon freezing-induced cell lysis, from severe mechanical damage in which the cells rupture; Schaberg et al. 2008; Kreyling et al. 2012; Muliele et al. 2015).

Each mode of death is likely to greatly influence how C and nutrients flow through the soil as well as influencing the size, activity and composition of the soil microbial community. The aim of the study was therefore to elucidate how root age and the mode of death influences root decomposition and thus the rate of C return to the atmosphere. We hypothesise that a Type III cell death would lead to a much more rapid release of CO<sub>2</sub> from soil in comparison to a Type II cell death due to the greater availability of labile C for the soil microbial community. We also hypothesise that in a Type II root death, most of the C release in the early stages is plant-derived rather than soil-microbially derived.

## Materials and methods

### Plant material

Plants of the dwarf shrub *Cistus monspeliensis* L. were propagated through cuttings and grown in hydroponic culture using half-strength Long Ashton nutrient solution (Hewitt 1966). Plants were grown at 20 °C, with a light intensity (photosynthetically active radiation) of 200 μmol m<sup>-2</sup> s<sup>-1</sup> at canopy height and 16 h photoperiod. After 6 months (plant height 40 cm), young white roots (<7 days old) and older highly pigmented (brown) roots (>30 days old) were harvested from the plants. Immediately after excision, sub-samples of each root type were placed into polypropylene microcentrifuge tubes and killed by immersion in liquid N<sub>2</sub> (de Neergaard et al. 2000). The cell sap was then recovered by the centrifugal-drainage procedure outlined in Hill et al. (2008), and the sap analysed for the following properties: total C and N using a Multi N/C 2100S analyser (AnalytikJena, Germany), total free amino acids (TFAA) using the

fluorometric *o*-phthaldialdehyde β-mercaptoethanol procedure of Jones et al. (2002), nitrate using the VCl<sub>3</sub> method of Miranda et al. (2001), ammonium using the salicylate-nitroprusside-hypochlorite procedure of Mulvaney (1996) and total soluble phenolics using the Folin Ciocalteu method of Swain and Hillis (1959). The roots were also analysed for the distribution of soluble and structural components (cellulose, hemicellulose, lignin and ash) using an ANKOM2000 automated fibre analyser (Ankom Technology, Macedon, NY; Karcher et al. 2015). Root nutrient content was determined after drying (80 °C, 16 h) by total reflection X-ray fluorescence spectroscopy using a S2-PICOFOX-TXRF (Bruker, Billerica, MA).

### Soil

A sandy clay loam textured soil (Eutric Cambisol) was collected (Ah horizon; 0–10 cm depth) from Abergwyngregyn, North Wales, UK (53° 14' N, 4° 01' W). Physical, chemical and biological properties of the soil are detailed in Hill et al. (2008). Immediately after collection, the soil was transferred to the laboratory, sieved (<2 mm) and stored at 5 °C. Soil moisture content was determined by oven drying (105 °C, 16 h).

### Root mineralisation

Newly excised white or pigmented roots (3 g, ca. 96 mg C) were placed into individual 50 cm<sup>3</sup> sterile polypropylene tubes. Half the roots were then frozen (−80 °C, 30 min) to instantly kill them. Roots were then mixed with field-moist soil (30 g) to give five treatments: (i) severed white roots plus soil, (ii) severed pigmented roots plus soil, (iii) ruptured white roots plus soil, (iv) ruptured pigmented roots plus soil and (v) soil-only control. There were three replicates of each treatment, and the level of soil disturbance was identical in all treatments. In addition, the four root treatments were also studied in an identical way but in the absence of soil (i.e. root-only controls). All treatments were incubated at 20 °C and CO<sub>2</sub> evolution measured using an automated, multichannel SR1 respirometer (PP Systems, Hitchin, UK). The SR1 respirometer has a built-in humidifier unit which prevents moisture loss from the samples. The CO<sub>2</sub> flux derived solely from roots when buried in soil (root<sub>soil</sub> CO<sub>2</sub> flux) was estimated by subtracting the background soil respiration (soil<sub>only</sub> CO<sub>2</sub> flux; unamended controls) from the total CO<sub>2</sub> flux values for each mesocosm.

### Statistical analysis

Differences in root chemistry were compared by one-way ANOVA, whilst differences in C mineralisation were compared using a repeated measure ANOVA with post-hoc testing using SPSS v20.0 (SPSS Inc., Chicago, IL). Cumulative CO<sub>2</sub>

evolution in the root treatments over the duration of the experiment were compared with a two-way ANOVA with the presence of soil and root type as factors.  $P < 0.05$  was used as the cut-off for statistical significance.

**Results**

Analysis of the two different root types revealed large differences in their chemistry (Table 1). Overall, the white roots had a much higher intrinsic nutrient content (soluble C, N, P and K) and much less structural C by weight, although the lignin concentration did not vary with root age, and the concentration of Ca was much higher in the older roots.

C mineralisation rates varied quantitatively and temporally between white and pigmented roots, and the mechanism of root death also influenced C mineralisation rates, especially during the first 72 h (Fig. 1a; Fig. S1;  $P < 0.001$ ). Overall, the rate of respiration was low in the unamended soil


**Table 1** Major nutrient composition of young white and older pigmented roots of *Cistus monspeliensis* L. Chemical characteristics of the cell sap and soluble components of white and pigmented roots

		Root type	
		White	Pigmented
<b>Cell sap</b>			
Total soluble C	(mg l <sup>-1</sup> )	3093 ± 181a <sup>c</sup>	1085 ± 186b
Total soluble N	(mg l <sup>-1</sup> )	758 ± 34a	231 ± 40b
Free amino acids	(mg l <sup>-1</sup> )	312 ± 16a	25 ± 12b
NH <sub>4</sub> <sup>+</sup> -N	(mg l <sup>-1</sup> )	62.5 ± 4.6	58.6 ± 1.1
NO <sub>3</sub> <sup>-</sup> -N	(mg l <sup>-1</sup> )	152 ± 7a	100 ± 3b
Soluble phenols	(mg l <sup>-1</sup> )	262 ± 1a	240 ± 2b
<b>Whole root</b>			
Detergent soluble <sup>a</sup>	(% DW) <sup>b</sup>	46.0 ± 0.7a	34.4 ± 0.5b
Hemicellulose	(% DW)	14.2 ± 0.5a	16.3 ± 0.1b
Cellulose	(% DW)	20.7 ± 1.1a	29.7 ± 1.0b
Lignin	(% DW)	18.7 ± 0.5	19.3 ± 0.5
Ash	(% DW)	0.38 ± 0.02	0.32 ± 0.05
Total C	(g kg <sup>-1</sup> DW)	425 ± 11	410 ± 6
Total N	(g kg <sup>-1</sup> DW)	44.5 ± 1.3a	28.7 ± 0.1b
C-to-N ratio		9.5 ± 0.1a	14.3 ± 0.2b
Total P	(g kg <sup>-1</sup> DW)	12.8 ± 0.9a	5.1 ± 0.3b
Total K	(g kg <sup>-1</sup> DW)	16.8 ± 1.2a	4.7 ± 0.3b
Total Ca	(g kg <sup>-1</sup> DW)	6.4 ± 0.4a	12.1 ± 0.4b

<sup>a</sup> Includes fats, oils and waxes and soluble cell contents (carbohydrates, lipids, pectin, starch, and soluble proteins)

<sup>b</sup> DW dry weight

<sup>c</sup> Means (±SEM, n = 3) followed by different letters indicate significant differences between root types at the  $P < 0.05$  level. The absence of letters indicates no significant difference



**Fig. 1** Cumulative production of CO<sub>2</sub> from soil after burial of either severed or ruptured roots of *Cistus monspeliensis* (a), and the production of CO<sub>2</sub> from the same roots in the absence of soil (b). The roots were either young (white) or old (pigmented). Values represent means ± SEM (n = 3). Values are expressed per mesocosm to allow direct comparison between treatments (i.e. 30 g soil, 30 g soil + 3 g roots or 3 g roots)

(0.48 ± 0.06 mg CO<sub>2</sub> kg<sup>-1</sup> h<sup>-1</sup>), with much greater CO<sub>2</sub> fluxes seen in the root-amended soils ( $P < 0.001$ ). Over the duration of the experiment (300 h), the total CO<sub>2</sub> flux was significantly different between the soil treatments ( $P < 0.001$ ; Fig. 1) and followed the series:

Severed white roots (42.6 ± 3.3 mg C) > ruptured pigmented roots (27.7 ± 0.4 mg C) = ruptured white roots (27.1 ± 0.5 mg C) > severed excised pigmented roots (10.1 ± 1.0 mg C) > soil only (3.0 ± 0.2 mg C)

Therefore, depending on the treatment, 7 to 41% of the total root-derived C was lost as CO<sub>2</sub> over the duration of the experiment.

When the CO<sub>2</sub> flux for the two severed root treatments was measured in the absence of soil (i.e. roots only), the respiratory fluxes were extremely similar to those of the root + soil treatments ( $P > 0.05$ ; Fig. 1b, Fig. 2). In contrast, in the two ruptured root-only treatments, the rate of CO<sub>2</sub> loss was initially (0–24 h) much less than the parallel treatment performed in soil ( $P < 0.01$  for white roots and  $P < 0.001$  for pigmented roots). Following this (24–96 h), the rate of CO<sub>2</sub> efflux became similar in both the soil and soil-free ruptured root treatments. Following this, a major difference was seen between CO<sub>2</sub> efflux in the soil and soil-free treatments but only in the ruptured white root treatment ( $P < 0.05$ ; Fig. 2). A two-way ANOVA showed no significant interaction of soil presence and root type on the total amount of CO<sub>2</sub> produced over the course of the experiment ( $F = 0.23$ ,  $P = 0.874$ ).


## Discussion

As expected, tissue chemistry varied substantially with root age. Our findings are consistent with previous studies in which the concentration of low molecular weight solutes (e.g. amino acids, organic acids and sugars), and their rate of turnover inside root tissues was greatest in the most actively growing root regions (Jones and Darrah 1996). The higher major nutrient concentrations (N, P, K) in young white roots in comparison

to pigmented roots reported here are also consistent with earlier studies showing nutrient translocation from older roots towards more physiologically active parts of the plant (Goldfarb et al. 1990; Volder et al. 2005). Our study also indicated that older pigmented roots had much greater structural, cell wall-associated C in comparison to younger roots. This brown pigmentation of older roots is concurrent with but not caused by suberisation and has been taken as an indicator of a general decline in root function (Comas et al. 2000; Cruz et al. 2004). Our results indicate that young roots should turnover much faster than older roots, and their lower C:N:P ratio should lead to the more rapid release of nutrients back into the soil.

Our data demonstrated that severed roots exhibited high rates of CO<sub>2</sub> efflux from soil. Almost all the additional CO<sub>2</sub> released from the severed roots immediately after burial in soil, however, could be attributed CO<sub>2</sub> loss from the roots themselves, rather than by breakdown of the roots by the soil microbial community. This is consistent with visual observation of root integrity by James et al. (1993) and Bingham and Rees (2008) who showed that excised roots continue to respire for up to 30 days after excision. Initially, respiration is fuelled by stored sugars (e.g. glucose, starch and sucrose; 0–12 h), after which point-free amino acids reserves get catabolised and used in respiration. When these labile C pools are depleted, a general rise in autoprotoleolytic activity is observed. Although the roots are undergoing progressive autophagy, evidence suggests that plasma membrane integrity is maintained for weeks after excision and that no large exudative flux of organic C and N solutes occurs during this time (Saglio and Pradet 1980; Fig. S2). In contrast to Bingham and Rees (2008), however, we showed a significant loss of NH<sub>4</sub><sup>+</sup> into the soil from the severed roots after excision (Fig. S2). We ascribe this to the excess production of NH<sub>4</sub><sup>+</sup> during proteolysis and the need to remove it from the cytosol to prevent it reaching toxic levels (Brouquisse et al. 1992, 1998). No such loss of amino acids or NO<sub>3</sub><sup>-</sup> into the soil was observed.

In contrast to roots which can remain alive after excision, roots exposed to low freezing temperatures undergo a catastrophic and immediate death as a result of ice crystal growth and disruption of the cell's membranes and cytoskeleton (Wesley-Smith et al. 2015). At this point, CO<sub>2</sub> efflux can only occur due to a residual loss of HCO<sub>3</sub><sup>-</sup> from the cells and CO<sub>2</sub> produced by the rhizosphere microbial community. Our results showed that CO<sub>2</sub> efflux from the ruptured roots in the absence of soil was initially very low; however, this rapidly increased over 12 h suggesting that this was due to the growth or activation of microbes present on the rhizoplane and endorhizosphere. When buried in soil, this pulse of CO<sub>2</sub> release from the ruptured roots was immediate, consistent with the rapid loss of low molecular weight solutes from the roots into the soil by passive diffusion and their rapid consumption and mineralization by a C-limited microbial community (Hill et al. 2008).



**Fig. 2** Ratio of the CO<sub>2</sub> output derived from severed or ruptured *Cistus monspeliensis* roots either in the presence (root<sub>soil</sub> CO<sub>2</sub> flux) or absence of soil (Root<sub>only</sub> CO<sub>2</sub> flux). The roots were either young (white) or old (pigmented). The background soil respiration (i.e. soil without roots present) has been subtracted from the soil + root treatments

The roots used here were grown in hydroponic culture and consequently no mycorrhizal symbionts were present. Further, the experiments were undertaken with fresh bulk soil which may have possessed a different microbial community from that naturally present in the *Cistus* rhizosphere. Under natural conditions, both of these represent additional factors that can influence the rate of root senescence and C loss from soil, particularly for older roots (Eissenstat et al. 2000). Theoretically, the addition of labile root C may induce either negative or positive priming of soil organic matter turnover. Based on our previous measurements of priming in the same soil using  $^{14}\text{C}$ -labelled organic matter and a range of organic materials (e.g. biochar, wood ash, compost, manure and shoot residues), however, these responses are normally very much lower (ca.  $-10$  to  $+15\%$  changes in basal respiration) than the large positive  $\text{CO}_2$  responses observed here (Fig. 1; Jones et al. 2011; Reed et al. unpubl.). We therefore expect that the contribution of priming to the overall  $\text{CO}_2$  fluxes observed here will be very low.

## Conclusions

This study clearly shows that root age and mode of death results in very different temporal patterns of C release from soil. Contrary to expectation, the rapid release of  $\text{CO}_2$  from soil after root excision could not be attributed to the microbial breakdown of root tissues. Our evidence, alongside that of others, suggests that this pulse of  $\text{CO}_2$  release is more related to root-induced autophagy in which the root attempts futilely to keep itself metabolically active. Similarly, roots which underwent an immediate death by freezing also resulted in a rapid loss of  $\text{CO}_2$  from soil. In this case, however, we ascribe this loss to the diffusion of solutes from ruptured cells into the soil which are subsequently mineralized by the soil microbial community. Surprisingly, the rate of C loss from soil was greater from the senescing excised root tips than those that died immediately. This suggests a flow of C out of the soil which largely bypasses the soil microbial community. Severed older roots were much less metabolically active and more resistant to microbial degradation than either young roots or older ruptured roots, leading to a greater persistence of C in the soil. From our findings, it is clear that more work is needed to characterise and quantify the different ways in which roots die and turnover in soil and how this subsequently impacts upon long-term soil C storage and nutrient cycling. In addition, a greater attention should be paid to the role of mycorrhizal symbionts and endorhizosphere microbial communities in the root decomposition process. Finally, more long-term studies under field conditions are needed to complement laboratory experiments such as those performed here.

**Acknowledgements** We would like to thank Dr. Paul Baker, Gordon Turner, Jonathan Roberts and Sarah Chesworth for the technical support. We also thank Douglas Godbold for the scientific input and the choice of plant species. Funding was provided by project ENV4970694 funded under the European Union FP4-ENV 2C programme.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Bingham JJ, Rees RM (2008) Senescence and N release from clover roots following permanent excision of the shoot. *Plant Soil* 303:229–240
- Brouquisse R, James F, Pradet A, Raymond P (1992) Asparagine metabolism and nitrogen distribution during protein degradation in sugar-starved maize root tips. *Planta* 188:384–395
- Brouquisse R, Gaudillère JP, Raymond P (1998) Induction of a carbon starvation-related proteolysis in whole maize plants submitted to light/dark cycles and to extended darkness. *Plant Physiol* 117:1281–1291
- Comas LH, Eissenstat DM, Lakso AN (2000) Assessing root death and root system dynamics in a study of grape canopy pruning. *New Phytol* 147:171–178
- Cruz C, Green JJ, Watson CA, Wilson F, Martins-Loução MA (2004) Functional aspects of root architecture and mycorrhizal inoculation with respect to nutrient uptake capacity. *Mycorrhiza* 14:177–184
- van Doorn WG, Beers EP, Dangl JL, Franklin-Tong VE, Gallois P, Hara-Nishimura I, Jones AM, Kawai-Yamada M, Lam E, Mundy J, Mur LAJ, Petersen M, Smertenko A, Taliensky M, Van Breusegem F, Wolpert T, Woltering E, Zhivotovsky B, Bozhkov PV (2011) Morphological classification of plant cell deaths. *Cell Death Differ* 18:1241–1246
- Eissenstat DM, Wells CE, Yannai RD, Whitbeck JL (2000) Building roots in a changing environment. *New Phytol* 147:33–42
- Ghidey F, Alberts EE (1993) Residue type and placement effects on decomposition: field study and model evaluation. *Trans Am Soc Agric Eng* 36:1611–1617
- Goebel M, Hobbie SE, Bulaj B, Zadworny M, Archibald DD, Oleksyn J, Reich PB, Eissenstat DM (2011) Decomposition of the finest root branching orders: linking belowground dynamics to fine root function and structure. *Ecol Monogr* 8:89–102
- Goldfarb D, Hendrick R, Pregitzer K (1990) Seasonal nitrogen and carbon concentrations in white, brown and woody fine roots of sugar maple (*Acer saccharum* Marsh). *Plant Soil* 126:144–148
- Han E, Kautz T, Perkins U, Uteau D, Peth S, Huang N, Hom R, Kopke U (2015) Root growth dynamics inside and outside of soil biopores as affected by crop sequence determined with the profile wall method. *Biol Fert Soils* 51:847–856
- Hewitt EJ (1966) Sand and water culture methods used in the study of plant nutrition. Technical Communication No. 22. Commonwealth Bureau, London
- Hill PW, Farrar JF, Jones DL (2008) Decoupling of microbial glucose uptake and mineralization in soil. *Soil Biol Biochem* 40:616–624
- Jackson RB, Mooney HA, Schulze ED (1997) A global budget for fine root biomass, surface area, and nutrient contents. *Proc Natl Acad Sci U S A* 94:7362–7366
- James F, Brouquisse R, Pradet A, Raymond P (1993) Changes in proteolytic activities in glucose-starved maize root tips. Regulation by sugars. *Plant Physiol Biochem* 31:845–856

- Jones DL, Darrah PR (1996) Re-sorption of organic compounds by roots of *Zea mays* L. and its consequences in the rhizosphere. 3. Characteristics of sugar influx and efflux. *Plant Soil* 178:153–160
- Jones DL, Owen AG, Farrar JF (2002) Simple method to enable the high resolution determination of total free amino acids in soil solutions and soil extracts. *Soil Biol Biochem* 34:1893–1902
- Jones DL, Hodge A, Kuzyakov Y (2004) Plant and mycorrhizal regulation of rhizodeposition. *New Phytol* 163:459–480
- Jones DL, Murphy DV, Khalid M, Ahmad W, Edwards-Jones G, DeLuca TH (2011) Short-term biochar-induced increase in soil CO<sub>2</sub> release is both biotically and abiotically mediated. *Soil Biol Biochem* 43:1723–1731
- Karcher MA, Iqbal Y, Lewandowski I, Senn T (2015) Comparing the performance of *Miscanthus × giganteus* and wheat straw biomass in sulfuric acid based pretreatment. *Bioresour Technol* 180:360–364
- Kreyling J, Persoh D, Werner S, Benzenberg M, Wollecke J (2012) Short-term impacts of soil freeze-thaw cycles on roots and root-associated fungi of *Holcus lanatus* and *Calluna vulgaris*. *Plant Soil* 353:19–31
- Miranda KM, Espey MG, Wink DA (2001) A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. *Nitric Oxide* 5:62–71
- Muliele MT, Biolders CL, van Asten PJA (2015) Short- and medium-term impact of manual tillage and no-tillage with mulching on banana roots and yields in banana-bean intercropping in the East African highlands. *Field Crops Res* 171:1–10
- Mulvaney RL (1996) Nitrogen—inorganic forms. In: Sparks DL (ed) *Methods of soil analysis*. Soil Science Society of America Inc. American Society of Agronomy Inc., Madison, pp 1123–1184
- de Neergaard E, Lyshede OB, Gahoonia TS, Care D, Hooker JE (2000) Anatomy and histology of roots and root-soil boundary. In: Smit AL, Bengough AG, Engels C, van Noordwijk M, Pellerin S, van de Geijn SC (eds) *Root methods*. Springer-Verlag, Berlin Heidelberg, pp 33–73
- Redin M, Guénon R, Recous S, Schmatz R, de Freitas LL, Aita C, Giacomini SJ (2014) Carbon mineralization in soil of roots from twenty crop species, as affected by their chemical composition and botanical family. *Plant Soil* 378:205–214
- Saglio PH, Pradet A (1980) Soluble sugars, respiration, and energy charge during aging of excised maize root-tips. *Plant Physiol* 66:516–519
- Schaberg PG, Hennon PE, D'Amore DV, Hawley GJ (2008) Influence of simulated snow cover on the cold tolerance and freezing injury of yellow-cedar seedlings. *Glob Change Biol* 14:1282–1293
- Silver WL, Miya RK (2001) Global patterns in root decomposition: comparisons of climate and litter quality effects. *Oecologia* 129:407–419
- Swain T, Hillis WE (1959) The phenolic constituents of *Prunus domestica*. I.—The quantitative analysis of phenolic constituents. *J Agric Food Sci* 10:63–68
- Tahir MM, Recous S, Aita C, Schmatz R, Pilecco GE, Giacomini SJ (2016) In situ roots decompose faster than shoots left on the soil surface under subtropical no-till conditions. *Biol Fert Soils* 52:853–865
- Volder A, Smart DR, Bloom AJ, Eissenstat DM (2005) Rapid decline in nitrate uptake and respiration with age in fine lateral roots of grapes: implications for root efficiency and competitive effectiveness. *New Phytol* 165:493–502
- Wesley-Smith J, Walters C, Pammenter NW, Berjak P (2015) Why is intracellular ice lethal? A microscopical study showing evidence of programmed cell death in cryo-exposed embryonic axes of recalcitrant seeds of *Acer saccharinum*. *Ann Bot* 115:991–1000