



## Correction to: Ex situ reconstruction of comminuted radial head fractures: is it truly worth a try?

Jens Everding<sup>1</sup> · Michael J. Raschke<sup>1</sup> · Patrick Polgart<sup>1</sup> · Niklas Grüneweller<sup>1</sup> · Dirk Wähnert<sup>1</sup> · Benedikt Schliemann<sup>1</sup>

Published online: 3 January 2020  
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

**Correction to:**  
**Archives of Orthopaedic and Trauma Surgery**  
**(2019) 139:1723–1729**  
<https://doi.org/10.1007/s00402-019-03250-3>

The original version of this article unfortunately contained a mistake. The second sentence in the second paragraph of the discussion section reports “Businger et al. [18] found excellent results in two patients with on table reconstruction who demonstrated an average range of motion of 0°–6°–141° for extension-flexion and an average MEPS of 99.2 points.”

The correct sentence should be:

“Businger et al. [18] found excellent results in six patients with on table reconstruction who demonstrated an average range of motion of 0°–6°–141° for extension-flexion and an average MEPS of 99.2 points.”

**Publisher’s Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The original article can be found online at <https://doi.org/10.1007/s00402-019-03250-3>.

---

✉ Jens Everding  
jens.everding@ukmuenster.de  
<http://www.traumacentrum.de>

<sup>1</sup> Department of Trauma, Hand and Reconstructive Surgery, University Hospital Münster, Albert-Schweitzer-Campus, Building W1, 48149 Munster, Germany