



Correction to: Hypofractionated sequential radiotherapy boost: a promising strategy in inoperable locally advanced pancreatic cancer patients

Gian Carlo Mattiucci^{1,2} · Luca Boldrini^{1,2} · Alessia Nardangeli² · Andrea D'Aviero¹  · Milly Buwenge³ · Francesco Cellini² · Francesco Deodato⁴ · Nicola Dinapoli² · Vincenzo Frascino² · Gabriella Macchia⁴ · Alessio Giuseppe Morganti³ · Vincenzo Valentini^{1,2}

Published online: 9 November 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to:
Journal of Cancer Research and Clinical Oncology
<https://doi.org/10.1007/s00432-020-03411-7>

The original article has been corrected.

In the original article published, the last sentence in the sixth paragraph of the introduction section is incorrect. The correct sentence is “This data, were confirmed also in an Italian phase II study, in which it has been demonstrated that gemcitabine-based chemoradiotherapy was correlated with improved overall survival, especially in pts who are clinically more fit to complete the foreseen treatment schedule (CRT) (Mattiucci et al. 2010)”.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00432-020-03411-7>.

✉ Andrea D'Aviero
andreadav89@gmail.com

- ¹ Università Cattolica del Sacro Cuore, Roma, Italy
- ² UOC Radioterapia Oncologica, Dipartimento Di Diagnostica Per Immagini, Radioterapia Oncologica ed Ematologia, Fondazione Policlinico Universitario “A. Gemelli” IRCCS, Roma, Italy
- ³ Radiation Oncology Center, Department of Experimental, Diagnostic and Specialty Medicine (DIMES), University of Bologna, S. Orsola-Malpighi Hospital, Bologna, Italy
- ⁴ Unità Operativa Di Radioterapia, Gemelli Molise Hospital, Campobasso, Italy