

Lower Bounds in Real Algebraic Geometry and Orientability of Real Toric Varieties

Evgenia Soprunova · Frank Sottile

Received: 10 November 2012 / Revised: 9 March 2013 / Accepted: 19 March 2013 /
Published online: 3 April 2013
© Springer Science+Business Media New York 2013

Abstract The real solutions to a system of sparse polynomial equations may be realized as a fiber of a projection map from a toric variety. When the toric variety is orientable, the degree of this map is a lower bound for the number of real solutions to the system of equations. We strengthen previous work by characterizing when the toric variety is orientable. This is based on work of Nakayama and Nishimura, who characterized the orientability of smooth real toric varieties.

Keywords Real toric variety · Polynomial system · Order polytope

Mathematics Subject Classification (1991) 14M25 · 57B20 · 57S10 · 14P99

1 Introduction

A ubiquitous phenomenon in enumerative real algebraic geometry is that many geometric problems possess a non-trivial lower bound on their number of real solutions. For example, at least 3 of the 27 lines on a real cubic surface are real as are at least 8 of the 12 rational cubics interpolating 8 real points in the plane [4, Prop. 4.7.3], but there are many, many other examples [1, 6, 7, 9, 16, 17, 19, 20, 23]. This phenomenon has the potential for significant impact on the applications of mathematics as a nontrivial lower

E. Soprunova
Department of Mathematics, Kent State University, Summit Street, Kent, OH 44242, USA
e-mail: soprunova@math.kent.edu
URL: <http://www.math.kent.edu/~soprunova/>

F. Sottile (✉)
Department of Mathematics, Texas A&M University, College Station, TX 77843, USA
e-mail: sottile@math.tamu.edu
URL: <http://www.math.tamu.edu/~sottile>

bound is an existence proof of real solutions. For this potential to be realized, methods need to be developed to predict when a system of polynomial equations or a geometric problem has a lower bound on its number of real solutions and to compute this bound.

We developed a theory of lower bounds on the number of real solutions to systems of sparse polynomials [20]. There, a system of polynomial equations was formulated as a fiber of a projection map from a toric subvariety of a sphere. When the toric variety is orientable, the absolute value of the degree of this projection map is a lower bound on the number of real solutions. Besides giving a condition implying this orientability, a method (foldable triangulations of the Newton polytope) was developed to compute the degree of certain maps, and a class of examples of polynomial systems (Wronski polynomial systems from posets) was presented to which this theory applied.

Further work [12, 13] on foldable triangulations has advanced our understanding of the bound they give. Others [1, 7, 11, 16, 17] have developed additional methods for proving lower bounds in real algebraic geometry and experimentation [10, 19, 20] has revealed many more likely examples of lower bounds.

We characterize which sparse polynomial systems possess a lower bound in the context of [20], by extending work of Nakayama and Nishimura [15], who characterized the orientability of small covers, which are topological versions of smooth real projective toric varieties. We characterize the orientability of the smooth points of any real toric variety, as well as toric subvarieties of a sphere, solving an important open problem from [20].

We review the construction of real toric varieties and spherical toric varieties in Sect. 2, where we formulate our results on orientability. Section 3 contains the mildly technical proof of these results. In Sect. 4 we use this characterization of orientability to strengthen results from [20] on the theory of lower bounds for the number of real solutions to systems of sparse polynomials.

2 Constructions of Real Toric Varieties

Real toric varieties appear in many applications of mathematics [2, 14, 18] and are interesting objects in their own right [5]. Davis and Januszkiewicz [3] introduced the notion of a small cover of a simple convex polytope as a generalization of smooth projective real toric varieties. We describe real toric varieties and small covers in terms of the gluing of explicit cell complexes and give a mild extension of Davis and Januszkiewicz's notion of a small cover (which are manifolds) to not necessarily smooth spaces. A projective toric variety may be lifted to the sphere over real projective space, and we also describe these spherical toric varieties in terms of the gluing of explicit cell complexes.

Nakayama and Nishimura [15] used this presentation of small covers to characterize their orientability, and similar arguments characterize the orientability of the smooth points of the above spaces.

Real toric varieties, singular small covers, and toric subvarieties of the sphere are obtained by gluing the real torus $\mathbb{T}^n := (\mathbb{R}^\times)^n$ or $\{\pm 1\} \times \mathbb{T}^n = \mathbb{T}^{n+1}/\mathbb{R}_\geq$ along copies

of \mathbb{T}^{n-1} , one copy for each vector in a set of integer vectors. There are further gluings in higher codimension, which presents these spaces as explicit cell complexes. They are smooth at the points of their dense torus \mathbb{T}^n (or at $\{\pm 1\} \times \mathbb{T}^n$) and the attached tori \mathbb{T}^{n-1} , and so their orientability is determined by the gluing along the tori \mathbb{T}^{n-1} .

Complex toric varieties are normal varieties over \mathbb{C} equipped with an action of an algebraic torus $(\mathbb{C}^\times)^n$ having a dense orbit. They are classified by rational fans Σ in \mathbb{R}^n , which encode their construction as a union of affine toric varieties U_σ , one for each cone $\sigma \in \Sigma$. A toric variety is a union of disjoint torus orbits \mathcal{O}_σ , one for each cone $\sigma \in \Sigma$, with $\dim \mathcal{O}_\sigma = n - \dim(\sigma)$. The dense orbit \mathcal{O}_0 coincides with the smallest affine patch U_0 , and both are associated to the origin 0 in the fan. See [8] for a complete description.

A toric variety has a canonical set Y of real points obtained from the real points of the orbits \mathcal{O}_σ of the construction [8, Ch. 4]. The dense orbit $\mathcal{O}_0(\mathbb{R}) \simeq \mathbb{T}^n$ is isomorphic to $(\mathbb{R}^\times)^n = (\mathbb{R}_{>0})^n \times \{\pm 1\}^n$, which has 2^n components, each a topological n -ball. The subgroup $\{\pm 1\}^n \subset \mathbb{T}^n$ acts on Y , permuting the components of $\mathcal{O}_0(\mathbb{R})$. The orbit space of Y under the group $\{\pm 1\}^n$ is isomorphic to the closure Y_{\geq} of any component of $\mathcal{O}_0(\mathbb{R})$ in the usual topology (not Zariski!) on Y . Each orbit $\mathcal{O}_\sigma(\mathbb{R})$ has a unique component contained in Y_{\geq} . We call this component F_σ a *face* of Y_{\geq} , which is isomorphic to $(\mathbb{R}_{>0})^{n-\dim(\sigma)}$. This endows Y_{\geq} with the structure of a cell complex that is dual to the fan Σ . That is, the intersection $\overline{F_\sigma} \cap \overline{F_\tau}$ of the closures of two faces is nonempty only if σ and τ lie in some cone of Σ , in which case it is the closure $\overline{F_\rho}$ where ρ is the minimal such cone.

The integer points in a cone σ of Σ form a subsemigroup of \mathbb{Z}^n whose image in $(\mathbb{Z}/2\mathbb{Z})^n = \{\pm 1\}^n$ is a subgroup $\overline{\sigma}$ of $\{\pm 1\}^n$. This subgroup $\overline{\sigma}$ is the isotropy subgroup of the face F_σ of Y_{\geq} . We will write $(-1)^v = ((-1)^{v_1}, \dots, (-1)^{v_n})$ for the image of $v \in \mathbb{Z}^n$ in $(\mathbb{Z}/2\mathbb{Z})^n = \{\pm 1\}^n$. This gives the following description of Y as a quotient space of $Y_{\geq} \times \{\pm 1\}^n$.

Proposition 2.1 *The real toric variety Y is obtained as the quotient of the cell complex $Y_{\geq} \times \{\pm 1\}^n$ by the equivalence relation where*

$$(p, \xi) \sim (q, \eta) \iff p = q \text{ and } \xi \overline{\sigma} = \eta \overline{\sigma}, \text{ where } p \text{ lies in the face } F_\sigma.$$

A *facet* of Y_{\geq} is a face F_σ corresponding to a one-dimensional cone σ . The real toric variety Y is smooth at points corresponding to facets, but may not be smooth along lower-dimensional faces. If Y_{\geq}° is the union of the dense face F_0 and its facets, then

$$Y^\circ := (Y_{\geq}^\circ \times \{\pm 1\}^n) / \sim$$

consists of smooth points of Y .

We generalize this construction. Let P be a finite ranked poset with minimal element 0 and rank at most n where two elements $\sigma, \tau \in P$ have at most one minimal upper bound in P . The cones σ in a rational fan in \mathbb{R}^n form such a poset. Suppose further that we have a collection $\mathcal{S} := \{\overline{\sigma} \mid \sigma \in P\}$ of subgroups of $\{\pm 1\}^n$ where $\overline{\sigma} \simeq \{\pm 1\}^{\text{rank}(\sigma)}$, and if $\sigma \subset \tau$, then $\overline{\sigma} \subset \overline{\tau}$. Finally, suppose that we have a cell complex Δ with cells (called faces) indexed by elements of P ,

$$\Delta = \coprod_{\sigma \in P} F_\sigma,$$

where each face F_σ is a cell of dimension $n - \text{rank}(\sigma)$, which we identify with the interior of the closed unit ball in $\mathbb{R}^{n-\text{rank}(\sigma)}$. We further suppose that:

- Δ is a subset of the closed ball $\overline{F_0}$ in \mathbb{R}^n ,
- the closure of a face F_σ in \mathbb{R}^n is homeomorphic to the closed ball of dimension $n - \text{rank}(\sigma)$, and
- given $\sigma, \tau \in P$, the closures of the faces F_σ and F_τ either do not meet (if σ and τ have no upper bound in P), or their intersection is the closure of the face F_ρ , where ρ is the least upper bound of σ and τ in P .

Definition 2.2 Given a ranked poset P , system \mathcal{S} of subgroups of $\{\pm 1\}^n$, and a cell complex Δ as above, the *small cover* $Y(\Delta, \mathcal{S})$ of Δ is the quotient

$$(\Delta \times \{\pm 1\}^n) / \sim,$$

where $(p, \xi) \sim (q, \eta)$ if and only if $p = q$ and $\xi\bar{\sigma} = \eta\bar{\sigma}$, where p lies in the face F_σ .

Observe that $Y(\Delta, \mathcal{S})$ is equipped with a natural action of $\{\pm 1\}^n$ whose orbit space is Δ , where the orbit of a face F_σ is identified with $F_\sigma \times \{\pm 1\}^n / \bar{\sigma} \simeq \mathbb{T}^{n-\text{rank}(\sigma)}$. In particular, it is a $\{\pm 1\}^n$ -equivariant compactification of \mathbb{T}^n .

A real toric variety Y associated to a fan Σ is a small cover where P is the set of cones in the fan, $\Delta = Y_{\geq}$, and $\mathcal{S} = \{\bar{\sigma} \mid \sigma \in \Sigma\}$.

The points of $Y(\Delta, \mathcal{S})$ corresponding to the big cell F_0 and to facets F_σ are points where $Y(\Delta, \mathcal{S})$ is a topological manifold. Write Δ° for the union of the big cell and the facets, and $Y^\circ(\Delta, \mathcal{S}) = (\Delta^\circ \times \{\pm 1\}^n) / \sim$ for this subset of the smooth points of $Y(\Delta, \mathcal{S})$.

Let $\Delta \subset \mathbb{R}^n$ be a n -dimensional polytope with integer vertices and normal fan Σ . Then the real toric variety Y_Σ associated to Σ has a projective embedding given by Δ . We may assume that the integer points $\Delta \cap \mathbb{Z}^n$ generate \mathbb{Z}^n . Let \mathbb{P}^Δ be the real projective space with coordinates indexed by $\Delta \cap \mathbb{Z}^n$ and $y^\alpha := y_1^{\alpha_1} \cdots y_n^{\alpha_n}$ the monomial with exponent α . Then we have an injection

$$\varphi_\Delta : \mathbb{T}^n \ni y \mapsto [y^\alpha \mid \alpha \in \Delta \cap \mathbb{Z}^n], \tag{2.1}$$

where $[\cdots]$ denotes homogeneous coordinates for \mathbb{P}^Δ . The closure Y_Δ of the image of this map is isomorphic to the real toric variety Y_Σ , and the cell complex Y_{\geq}° is identified with the polytope Δ .

The unit sphere $\mathbb{S}^\Delta \subset \mathbb{R}^\Delta$ has a two-to-one map to \mathbb{P}^Δ , and we define Y_Δ^+ to be the pullback of Y_Δ along this map. The sphere \mathbb{S}^Δ has homogeneous coordinates $(x_\alpha \mid \alpha \in \Delta \cap \mathbb{Z}^n)$, where we identify points with a positive constant of proportionality. The group $\{\pm 1\}^{n+1}$ acts on \mathbb{S}^Δ with the last coordinate acting through global multiplication by ± 1 and the remaining coordinates $\{\pm 1\}^n$ through the map φ_Δ (2.1),

$$(g, g_{n+1}) \cdot (x_\alpha \mid \alpha \in \Delta \cap \mathbb{Z}^n) = (g_{n+1}g^\alpha x_\alpha \mid \alpha \in \Delta \cap \mathbb{Z}^n).$$

The faces of Y_Δ^+ are its intersections with coordinate subspaces \mathbb{S}^F of \mathbb{S}^Δ corresponding to faces F of Δ ,

$$\mathbb{S}^F := \{(x_\alpha \mid \alpha \in \Delta \cap \mathbb{Z}^n) \mid x_\alpha = 0 \text{ if } \alpha \notin F \cap \mathbb{Z}^n\}.$$

The isotropy subgroup of \mathbb{S}^F is

$$\{(g, g_{n+1}) \mid g_{n+1}g^\alpha = 1 \text{ for } \alpha \in F \cap \mathbb{Z}^n\}.$$

Vectors b in the normal cone σ_F to a face F of Δ have constant dot product with elements of F —define $b \cdot F$ to be this constant. Then the subgroup

$$\bar{\sigma}_F^+ := \{(-1)^{(b, b \cdot F)} \mid b \in \sigma_F\} \subset \{\pm 1\}^{n+1}$$

is the isotropy group of \mathbb{S}^F , and therefore of the corresponding face of Y_Δ^+ .

Proposition 2.3 *The spherical toric variety Y_Δ^+ is obtained as the quotient of the cell complex $\Delta \times \{\pm 1\}^{n+1}$ by the equivalence relation*

$$(p, \xi) \sim (q, \eta) \iff p = q \text{ and } \xi \bar{\sigma}_F^+ = \eta \bar{\sigma}_F^+, \text{ where } p \text{ lies in the face } F.$$

3 Characterization of Orientability

We follow Nakayama and Nishimura [15] to characterize the orientability of a general small cover and of spherical toric varieties, and determine their numbers of components.

Theorem 3.1 *Let $Y(\Delta, S)$ be a small cover of dimension n .*

- (1) $Y^\circ(\Delta, S)$ is orientable if and only if there exists a basis of $\{\pm 1\}^n$ such that for every $\sigma \in P$ of rank 1, the generator of $\bar{\sigma} \simeq \{\pm 1\}$ is a product of an odd number of basis vectors.
- (2) The components of $Y^\circ(\Delta, S)$ are naturally indexed by $\{\pm 1\}^n / \langle \bar{\sigma} \mid \text{rank}(\sigma) = 1 \rangle$.

Thus $Y^\circ(\Delta, S)$ has 2^{n-k} connected components, where $2^k = |\langle \bar{\sigma} \mid \text{rank}(\sigma) = 1 \rangle|$.

Proof For each $\sigma \in P$ with rank 1, let g_σ be the generator of $\bar{\sigma} \simeq \mathbb{Z}/2\mathbb{Z}$. Then $Y^\circ := Y^\circ(\Delta, S)$ is obtained by gluing (Δ, ξ) and (Δ, η) along F_σ whenever $\xi = \eta g_\sigma$ for some $\sigma \in P$ of rank 1, so the connected components of Y° correspond to the orbits of Y° under the action of $\langle \bar{\sigma} \mid \text{rank}(\sigma) = 1 \rangle$.

The space Y° is orientable if and only if $H_n(Y^\circ, \mathbb{Z}) \neq \{0\}$. This group is the kernel $\ker \partial$ of the differential in the cellular chain complex of the cell complex Y° ,

$$C_n \xrightarrow{\partial} C_{n-1}.$$

Here C_n is the free abelian group generated by

$$\{\Delta\} \times \{\pm 1\}^n = \{(\Delta, \xi) \mid \xi \in \{\pm 1\}^n\}$$

and C_{n-1} is the free abelian group generated by

$$\{[F_\sigma, \xi] \mid \sigma \in P, \text{rank}(\sigma) = 1, \xi \in \{\pm 1\}^n\} / \sim,$$

where $[F_\sigma, \xi] \sim [F_\sigma, \xi g_\sigma]$. Orient each facet F_σ so that

$$\partial(\Delta) = \sum_{\text{rank}(\sigma)=1} F_\sigma.$$

Consider an n -cycle

$$X = \sum_{\xi \in \{\pm 1\}^n} n_\xi \cdot (\Delta, \xi) \in C_n$$

on Y° , where $n_\xi \in \mathbb{Z}$. Then

$$\partial(X) = \sum_{\xi \in \{\pm 1\}^n} n_\xi \sum_{\text{rank}(\sigma)=1} [F_\sigma, \xi] = \sum_{\text{rank}(\sigma)=1} \sum_{\xi \in \{\pm 1\}^n / \langle g_\sigma \rangle} (n_\xi + n_{\xi g_\sigma}) [F_\sigma, \xi].$$

Hence an n -cycle X lies in $\ker \partial$ if and only if $n_\xi = -n_{\xi g_\sigma}$ for all ξ in $\{\pm 1\}^n$ and σ of rank 1. Equivalently, $n_\xi = (-1)^k n_{\xi g_{\sigma_1} \dots g_{\sigma_k}}$ for all $\xi \in \{\pm 1\}^n$ and σ_i of rank 1.

We show that $\ker \partial$ is non-trivial if and only if there exists a basis e_1, \dots, e_n of $\{\pm 1\}^n$ such g_σ is a product of an odd number of basis vectors, for each element $\sigma \in P$ of rank one. Let \mathbb{O} be the set of generators g_σ of $\bar{\sigma}$ for rank one elements $\sigma \in P$.

Suppose that there exists a basis e_1, \dots, e_n of $\{\pm 1\}^n$ such that each $g_\sigma \in \mathbb{O}$ is a product of an odd number of basis vectors. For $\xi \in \{\pm 1\}^n$ define n_ξ to be 1 if ξ is a product of an even number of the e_i and -1 if it is a product of an odd number of the e_i . Then $n_\xi = -n_{\xi g_\sigma}$ for all ξ and σ , so $\ker \partial$ is non-trivial and hence Y° is orientable. Since the number of connected components is 2^{n-k} , the kernel is isomorphic to $\mathbb{Z}^{2^{n-k}}$.

If there is no such basis of $\{\pm 1\}^n$, then there is some $g_\sigma \in \mathbb{O}$ which is a product of an even number of other elements in \mathbb{O} , for otherwise we can reduce \mathbb{O} to a linearly independent set and then extend it to a basis of $\{\pm 1\}^n$. We get $g_\sigma = g_{\sigma_1} \dots g_{\sigma_{2k}}$ and hence $1 = g_\sigma g_{\sigma_1} \dots g_{\sigma_{2k}}$, so for every ξ we get

$$n_\xi = (-1)^{2k+1} n_{\xi g_\sigma g_{\sigma_1} \dots g_{\sigma_{2k}}} = -n_\xi,$$

which implies that $n_\xi = 0$ and hence $\ker \partial = 0$ and so Y° is non-orientable. □

We restate the orientability criteria of Theorem 3.1 for real toric varieties.

Theorem 3.2 *Let Y be a real toric variety defined by a fan Σ . Then Y° is orientable if and only if there exists a basis of $\{\pm 1\}^n$ such that $(-1)^b$ is a product of an odd number of basis vectors, for each primitive vector b lying on a ray of Σ .*

The condition of Theorem 3.2 is easily checked.

Lemma 3.3 *Given $A \subset \{\pm 1\}^n$, the condition that there exists a basis of $\{\pm 1\}^n$ such that each vector in A is a product of an odd number of basis vectors, is equivalent to the condition that no product of an odd number of vectors in A is equal to 1 in $\{\pm 1\}^n$.*

Proof If we had $v_1 \cdots v_{2k+1} = 1$, then $v_{2k+1} = v_1 \cdots v_{2k}$, and expressing each v_i as the product of an odd number of basis elements of $\{\pm 1\}^n$ yields a contradiction. For the other implication, reduce A to a linearly independent set A' and then extend A' to a basis of $\{\pm 1\}^n$. If there were a vector in $A \setminus A'$ which is a product of an even number of vectors $v = v_1 \cdots v_{2k}$, we would have then had $v \cdot v_1 \cdots v_{2k} = 1$. \square

We may check if the condition is satisfied by reducing A to a linearly independent set A' and checking if each vector in $A \setminus A'$ is a product of an odd number vectors in A' .

The analog of Theorem 3.1 for spherical toric varieties has a similar proof.

Theorem 3.4 *Let $Y_\Delta^+ \subset \mathbb{S}^\Delta$ be a spherical toric variety defined by a full-dimensional lattice polytope $\Delta \subset \mathbb{R}^n$.*

- (1) Y_Δ^+ is orientable if and only if there exists a basis of $\{\pm 1\}^{n+1}$ such that for each facet F of Δ with primitive normal vector b , the element $(-1)^{(b, b \cdot F)}$ is a product of an odd number of basis elements.
- (2) The components of Y_Δ^+ are naturally indexed by

$$\{\pm 1\}^{n+1} / \langle (-1)^{(b, b \cdot F)} \mid b \text{ is a primitive normal vector to a facet } F \text{ of } \Delta \rangle.$$

4 Examples and Applications to Lower Bounds

We settle questions of orientability left open in [20] and explain our motivation from the study of real solutions to systems of polynomials. We begin with an example.

4.1 Cross Polytopes

The cross polytope is the convex hull of the basis vectors e_1, \dots, e_n in \mathbb{R}^n and their negatives $-e_1, \dots, -e_n$. When $n > 1$ the corresponding toric variety is singular. The rays of its normal fan have generators $(\pm 1, \dots, \pm 1)$, all with the same image in $(\mathbb{Z}/2\mathbb{Z})^n$. The hypotheses of Theorem 3.1 hold, and so the corresponding real toric variety is orientable and its smooth points have 2^{n-1} connected components. Figure 1 displays the cross polytope when $n = 2$ and an embedding in \mathbb{R}^3 of the corresponding real toric variety.

This example was treated in detail in [21, § 7].

4.2 Order Polytopes

The *order polytope* $O(P)$ [22] of a finite poset P is

$$O(P) := \{y \in [0, 1]^P \mid a \leq b \text{ in } P \Rightarrow y_a \leq y_b\}.$$




Fig. 1 Two-dimensional cross polytope and double pillow

The integer points of $O(P)$ are its vertices and they correspond to the order ideals of P .

Theorem 4.1 $Y_{O(P)}$ is orientable if and only if all maximal chains of P have odd length.

Proof Lemma 4.9 of [20] (or rather its proof) implies that $Y_{O(P)}$ is orientable if all maximal chains of P have odd length. We establish the converse.

The order polytope has three types of facets

$$\begin{aligned} y_a &= 0 && \text{for } a \in P \text{ minimal,} \\ y_b &= 1 && \text{for } b \in P \text{ maximal,} \\ y_b - y_a &= 0 && \text{for } b \text{ covering } a \text{ (} a < b \text{) in } P. \end{aligned}$$

Replacing $=$ by \geq gives valid inequalities for $O(P)$, which we write in matrix form

$$O(P) := \{y \in \mathbb{R}^P \mid \mathcal{A}y \geq c\}. \tag{4.1}$$

By Theorem 3.2, $Y_{O(P)}$ is orientable if and only if there is a basis of the row space of \mathcal{A} , reduced modulo 2, such that each row is a sum of an odd number of basis vectors.

Fix a maximal chain $a_1 < \dots < a_k$ in P . The corresponding facets of $O(P)$ are

$$y_{a_1} = 0, \quad y_{a_2} - y_{a_1} = 0, \quad \dots, \quad y_{a_k} - y_{a_{k-1}} = 0, \quad y_{a_k} = 1,$$

and the corresponding rows of the matrix \mathcal{A} (modulo 2) are

$$\begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 & 0 & 0 & \dots & 0 \\ 0 & 1 & 1 & \dots & 0 & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 1 & 1 & 0 & \dots & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 & 0 & \dots & 0 \end{bmatrix},$$

where the non-zero columns correspond to a_1, \dots, a_k . This gives $k+1$ rows whose sum is zero modulo 2. If k is even, Lemma 3.3 implies that $Y_{O(P)}$ is non-orientable. \square

A poset P is *ranked modulo 2* if all maximal chains in P have the same parity.

Theorem 4.2 *A spherical toric variety $Y_{O(P)}^+$ is orientable if and only if P is ranked modulo 2.*

Proof By Lemma 4.9 of [20], $Y_{O(P)}^+$ is orientable if it is ranked modulo 2.

Suppose that P is not ranked modulo 2. We exhibit an odd number of rows of the augmented matrix $[\mathcal{A} : c]$ whose sum is zero modulo 2, which shows that $Y_{O(P)}^+$ is not orientable, by Theorem 3.4 and Lemma 3.3, as these rows have the form $(b, b \cdot F)$ for b a primitive normal to a facet of the order polytope.

The order polytope is defined by the facet inequalities (4.1). For a maximal chain $a_1 \prec \dots \prec a_k$ in P , the corresponding rows of the augmented matrix $[\mathcal{A} : c]$ are

$$\begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & \dots & 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 1 & \dots & 0 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 1 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 & 0 & \dots & 0 & 1 \end{bmatrix}.$$

Observe that the sum of these rows is $[0 : 1]$. Each row of $[\mathcal{A} : c]$ has the form $(b, b \cdot F)$ (modulo 2), where b is a primitive normal vector to a facet F of Δ .

Since P is not ranked modulo 2, it has two maximal chains of different parities. Summing the rows of $[\mathcal{A} : c]$ which correspond to facets given by the two chains gives a sum of an odd number of rows of $[\mathcal{A} : c]$ which is equal to zero modulo 2. \square

4.3 Real Solutions to Systems of Equations

In [20] we considered systems,

$$f_1(x_1, \dots, x_n) = f_2(x_1, \dots, x_n) = \dots = f_n(x_1, \dots, x_n) = 0, \tag{4.2}$$

where each f_i is a real polynomial whose exponent vectors lie in $\Delta \cap \mathbb{Z}^n$, for a fixed lattice polytope Δ , called the *Newton polytope* of the system. When the exponent vectors $\Delta \cap \mathbb{Z}^n$ affinely span \mathbb{Z}^n , the solutions to (4.2) correspond to a linear section $L \cap Y_\Delta$ of the real projective toric variety Y_Δ corresponding to Δ . Here $L \subset \mathbb{RP}^\Delta$ is a linear subspace of codimension n . Projecting from a general codimension one linear subspace E of L , we may realize the solutions to (4.2) as the fibers of a map

$$\pi_E : Y_\Delta \longrightarrow \mathbb{RP}^n,$$

to real projective space. If n is odd, then \mathbb{RP}^n is orientable. If Y_Δ is also orientable, then fixing orientations, the map π_E has a degree whose absolute value gives a lower

bound on the cardinality of a fiber of π_E , and thus on the number of real solutions to (4.2).

More generally, we may lift this projection to the spherical toric varieties

$$\pi_E^+ : Y_\Delta^+ \longrightarrow \mathbb{S}^n. \quad (4.3)$$

If Y_Δ^+ is orientable, we fix an orientation and the absolute value of the degree of π_E^+ is a lower bound on the number of solutions to the system (4.2). Changing orientations in each component if necessary, we may assume that the degree is divisible by the number of components of $(Y_\Delta^+)^{\circ}$.

This has the following consequence for lower bounds to systems of polynomial equations.

Theorem 4.3 *Suppose that we have a system of polynomials (4.2) with Newton polytope Δ where $\Delta \cap \mathbb{Z}^n$ affinely spans \mathbb{Z}^n whose solutions are a fiber of a projection map π_E^+ (4.3). If there is a basis for $\{\pm 1\}^{n+1}$ such that $(-1)^{(b, b \cdot F)}$ is a product of an odd number of basis elements for every primitive normal vector b to a facet F of Δ , then the absolute value of the degree of the map π_E^+ is a lower bound for the number of real solutions to (4.2), and this lower bound is a multiple of the number of components of $(Y_\Delta^+)^{\circ}$.*

Moreover, the map π_E^+ does not have a degree if this condition is not satisfied.

Remark 4.4 We did not need to consider the parity of n , for the condition of Theorem 3.2 implies that of Theorem 3.4. (A vector lies in a ray of the normal fan Σ to Δ if and only if it is normal to a facet F of Δ .)

4.4 Conclusions

We characterized the orientability of Y_Δ and Y_Δ^+ , which implies that the corresponding polynomial system has lower bounds on its number of real solutions, expressed as the degree of a projection π_E or π_E^+ . These degrees have been computed for polynomial systems from posets [20] and those from foldable triangulations [12, 13, 20]. Our characterization of orientability replaces the condition in [20] that a variety is Cox-oriented and therefore strengthens the results of [20], particularly Theorem 3.5.

Acknowledgments The research of Sottile was supported in part by NSF grant DMS-1001615, and the Mittag-Leffler Institut.

References

1. Azar, M., Gabrielov, A.: Lower bounds in B. and M. Shapiro conjecture. *Discrete Comput. Geom.* **46**(4), 636–659 (2011)
2. Craciun, G., Dickenstein, A., Shiu, A., Sturmfels, B.: Toric dynamical systems. *J. Symb. Comput.* **44**(11), 1551–1565 (2009)
3. Davis, M.W., Januszkiewicz, T.: Convex polytopes, Coxeter orbifolds and torus actions. *Duke Math. J.* **62**(2), 417–451 (1991)
4. Degtyarev, A.I., Kharlamov, V.M.: Topological properties of real algebraic varieties: Rokhlin's way. *Uspekhi Mat. Nauk* **55**(4(334)), 129–212 (2000)

5. Delaunay, C.: Real structures on compact toric varieties. Ph.D. thesis, Université Louis Pasteur, Strasbourg (2004)
6. Eremenko, A., Gabrielov, A.: Degrees of real Wronski maps. *Discrete Comput. Geom.* **28**(3), 331–347 (2002)
7. Finashin, F., Kharlamov, V.: Abundance of real lines on real projective hypersurfaces. *Int. Math. Res. Notices* (2012). doi:[10.1093/imrn/rns135](https://doi.org/10.1093/imrn/rns135)
8. Fulton, W.: Introduction to Toric Varieties. *Annals of Mathematics Studies*, vol. 131. Princeton University Press, Princeton (1993)
9. García-Puente, L., Hein, N., Hillar, C., Martín del Campo, A., Ruffo, J., Sottile, F., Teitler, Z.: The secant conjecture in the real Schubert calculus. *Exp. Math.* **21**(3), 252–265 (2012)
10. Hein, N., Sottile, F.: Lower bounds in real Schubert calculus. <http://arXiv.org/1211.7160> (2013)
11. Hein, N., Sottile, F., Zelenko, I.: A congruence modulo four in real Schubert calculus. <http://arXiv/abs/1211.7160> (2012)
12. Joswig, M., Witte, N.: Products of foldable triangulations. *Adv. Math.* **210**(2), 769–796 (2007)
13. Joswig, M., Ziegler, G.: Foldable triangulations of lattice polygons. *Am. Math. Mon.* to appear. <http://arXiv/abs/1207.6865> (2012)
14. Krasauskas, R.: Toric surface patches: advances in geometrical algorithms and representations. *Adv. Comput. Math.* **17**(1–2), 89–113 (2002)
15. Nakayama, H., Nishimura, Y.: The orientability of small covers and coloring simple polytopes. *Osaka J. Math.* **42**(1), 243–256 (2005)
16. Okonek, Ch., Teleman, A.: Intrinsic signs and lower bounds in real algebraic geometry. *J. Reine Angew. Math.* to appear. <http://arXiv/abs/1112.3851> (2012)
17. Okonek, Ch., Teleman, A.: A wall crossing formula for degrees of real central projections. <http://arXiv/abs/1206.4271> (2012)
18. Pachter, L., Sturmfels, B. (eds.): *Algebraic Statistics for Computational Biology*. Cambridge University Press, New York (2005)
19. Ruffo, J., Sivan, Y., Soprunova, E., Sottile, F.: Experimentation and conjectures in the real Schubert calculus for flag manifolds. *Exp. Math.* **15**(2), 199–221 (2006)
20. Soprunova, E., Sottile, F.: Lower bounds for real solutions to sparse polynomial systems. *Adv. Math.* **204**(1), 116–151 (2006)
21. Sottile, F.: Toric ideals, real toric varieties, and the moment map. In: *Topics in Algebraic Geometry and Geometric Modeling*. Contemporary Mathematics, vol. 334. American Mathematical Society, Providence, RI, pp. 225–240 (2003)
22. Stanley, R.P.: Two poset polytopes. *Discrete Comput. Geom.* **1**(1), 9–23 (1986)
23. Welschinger, J.-Y.: Invariants of real rational symplectic 4-manifolds and lower bounds in real enumerative geometry. *C. R. Math. Acad. Sci. Paris* **336**(4), 341–344 (2003)