

Erratum to: Prevalence of preoperative alcohol abuse among patients seeking weight-loss surgery

Omar Yusef Kudsi · Karen Huskey ·
Shannon Grove · George Blackburn ·
Daniel B. Jones · Christina C. Wee

Published online: 18 December 2012
© Springer Science+Business Media New York 2012

Erratum to: Surg Endosc
DOI 10.1007/s00464-012-2568-x

On the first page of the article, second paragraph, third line, the response rate should be 70 %, not 75 %.

The same correction to the response rate also applies to the text on the second page, third paragraph, second line.

In addition, C. C. Wee should also be considered as the corresponding author.

The online version of the original article can be found under
doi:[10.1007/s00464-012-2568-x](https://doi.org/10.1007/s00464-012-2568-x).

O. Y. Kudsi · D. B. Jones
Division of Minimally Invasive Surgery, Department of Surgery,
Beth Israel Deaconess Medical Center, Harvard Medical School,
Boston, MA 02215, USA
e-mail: okudsi@bidmc.harvard.edu

K. Huskey · S. Grove · C. C. Wee (✉)
Division of General Medicine and Primary Care, Department of
Medicine, Beth Israel Deaconess Medical Center, Harvard
Medical School, Boston, MA 02215, USA
e-mail: cwee@bidmc.harvard.edu

G. Blackburn
Center for the Study of Nutrition, Beth Israel Deaconess Medical
Center, Harvard Medical School, Boston, MA 02215, USA

C. C. Wee
Division of General Medicine and Primary Care, Beth Israel
Deaconess Medical Center, 330 Brookline Ave, Boston,
MA 02215, USA