



Correction to: Clinical practice recommendations for the diagnosis and management of Alport syndrome in children, adolescents, and young adults—an update for 2020

Clifford E. Kashtan¹ · Oliver Gross²

Published online: 12 January 2021

© IPNA 2021

Correction to: *Pediatr Nephrol*

<https://doi.org/10.1007/s00467-020-04819-6>

In our recent paper “Clinical Practice Recommendations for the Diagnosis and Management of Alport Syndrome in Children, Adolescents and Young Adults – An Update for 2020” [1] the definition of microalbuminuria is misstated as a urine microalbumin-creatinine ratio of 30 mg/mg. The correct value is 30 mg/g. We apologize for this error.

References

1. Kashtan CE, Gross O (2020) Clinical practice recommendations for the diagnosis and management of Alport syndrome in children, adolescents, and young adults-an update for 2020. *Pediatr Nephrol*. <https://doi.org/10.1007/s00467-020-04819-6>.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00467-020-04819-6>

✉ Clifford E. Kashtan
kash001@umn.edu

¹ Department of Pediatrics, Division of Pediatric Nephrology, University of Minnesota Medical School, 2450 Riverside Avenue, Minneapolis, MN 55454, USA

² Department of Nephrology and Rheumatology, University Medical Center Goettingen, Goettingen, Germany