

Erratum to: Potential volume for CO₂ deep ocean sequestration: an assessment of the area located on western Pacific Ocean

David Ching-Fang Shih¹ · Yih-Min Wu² · Jyr-Ching Hu²

Published online: 25 November 2015
© Springer-Verlag Berlin Heidelberg 2015

**Erratum to: Stoch Environ Res Risk Assess (2010)
24:705–711
DOI 10.1007/s00477-009-0357-4**

Two mistakes require to be modified.

1. The last sentence in **Abstract** section is corrected as
“On the other hand, if it accounts for the bottom of the sea floor as the reference and the accumulated CO₂ is stored from the depth at -4900 m using bottom-up

- style, it requires about 37 and 61 m for the storage of 20 years and a century respectively.”
2. The last sentence in **Conclusions** section is corrected as
“On the other hand, if CO₂ storage can be conducted from the sea bottom using bottom-up style at the reference depth at -4900 m, it requires about 37 and 61 m for the storage of 20 and 100 years respectively.”

The online version of the original article can be found under
doi:[10.1007/s00477-009-0357-4](https://doi.org/10.1007/s00477-009-0357-4).

✉ David Ching-Fang Shih
cfshih@iner.gov.tw

¹ Institute of Nuclear Energy Research, AEC,
P.O. Box 3-7, Longtan 32546, Taiwan, ROC

² Department of Geosciences, National Taiwan University,
Taipei 10617, Taiwan, ROC