

## Erratum to: Prevalence and factors associated with polypharmacy in older people with cancer

Justin P. Turner · Sepehr Shakib · Nimit Singhal ·  
Jonathon Hogan-Doran · Robert Prowse · Sally Johns ·  
J. Simon Bell

Published online: 19 April 2014  
© Springer-Verlag Berlin Heidelberg 2014

**Erratum to: Support Care Cancer**  
**DOI 10.1007/s00520-014-2171-x**

The original version of this paper unfortunately contains errors, the fourth sentence of the second paragraph in the results section incorrectly reads as:

“The most prevalent comorbidities were hypertension (62.0 %), hypercholesterolemia (34.4 %), arthritis (31.1 %), diabetes (27.0 %), airways diseases (26.2 %), gastrointestinal disorders (22.0 %) and coronary artery disease (17.9 %).”

Where it should have read as:

“The most prevalent comorbidities were hypertension (58.4 %), hypercholesterolemia (32.5 %), arthritis (29.4 %), diabetes (25.5 %), airways diseases (24.6 %), gastrointestinal disorders (22.0 %) and coronary artery disease (16.9 %).”

---

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00520-014-2171-x>.

---

J. P. Turner · J. S. Bell  
School of Pharmacy and Medical Sciences, University of South  
Australia, Adelaide, South Australia, Australia

J. P. Turner (✉) · J. S. Bell  
Centre for Medicine Use and Safety, Faculty of Pharmacy and  
Pharmaceutical Sciences, Monash University, 381 Royal Parade,  
Parkville, 3052 Melbourne, Victoria, Australia  
e-mail: [justin.turner@unisa.edu.au](mailto:justin.turner@unisa.edu.au)

S. Shakib · S. Johns  
Department of Clinical Pharmacology, Royal Adelaide Hospital  
Adelaide, Adelaide, South Australia, Australia

N. Singhal · J. Hogan-Doran  
University of Adelaide and Department of Medical Oncology, Royal  
Adelaide Hospital, Adelaide, South Australia, Australia

R. Prowse  
Department of Geriatric and Rehabilitation Medicine, Royal  
Adelaide Hospital, Adelaide, South Australia, Australia