

## Erratum to: Hemodynamic effect of the left gastric artery on esophageal varices in patients with cirrhosis

Soichiro Kiyono<sup>1</sup> · Hitoshi Maruyama<sup>1</sup> · Takayuki Kondo<sup>1</sup> · Tadashi Sekimoto<sup>1</sup> · Taro Shimada<sup>1</sup> · Masanori Takahashi<sup>1</sup> · Osamu Yokosuka<sup>1</sup>

Published online: 17 August 2016  
© Japanese Society of Gastroenterology 2016

### Erratum to: J Gastroenterol DOI 10.1007/s00535-015-1157-x

The author would like to correct the errors in the publication of the original article. The corrected details are given below for your reading.

In the “Abstract” section, the second sentence beneath the heading “Results” should read as

EV were demonstrated in 45/45 patients, with a mean detection time of 6.9 s (2–21), which was longer in patients with variceal recurrence (7.0 s) than in those without (5.6 s,  $P = 0.048$ ).

Secondly, the second sentence in the page 6 beneath the heading “Relationship between the LGA findings and post-treatment variceal recurrence” should read as

The mean detection time was significantly longer in patients with recurrence (7.0 s) than in those without (5.6 s,  $P = 0.048$ ).

---

The online version of the original article can be found under doi:[10.1007/s00535-015-1157-x](https://doi.org/10.1007/s00535-015-1157-x).

---

✉ Hitoshi Maruyama  
maru-cib@umin.ac.jp

<sup>1</sup> Department of Gastroenterology and Nephrology, Graduate School of Medicine, Chiba University, Chiba, Japan