


Correction to: Telomere length of gallbladder epithelium is shortened in patients with congenital biliary dilatation: measurement by quantitative fluorescence in situ hybridization

Yuto Aoki^{1,2}  · Junko Aida² · Youichi Kawano^{2,3} · Ken-ichi Nakamura² · Naotaka Izumiyama-Shimomura² · Naoshi Ishikawa² · Tomio Arai⁴ · Yoshiharu Nakamura¹ · Nobuhiko Taniai¹ · Eiji Uchida¹ · Kaiyo Takubo² · Toshiyuki Ishiwata²

Published online: 5 December 2017
© Japanese Society of Gastroenterology 2017

Correction to: J Gastroenterol

<https://doi.org/10.1007/s00535-017-1411-5>

In the original publication of this article, Fig. 5 was published with incorrect color of the approximate lines of CBD and Control.

The corrected figure is given in the following page.

The original article can be found online at <https://doi.org/10.1007/s00535-017-1411-5>.

✉ Yuto Aoki
yutoaoki@nms.ac.jp

✉ Toshiyuki Ishiwata
tishiwat@tmig.or.jp

¹ Department of Gastrointestinal and Hepato-Biliary-Pancreatic Surgery, Nippon Medical School, 1-1-5 Sendagi, Bunkyo-ku, Tokyo 113-8603, Japan

² Research Team for Geriatric Pathology and Department of Pathology, Tokyo Metropolitan Geriatric Hospital and Institute of Gerontology, Tokyo, Japan

³ Department of Surgery, Nippon Medical School Chiba Hokusoh Hospital, Chiba, Japan

⁴ Department of Pathology, Tokyo Metropolitan Geriatric Hospital, Tokyo, Japan

