

## Erratum to: The efficacy of continuous subcostal transversus abdominis plane block for analgesia after living liver donation: a retrospective study

Akihiko Maeda<sup>1,2</sup>  · Sho Carl Shibata<sup>2</sup> · Hiroshi Wada<sup>3</sup> · Shigeru Marubashi<sup>3</sup> · Takahiko Kamibayashi<sup>2</sup> · Hidetoshi Eguchi<sup>3</sup> · Yuji Fujino<sup>2</sup>

Published online: 16 February 2016  
© Japanese Society of Anesthesiologists 2016

**Erratum to: J Anesth (2016) 30:39–46**  
**DOI 10.1007/s00540-015-2085-x**

The authors noticed an error in the last sentence of the second paragraph in the ‘Result’ section of the article, and would like to correct it as follows:

“The administration of nonopioid analgesics (Table 2), PT-INR, and duration of postoperative hospital stay (Table 3) did not differ between the groups.”

---

The online version of the original article can be found under doi:[10.1007/s00540-015-2085-x](https://doi.org/10.1007/s00540-015-2085-x).

---

✉ Akihiko Maeda  
amaeda@anes.med.osaka-u.ac.jp

- <sup>1</sup> Present Address: Department of Anesthesiology, National Hospital Organization Osaka National Hospital, 2-1-14 Hoenzaka, Chuo-ku, Osaka, Osaka 540-0006, Japan
- <sup>2</sup> Department of Anesthesiology and Intensive Care Medicine, Graduate School of Medicine, Osaka University, Suita, Osaka, Japan
- <sup>3</sup> Department of Gastroenterological Surgery, Graduate School of Medicine, Osaka University, Suita, Osaka, Japan