



## Correction to: Attenuation of cortical activity triggering descending pain inhibition in chronic low back pain patients: a functional magnetic resonance imaging study

Yohei Matsuo<sup>1</sup> · Jiro Kurata<sup>2</sup>  · Miho Sekiguchi<sup>1</sup> · Katsuhiko Yoshida<sup>1</sup> · Takuya Nikaido<sup>1</sup> · Shin-ichi Konno<sup>1</sup>

Published online: 19 January 2018  
© Japanese Society of Anesthesiologists 2018

**Correction to: J Anesth (2017) 31:523–530**  
<https://doi.org/10.1007/s00540-017-2343-1>

Inadvertently, the Fig. 7 was published incorrectly in the original publication of the article. The correct Fig. 7 should be as below:

---


The original article can be found online at <https://doi.org/10.1007/s00540-017-2343-1>.

---

✉ Jiro Kurata  
jkurata@plum.plala.or.jp

<sup>1</sup> Department of Orthopaedic Surgery, Fukushima Medical University School of Medicine, 1 Hikariga-oka, Fukushima City 960-1295, Japan

<sup>2</sup> Department of Anesthesiology and Pain Clinic, Tokyo Medical and Dental University Hospital of Medicine, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan



**Fig. 7** Correlations between the MPQ score and %BOLD signal changes. Activations at both the ACC and DLPFC were negatively correlated with the MPQ scores. *MPQ* scores of the McGill Pain

Questionnaire Short Form, *CP* chronic pain patients, *HC* healthy control subjects, *L* left, *ACC* anterior cingulate cortex, *DLPFC* dorsolateral prefrontal cortex