


Erratum to: Long-term results of surgery for lumbar spinal stenosis: a randomised controlled trial

Pär Slätis · Antti Malmivaara · Markku Heliövaara ·
Päivi Sainio · Arto Herno · Jyrki Kankare · Seppo Seitsalo ·
Kaj Tallroth · Veli Turunen · Paul Knekt · Heikki Hurri

Published online: 5 August 2011
© Springer-Verlag 2011

Erratum to: Eur Spine J (2011) 20:1174–1181 DOI 10.1007/s00586-010-1652-y

Figure 3 indicating the magnitude of perceived leg pain during the follow up period is inadvertently a copy of Fig. 2 in the same article. The correct Fig. 3 is attached here and should replace the published Fig. 3.



The online version of the original article can be found under
doi:[10.1007/s00586-010-1652-y](https://doi.org/10.1007/s00586-010-1652-y).

P. Slätis · S. Seitsalo · K. Tallroth · H. Hurri
ORTON Orthopaedic Hospital, Invalid Foundation,
Helsinki, Finland

A. Malmivaara (✉)
Centre for Health Economics,
National Institute for Health and Welfare,
P.O. Box 30, 00271 Helsinki, Finland
e-mail: antti.malmivaara@thl.fi

M. Heliövaara · P. Sainio · P. Knekt
National Institute for Health and Welfare,
P.O. Box 30, 00271 Helsinki, Finland

A. Herno · V. Turunen
Department of Orthopaedic Surgery,
Kuopio University Hospital, Kuopio, Finland

J. Kankare
Department of Orthopaedic and Trauma Surgery (JK),
Helsinki University Hospital, Helsinki, Finland