



Correction to: Staged surgery for tandem cervical and lumbar spinal stenosis: Which should be treated first?

Chi-An Luo¹ · Arun-Kumar Kaliya-Perumal^{1,2} · Meng-Ling Lu³ · Lih-Huei Chen¹ · Wen-Jer Chen^{1,4} · Chi-Chien Niu¹ 

Published online: 3 February 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2020

Correction to: European Spine Journal (2019) 28:61–68
<https://doi.org/10.1007/s00586-018-5795-6>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The authors would like to acknowledge the following funding information that was missing in the original article.

This study was funded by Chang-Gung Memorial Hospital (Grant CORPG3H0261, representing hospital funding). The funding body did not have any role in the design of the study or in data collection, analysis and interpretation or in writing the manuscript.

The original article can be found online at <https://doi.org/10.1007/s00586-018-5795-6>.

✉ Chi-Chien Niu
niuchien@adm.cgmh.org.tw

Chi-An Luo
j5555yrre@gmail.com

Arun-Kumar Kaliya-Perumal
dr.arunkumar.orth@gmail.com

Meng-Ling Lu
lumengling@gmail.com

Lih-Huei Chen
lhchen213216@gmail.com

Wen-Jer Chen
wenjchen4012@gmail.com

¹ Department of Orthopaedic Surgery, Spine Section, Bone and Joint Research Center, Chang Gung Memorial Hospital, Linkou, Chang Gung University College of Medicine, No. 5, Fusing Street, Gueishan, Taoyuan 333, Taiwan

² Department of Orthopaedic Surgery, Melmaruvathur Adhiparasakthi Institute of Medical Sciences and Research, Affiliated to the Tamil Nadu Dr. MGR Medical University, Chennai, Tamil Nadu, India

³ Department of Orthopaedic Surgery, Spine Section, Chang Gung Memorial Hospital, Kaohsiung 833, Taiwan

⁴ Department of Orthopaedic Surgery, Chung Shan Hospital, Taipei 106, Taiwan