

Erratum to: Advantages of FDG-PET/CT over CT alone in the preoperative assessment of lymph node metastasis in patients with esophageal cancer

Ryuichi Karashima · Masayuki Watanabe · Yu Imamura · Satoshi Ida · Yoshifumi Baba · Shiro Iwagami · Yuji Miyamoto · Yasuo Sakamoto · Naoya Yoshida · Hideo Baba

Published online: 18 February 2015
© Springer Japan 2015

Erratum to: Surg Today (2014)
DOI 10.1007/s00595-014-0965-6

Two inaccurate technical descriptions appeared in above titled article.

In the “Imaging studies” subsection, the dose of intravenously injected ^{18}F -FDG was misdescribed as 370 MBq (10 mCi). Actual dose of ^{18}F -FDG was 185–296 MBq (5–8 mCi). In addition, injection rate of nonionic contrast medium during CT imaging was misdescribed as 2.5 ml/s, and accurate injection rate was 2.0 ml/s.

The authors sincerely apologize for these errors.

The online version of the original article can be found under doi:[10.1007/s00595-014-0965-6](https://doi.org/10.1007/s00595-014-0965-6).

R. Karashima · M. Watanabe · Y. Imamura · S. Ida · Y. Baba · S. Iwagami · Y. Miyamoto · Y. Sakamoto · N. Yoshida · H. Baba
Department of Gastroenterological Surgery, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan

M. Watanabe (✉)
Department of Gastroenterological Surgery, Cancer Institute Hospital of Japanese Foundation for Cancer Research,
3-8-31 Ariake, Koto-ku, Tokyo 135-8550, Japan
e-mail: masayuki.watanabe@jfc.or.jp