

## Association between new indices in the locomotive syndrome risk test and decline in mobility: third survey of the ROAD study

Noriko Yoshimura<sup>1</sup> · Shigeyuki Muraki<sup>2</sup> · Hiroyuki Oka<sup>3</sup> · Sakae Tanaka<sup>4</sup> · Toru Ogata<sup>5</sup> · Hiroshi Kawaguchi<sup>6</sup> · Toru Akune<sup>5</sup> · Kozo Nakamura<sup>5</sup>

Received: 5 December 2014 / Accepted: 29 May 2015 / Published online: 25 June 2015  
© The Author(s) 2015. This article is published with open access at Springerlink.com

### Abstract

**Background** We aimed to clarify the association between new indices in a locomotive syndrome risk test and decline in mobility.

**Methods** In the third survey of the Research on Osteoarthritis/osteoporosis Against Disability (ROAD) study, data on the indices were obtained from 1575 subjects (513 men, 1062 women) of the 1721 participants in mountainous and coastal areas. As outcome measures for decline in mobility, we used the five-times-sit-to-stand test (FTSST) and walking speed with cutoff values of 12 s and 0.8 m/s, respectively.

**Electronic supplementary material** The online version of this article (doi:10.1007/s00776-015-0741-5) contains supplementary material, which is available to authorized users.

✉ Noriko Yoshimura  
yoshimuran-ort@h.u-tokyo.ac.jp

<sup>1</sup> Department of Joint Disease Research, 22nd Century Medical and Research Center, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-8655, Japan

<sup>2</sup> Department of Clinical Motor System Medicine, 22nd Century Medical and Research Center, The University of Tokyo, Tokyo 113-8655, Japan

<sup>3</sup> Department of Medical Research and Management for Musculoskeletal Pain, 22nd Century Medical and Research Center, The University of Tokyo, Tokyo 113-8655, Japan

<sup>4</sup> Department of Orthopaedic Surgery, Sensory and Motor System Medicine, Graduate School of Medicine, The University of Tokyo, Tokyo 113-8655, Japan

<sup>5</sup> National Rehabilitation Center for Persons with Disabilities, Saitama 359-0042, Japan

<sup>6</sup> JCHO Tokyo Shinjuku Medical Center, Tokyo 162-8542, Japan

**Results** We first estimated the prevalence of the indices in locomotive syndrome risk test stage 1, including two-step test score <1.3, difficulty with one-leg standing from a 40-cm-high seat in the stand-up test, and 25-question GLFS score  $\geq 7$ , which were found to be 57.4, 40.6, and 22.6 %, respectively. Next, we investigated the prevalence of the indices in locomotive syndrome risk test stage 2, including two-step test score <1.1, difficulty with standing from a 20-cm-high seat using both legs in the stand-up test, and 25-question GLFS score  $\geq 16$ , which were found to be 21.1, 7.9, and 10.6 %, respectively. Logistic regression analysis using slow FTSST time or slow walking speed as the objective factor, and presence or absence of indices as the independent factor, after adjusting for confounders, showed all three indices in both stages 1 and 2 were significantly and independently associated with immobility. Finally, we clarified the risk of immobility according to an increasing number of indices in both stages 1 and 2 and found that the odds ratio for both slow FTSST time and slow walking speed increased exponentially.

**Conclusion** We found that the three indices independently predicted immobility and that accumulation of indices increased the risk of immobility exponentially.

### Introduction

According to the most recent National Livelihood Survey by the Ministry of Health, Labour, and Welfare in Japan, osteoporotic fracture and falls is ranked fourth and osteoarthritis is ranked fifth among conditions that cause disability and subsequently require support with regard to activities of daily living [1]. Given the increasing proportion of elderly individuals in the Japanese population, a comprehensive and evidence-based prevention strategy for musculoskeletal

diseases is urgently required. In 2007, the Japanese Orthopaedic Association (JOA) proposed that the term “locomotive syndrome” should be adopted to designate a condition requiring nursing care, or being at risk of developing such a condition, because of a decline in mobility resulting from a disorder of the locomotive system, which consists of bones, joints, muscles, and nerves [2]. Weakness of such locomotive components causes difficulty in mobility—defined as the ability to stand, walk, run, climb stairs, and perform other physical functions essential to daily life.

As candidate indices to assess the risk of locomotive syndrome, in 2013, the JOA proposed the following three tests: two-step test, stand-up test, and 25-question geriatric locomotive function scale (GLFS) [3]. With regard to the stand-up test, more than 50 % of subjects younger than 70 years old can stand up on one leg from a 40-cm-high seat [3]. The 25-question GLFS has already been assessed regarding its sensitivity and specificity for prediction of disability and was assigned a cutoff value of 16 by Seichi et al. [4]. However, there is little information regarding reference and/or cutoff values for the two-step test.

Recently, the JOA determined clinical decision limits of these three indices for assessing risk of locomotive syndrome [5]. In their proposal, clinical decision limits were established in two stages as follows:

Stage 1:

1. Two-step test score <1.3.
2. Difficulty with one-leg standing from a 40-cm-high seat in the stand-up test (either leg).
3. 25-question GLFS score  $\geq 7$ .

When a subject meets any of the above-mentioned conditions, he/she is diagnosed as starting to decline in mobility.

Stage 2:

1. Two-step test score <1.1.
2. Difficulty with standing from a 20-cm-high seat using both legs in the stand-up test.
3. A 25-question GLFS score  $\geq 16$ .

When a subject meets any of the above-mentioned conditions, he/she is diagnosed as progressing to a decline in mobility.

However, no report has evaluated such indices using data of the general population. From 2005 to 2007, we started a large-scale, population-based cohort investigation entitled the Research on Osteoarthritis/osteoporosis Against Disability (ROAD) study, consisting of 3040 participants in three communities located in urban, mountainous, and

coastal areas. Following the baseline study, we performed a second survey in the same communities from October 2008 to January 2010, followed by a third survey from October 2012 to December 2013. In the third survey, participants completed the two-step test, stand-up test, and 25-question GLFS. In the present report, using data from the third survey of the ROAD study, we assessed the usefulness of these new indices for predicting immobility, which causes subsequent disability.

## Participants and methods

### Participants

Measurements were obtained from participants of the third survey of the ROAD study. The ROAD study, which began in 2005, is a nationwide prospective study comprising population-based cohorts established in several communities in Japan. Recruitment methods for this study have been described in detail elsewhere [6, 7]. To date, we have created a baseline database including clinical and genetic information of 3040 inhabitants (1061 men; 1979 women) aged 23–95 years who were recruited from listings of resident registrations in three communities. All participants provided written informed consent, and the study was conducted with approval from the ethics committees of the participating institutions.

The third survey of the ROAD study began in 2012 and was completed in 2013. All participants in the baseline study and second survey were invited to participate in the third survey. Besides former participants, inhabitants aged  $\geq 40$  years who were willing to attend the ROAD survey performed in 2012–2013 also were included as participants in the third survey. As a result, a total of 2566 (837 men, 1729 women; urban area, 845 individuals; mountainous area, 769 individuals; coastal area, 952 individuals) residents participated in the third survey.

In the present study, we used data from 1575 subjects (513 men; 1062 women) who completed the stand-up test, two-step test, and 25-question GLFS for disability among all 1721 participants in mountainous and coastal areas in the third survey.

At the third survey, participants completed an interviewer-administered questionnaire. Five interviewers, who had been trained by an expert (NY), were provided for this study. The questionnaire consisted of 200 items that included lifestyle information, such as primary occupation, smoking habits, alcohol consumption, physical activity, medical history, and prescription medication. Anthropometric measurements included height (cm), weight (kg), body mass index [BMI, weight (kg)/height (m)<sup>2</sup>], and hand grip strength (kg).

### Indices for risk of decline in mobility resulting from locomotive syndrome

In the present study, participants performed the following tests for assessment of decline in mobility.

#### *Two-step test*

This test measures the stride length to assess walking ability, including muscle strength, balance, and flexibility of the lower limbs. The two-step test was performed using the following procedure [3, 8, 9]: (1) subjects determined the starting line and stood with the toes of both feet behind it; (2) subjects were instructed to take two long steps (as long as possible) and then align both feet; (3) the length of the two steps from the starting line to the tips of the subject's toes where he/she stopped was measured. The two-step test score was calculated using the following formula: length of the two steps (cm)  $\div$  height (cm).

#### *Stand-up test*

This test assesses leg strength by having the subject stand up on one or both legs from a specified height. After preparation of four seats of different heights—40, 30, 20, and 10 cm—the subject stood up from each seat (in descending height order), first with both legs then with one leg. If the subject could stand up without leaning back to gain momentum and maintain the posture for 3 s, then he/she was diagnosed as having passed that height level [3, 9]. In the present study, if the subject was unable to stand up on one leg (right or left) from a height of 40 cm, then his/her stand-up test was considered failed.

#### *The 25-Question GLFS*

As mentioned above, the 25-question GLFS was developed by Seichi et al. [4]. It is a self-administered, comprehensive measure, consisting of 25 items that include four questions regarding pain during the last month, 16 questions regarding activities of daily living during the last month, three questions regarding social functions, and two questions regarding mental health status during the last month. These 25 items are graded with a five-point scale, from no impairment (0 points) to severe impairment (4 points), and then arithmetically added to produce a total score (minimum = 0, maximum = 100). Thus, a higher score is associated with worse locomotive function. Validity of the scale has been assessed, and a cutoff point of 16 was determined to have the highest sensitivity and specificity for indication of disability resulting from locomotive syndrome [3, 4].

### Indices for decline in mobility resulting from locomotive syndrome

Because the present study utilized a cross-sectional design, not a longitudinal follow-up design, we could not evaluate the ability of the stand-up test or two-step test for prediction of disability resulting from locomotive syndrome. Therefore, in the present study, we used the following outcome measures as indices for decline in mobility.

#### *Five-times-sit-to-stand test*

There are several reports that inability to rise from a chair five times within a determined time is associated with increased disability and morbidity [10–12]. We have also reported that the longer the standing time is, the higher the incidence of disability [13]. The five-times-sit-to-stand test (FTSST) was performed according to the following procedure: (1) using a straight-back chair with a solid seat, participants were asked to sit on the chair with their arms folded across their chest; (2) participants were instructed to stand up and sit down as quickly as possible five times, keeping their arms folded across their chest; (3) the time when the participant stood for the fifth time was measured. In the present study, we used a cutoff value of 12 s to indicate a decline in mobility [14].

#### *The 6-m walking time*

As another outcome measure for decline in mobility, participants walked a 6-m course at their usual speed. The method of measurement of walking time was identical to that performed in the large-scale cohort study entitled Osteoporotic Fractures in Men (MrOS), which started prior to the ROAD study [15]. In the present study, we used a cutoff value of 0.8 m/s to indicate a decline in mobility [16].

In the present study, among the above-mentioned indices, mean scores and SDs for the two-step test were calculated according to participants' sex and age strata (<40, 40s, 50s, 60s, 70s, and  $\geq 80$  years). Then, we estimated the prevalence of each index in stages 1 and 2. Finally, we assessed the association between the cumulative number of indices and decline in mobility using multivariate analysis.

### Statistical analysis

All statistical analyses were performed using STATA statistical software (STATA, College Station, TX, USA). Differences in proportions were compared using the chi-square test. The significance of differences in continuous variables was evaluated using analysis of variance for comparisons

among multiple groups or Scheffe's least significant difference test for pairs of groups. All  $p$  values and 95 % confidence intervals are two-sided. A  $p$  value of  $<0.05$  was considered statistically significant.

Logistic regression analysis was used to test the association of each factor with the presence or absence of a decline in mobility. In the analysis, we used presence of immobility according to the FTSSST time ( $>12$  s = 1;  $\leq 12$  s = 0) and usual walking speed ( $<0.8$  m/s = 1;  $\geq 0.8$  m/s = 0) as the objective variable, and presence or absence of new indices in stages 1 and 2 as explanatory variables, after adjusting for age (+1 year), sex (men = 0, women = 1), BMI (+1 kg/m<sup>2</sup>), and regional difference (mountainous area = 0; coastal area = 1). Other factors were considered in the multivariate model after simple linear analysis; those used as explanatory factors are described in "Results."

## Results

Summary characteristics, including age and BMI of the participants, are shown in Table 1. Two-thirds of the 1575 participants were women, and the mean age of women participants was 1 year less than that of men participants; however, this difference was not significant. By contrast, there was a significant difference in BMI between sexes

( $p < 0.0001$ ). Table 1 also shows the age and sex distributions of mean FTSSST time and walking speed. Both values tended to be significantly slower in participants aged in their 70s and 80s in both men and women, and there were no significant differences between sexes. Table 1 also shows the age and sex distributions of mean two-step test scores. Mean two-step test score was 1.25 (SD 0.20) in men and 1.23 (SD 0.21) in women; this difference was significant ( $p < 0.0001$ ). Age differences indicated that the two-step test score was significantly lower according to age in both men and women ( $p < 0.05$ ).

First, the prevalence of the indices in stage 1 and their association with decline in mobility described by slow FTSSST time and slow walking speed were assessed (Table 2). Overall, the prevalence of two-step test score  $<1.3$ , difficulty with one-leg standing from a 40-cm-high seat in the stand-up test, and 25-question GLFS score  $\geq 7$  were 57.4, 40.6, and 22.6 %, respectively.

Prevalence of two-step test score  $<1.3$  in subjects aged in their 30s and younger, 40s, 50s, 60s, 70s, and 80 years and older were 17.0, 28.5, 32.6, 51.5, 76.2, and 90.0 %, respectively, indicating that the prevalence increased according to age; even in subjects aged in their 40s and 50s, the prevalence was more than 30 %. Prevalence of subjects who could not stand with one leg from a 40-cm-high seat who were aged in their 30s and younger, 40s, 50s,

**Table 1** Mean (SD) values for age, body mass index (BMI), five-times-sit-to-stand test (FTSSST) time, walking speed, and two-step test score of participants classified by age and sex

Age strata (years)	<i>n</i>	Age (years)	BMI (kg/m <sup>2</sup> )	FTSSST time (s)	Walking speed (m/s)	Two-step test score
<b>Men</b>						
<40	23	32.8 (4.8)	24.5 (3.3)	6.96 (1.33)	1.26 (0.22)	1.49 (0.14)
40–49	38	44.7 (3.1)	25.4 (5.1)	6.79 (2.41)	1.25 (0.25)	1.41 (0.15)
50–59	82	55.2 (2.5)	24.2 (3.3)	7.11 (1.47)	1.25 (0.26)	1.36 (0.13) <sup>a</sup>
60–69	137	64.3 (2.7)	23.8 (3.4)	8.10 (2.51)	1.16 (0.26)	1.29 (0.15) <sup>ab</sup>
70–79	139	74.3 (2.8)	23.4 (2.9)	8.72 (2.18) <sup>bc</sup>	1.02 (0.24) <sup>abcd</sup>	1.20 (0.16) <sup>abcd</sup>
$\geq 80$	94	83.8 (3.1)	22.3 (3.0)	11.48 (4.72) <sup>abcde</sup>	0.81 (0.28) <sup>abcde</sup>	1.06 (0.22) <sup>abcde</sup>
Total	513	66.2 (13.7)	23.6 (3.4)	8.57 (3.17)	1.08 (0.30)	1.25 (0.20)
<b>Women</b>						
<40	36	34.4 (4.8)	20.7 (3.0)	7.11 (1.26)	1.28 (0.17)	1.40 (0.14)
40–49	85	44.9 (2.9)	21.9 (3.2)	7.19 (1.64)	1.25 (0.22)	1.35 (0.11)
50–59	195	54.7 (2.9)	23.0 (4.1)	7.10 (1.94)	1.26 (0.22)	1.35 (0.13)
60–69	309	64.7 (2.9)	22.8 (3.4)	7.90 (2.31)	1.18 (0.23) <sup>c</sup>	1.28 (0.18) <sup>abc</sup>
70–79	303	74.3 (2.9)	23.3 (3.3)	9.44 (3.57) <sup>abcd</sup>	1.02 (0.28) <sup>abcd</sup>	1.16 (0.18) <sup>abcd</sup>
$\geq 80$	134	83.1 (3.0)	22.0 (3.4)	11.89 (4.60) <sup>abcde</sup>	0.75 (0.28) <sup>abcde</sup>	0.97 (0.23) <sup>abcde</sup>
Total	1062	65.3 (12.6)	22.7 (3.5)	8.58 (3.31)	1.11 (0.30)	1.23 (0.21)

<sup>a</sup> Significantly different ( $p < 0.05$ ) from values of those aged  $<40$  years

<sup>b</sup> Significantly different ( $p < 0.05$ ) from values of those aged in their 40s

<sup>c</sup> Significantly different ( $p < 0.05$ ) from values of those aged in their 50s

<sup>d</sup> Significantly different ( $p < 0.05$ ) from values of those aged in their 60s

<sup>e</sup> Significantly different ( $p < 0.05$ ) from values of those aged in their 70s

**Table 2** Prevalence of indices in the locomotive syndrome risk test (stage 1): two-step test score <1.3, difficulty with one-leg standing from 40-cm-high seat in the stand-up test (either leg), and 25-ques-tion geriatric locomotive function scale (GLFS) score  $\geq 7$  in participants classified by age and sex

Age strata (years)	Age (years) mean (SD)	Two-step test score <1.3 (%)	Difficulty with one-leg standing from 40-cm-high seat (either leg) (%)	25-question GLFS score $\geq 7$ (%)
<b>Men</b>				
<40	32.8 (4.8)	13.0	4.4	4.4
40–49	44.7 (3.1)	21.1	15.8	10.8
50–59	55.2 (2.5)	34.6	15.9	7.4
60–69	64.3 (2.7)	49.3	30.7	12.0
70–79	74.3 (2.8)	71.7	47.8	19.9
$\geq 80$	83.8 (3.1)	84.6	78.0	44.0
Total	66.2 (13.7)	55.6	39.1	18.8
<b>Women</b>				
<40	34.4 (4.8)	19.4	11.1	0.0
40–49	44.9 (2.9)	31.8	12.9	8.3
50–59	54.7 (2.9)	31.8	23.6	13.0
60–69	64.7 (2.9)	52.4	33.9	19.7
70–79	74.3 (2.9)	78.3	56.2	31.6
$\geq 80$	83.1 (3.0)	93.8	78.1	54.3
Total	65.3 (12.6)	58.2	41.3	24.5*

\* Significantly different ( $p < 0.05$ ) from values of men**Table 3** Effect of presence of indices in the locomotive syndrome risk test (stage 1) for decline in mobility described by slow five-times-sit-to-stand test (FTSST) time and slow walking speed

Indices for decline in mobility	Reference	OR (95 % CI)	<i>p</i> value
<b>FTSST time &gt;12 s</b>			
Two-step test score <1.3	Yes vs. no	3.28 (1.81–5.97)	<0.001
Difficulty with one-leg standing from 40-cm-high seat (either leg)	Yes vs. no	1.78 (1.17–2.69)	0.007
25-question GLFS score $\geq 7$	Yes vs. no	2.51 (1.73–3.62)	<0.001
<b>Walking speed &lt;0.8 m/s</b>			
Two-step test score <1.3	Yes vs. no	4.24 (2.18–8.22)	<0.001
Difficulty with one-leg standing from 40-cm-high seat (either leg)	Yes vs. no	2.01 (1.35–3.16)	0.001
25-question GLFS score $\geq 7$	Yes vs. no	2.65 (1.82–3.86)	<0.001

After adjusting for age, sex, body mass index, and regional difference. Presence or absence of indices for stage 1 also was mutually adjusted  
*CI* confidence interval, *GLFS* geriatric locomotive function scale, *OR* odds ratio

60s, 70s, and 80 years and older were 8.5, 13.8, 21.3, 32.9, 53.5, and 78.1 %, respectively, indicating that the prevalence increased according to age, similar to the two-step test; even in subjects aged in their 40s and 50s, the prevalence was around 20 %. Prevalence of 25-question GLFS score  $\geq 7$  in participants aged in their 30s and younger, 40s, 50s, 60s, 70s, and 80 years and older were 1.7, 9.1, 11.4, 17.4, 27.9, and 50.0 %, respectively, indicating the overall prevalence was lower than that of the other two indices, but it increased synergistically in those in their 80s and older. Regarding the sex difference of the indices in stage 1,

although there were no significant differences between men and women with regard to two-step test score <1.3 and difficulty with one-leg standing from a 40-cm-high seat in the stand-up test, the prevalence of 25-question GLFS score  $\geq 7$  in women was significantly higher than that in men ( $p < 0.05$ ).

Table 3 shows the results of logistic regression analysis using immobility described by slow FTSST time or slow walking speed as the objective factor and the presence or absence of new indices in stage 1 for a decline in mobility as explanatory factors, after adjusting for age (+1 year),

sex (men = 0; women = 1), BMI (+1 kg/m<sup>2</sup>), and regional difference (mountainous area = 0; coastal area = 1). The analysis revealed that all three indices in stage 1 independently predicted immobility described by both slow FTSSST time and slow walking speed.

Table 4 shows the association between accumulation of the indices in stage 1 and decline in mobility described by slow FTSSST time or slow walking speed, after adjusting for age (+1 year), sex (men = 0; women = 1), BMI (+1 kg/m<sup>2</sup>), and regional difference (mountainous area = 0;

**Table 4** Effect of accumulation of indices (stage 1) for decline in mobility

No. of indices	OR (95 % CI)	p value
<b>Five-times-sit-to-stand test time &gt;12 s</b>		
0 = reference	1.0	–
1	2.19 (0.98–4.87)	0.055
2	2.90 (1.30–6.47)	0.009
3	11.59 (5.18–25.93)	<0.001
<b>Walking speed &lt;0.8 m/s</b>		
0 = reference	1.0	–
1	5.73 (1.71–19.16)	0.005
2	9.82 (2.96–32.52)	<0.001
3	32.21 (9.64–107.7)	<0.001

After adjusting for age, sex, body mass index, and regional difference  
CI confidence interval, OR odds ratio

**Table 5** Prevalence of indices in the locomotive syndrome risk test (stage 2): two-step test score <1.1, difficulty with standing from 20-cm-high seat using both legs in the stand-up test, and 25-question

Age strata (years)	Age, mean (SD) years	Two-step test score <1.1 (%)	Difficulty with standing from 20-cm-high seat (%)	25-question GLFS score ≥16 (%)
<b>Men</b>				
<40	32.8 (4.8)	0.0	0.0	0.0
40–49	44.7 (3.1)	2.6	2.6	0.0
50–59	55.2 (2.5)	3.7	0.0	1.2
60–69	64.3 (2.7)	8.8	3.7	6.0
70–79	74.3 (2.8)	23.9	2.9	8.1
≥80	83.8 (3.1)	58.2	16.5	27.5
Total	66.2 (13.7)	20.1	4.9	9.0
<b>Women</b>				
<40	34.4 (4.8)	0.0	0.0	0.0
40–49	44.9 (2.9)	1.2	0.0	2.4
50–59	54.7 (2.9)	3.1	1.0	4.2
60–69	64.7 (2.9)	12.4	5.5	4.6
70–79	74.3 (2.9)	30.4	13.1	15.1
≥80	83.1 (3.0)	71.1	31.3	39.4
Total	65.3 (12.6)	21.6	9.4*	11.4

\* Significantly different (p < 0.01) from values of men

coastal area = 1). The analysis revealed that according to an increasing number of indices, the odds ratio of both slow FTSSST time and slow walking speed increased exponentially.

Next, the prevalence of the indices in stage 2 and their association with decline in mobility described by a slow FTSSST time and slow walking speed were assessed (Table 5). Overall, the prevalence of two-step test score <1.1, difficulty with standing from a 20-cm-high seat using both legs in the stand-up test, and 25-question GLFS score ≥16 were 21.1, 7.9, and 10.6 %, respectively.

Prevalence of two-step test score <1.1 in subjects aged in their 30s and younger, 40s, 50s, 60s, 70s, and 80 years and older were 0.0, 1.6, 3.3, 11.3, 28.4, and 65.8 %, respectively, indicating that the prevalence was less than 5 % in those aged in their 50s and younger, around 10 % in those aged in their 60s, but more than 50 % in those aged 80 years and older. Prevalence of subjects who could not stand from a 20-cm-high seat using both legs who were aged in their 30 s and younger, 40s, 50s, 60s, 70s, and 80 years and older were 0.0, 0.8, 0.7, 5.0, 9.9, and 25.1 %, respectively, indicating that the prevalence was less than 5 % in those aged in their 60s and younger, around 10 % in those aged in their 70s, but dramatically increased in those aged 80 years and older. Prevalence of 25-question GLFS score ≥16 in participants aged in their 30s and younger, 40s, 50s, 60s, 70s, and 80 years and older were 0.0, 1.7, 3.3, 5.0, 12.9, and

geriatric locomotive function scale (GLFS) score ≥16 in participants classified by age and sex

**Table 6** Effect of presence of indices in the locomotive syndrome risk test (stage 2) for decline in mobility described by slow five-times-sit-to-stand test (FTSST) time and slow walking speed

Indices for decline in mobility	Reference	OR (95 % CI)	<i>p</i> value
FTSST time >12 s			
Two-step test score <1.1	Yes vs. no	3.03 (1.97–4.66)	<0.001
Difficulty with standing from 20-cm-high seat %	Yes vs. no	5.87 (3.48–9.89)	<0.001
25-question GLFS score $\geq 16$	Yes vs. no	2.83 (1.77–4.54)	<0.001
Walking speed <0.8 m/s			
Two-step test score <1.1	Yes vs. no	4.19 (2.75–6.39)	<0.001
Difficulty with standing from 20-cm-high seat %	Yes vs. no	3.40 (1.99–5.82)	<0.001
25-question GLFS score $\geq 16$	Yes vs. no	3.49 (2.15–5.65)	<0.001

After adjusting for age, sex, body mass index, and regional difference. Presence or absence of indices for stage 2 also was mutually adjusted  
*CI* confidence interval, *GLFS* geriatric locomotive function scale, *OR* odds ratio

34.4 %, respectively, indicating that the overall prevalence increased according to age; the prevalence was less than 5 % in subjects aged in their 60s and younger, around 10 % in those aged in their 70s, but dramatically increased in those aged 80 years and older. Regarding the sex difference of the indices in stage 2, although there were no significant differences between men and women with regard to two-step test score <1.1 and 25-question GLFS score  $\geq 16$ , the prevalence of difficulty with standing from a 20-cm-high seat using both legs in the stand-up test in women was significantly higher than that in men ( $p < 0.01$ ).

Table 6 shows the results of logistic regression analysis using immobility described by a slow FTSST time or slow walking speed as the objective factor and the presence or absence of new indices in stage 2 for decline in mobility as explanatory factors, after adjusting for age (+1 year), sex (men = 0; women = 1), BMI (+1 kg/m<sup>2</sup>), and regional difference (mountainous area = 0; coastal area = 1). The analysis revealed that all three indices in stage 2 independently predicted immobility described by both a slow FTSST time and slow walking speed. The odds ratios of all indices in stage 2 for slow FTSST time and slow walking speed were greater than those of all indices in stage 1.

Table 7 shows the association between accumulation of the indices in stage 2 and decline in mobility described by slow FTSST time or slow walking speed, after adjusting for age (+1 year), sex (men = 0; women = 1), BMI (+1 kg/m<sup>2</sup>), and regional difference (mountainous area = 0; coastal area = 1). The analysis revealed that according to an increasing number of indices, the odds ratio of both slow FTSST time and slow walking speed increased exponentially. The odds ratios of accumulation of the indices in stage 2 for slow FTSST time and slow walking speed were greater than those of accumulation of the indices in stage 1.

**Table 7** Effect of accumulation of new indices (stage 2) for decline in mobility

No. of new indices	OR (95 % CI)	<i>p</i> value
Five-times-sit-to-stand test time >12 s		
0 = reference	1.0	–
1	2.99 (1.85–4.84)	<0.001
2	12.35 (7.08–21.54)	<0.001
3	46.87 (19.37–113.45)	<0.001
Walking speed <0.8 m/s		
0 = reference	1.0	–
1	3.50 (2.19–5.59)	<0.001
2	12.52 (7.08–22.13)	<0.001
3	61.93 (24.92–153.87)	<0.001

After adjusting for age, sex, body mass index, and regional difference  
*CI* confidence interval, *OR* odds ratio

## Discussion

In the present study, we clarified the associations of three new indices for immobility, including the two-step test, stand-up test, and 25-question GLFS, represented by slow FTSST time and slow walking speed. We first tested the associations among the three indices by using the clinical decision limits for locomotive syndrome risk test stage 1. Next, we tested the three indices in stage 2. We clarified the age and sex distributions of the prevalence of these three indices and found that the three indices in both stages 1 and 2 significantly and independently predicted a decline in mobility and that the accumulation of such indices increased the risk of immobility exponentially.

First, we used both FTSST time and 6-m walking speed as outcome measures of immobility. As mentioned in “Methods”, these two indices are both regarded as predictors for morbidity and disability [10–12, 16]. In the ROAD

study, we reported that the longer the standing time, the higher the incidence of disability [13] and that slow walking speed was also a predictor for the occurrence of disability [13]. Although we could not clarify the direct associations between these new indices and occurrence of disability because this study design was cross-sectional, these surrogate indices could help predict disability in the near future. Therefore, if we could clarify the significant associations between these new indices and FTSST time and 6-m walking speed, we might clarify the ability to predict disability indirectly. Based on this hypothesis, we used cutoff values of 12 s for FTSST time [13] and 0.8 m/s for 6-m walking speed [16].

The two-step test was developed to assess walking ability, including muscle strength, balance, and flexibility of the lower limbs. This test was first developed by Muranaga and Hirano in 2003 [3, 8]. They compared two-step test scores of 108 healthy volunteers (38 men, 70 women; mean age 59.0 years) with those of 108 hospital outpatients (56 men, 52 women; mean age 60.3 years) and found that the two-step test score was significantly associated with the risk of falling and degree of independence in daily life [7]. In the present study, we clarified mean two-step test scores in participants classified by age and sex and found mean scores of 2.07 in men and 1.87 in women. Scores of men were significantly higher than those of women, and age differences indicated that scores were significantly lower according to age in both men and women. These age and sex tendencies are consistent with those reported in a previous study [3].

Regarding the prevalence of the indices in locomotive syndrome risk test stages 1 and 2, we found that all prevalences increased with age. However, the distribution of each index seemed to differ. In stage 1, the prevalence was highest for a two-step test score  $<1.3$ , followed by difficulty with one-leg standing from a 40-cm-high seat in the stand-up test and 25-question GLFS score  $\geq 7$ . By contrast, in stage 2, the prevalence also was highest for a two-step test score  $<1.1$ , but the prevalence of a 25-question GLFS score  $\geq 16$  was higher than that of difficulty with standing from a 20-cm-high seat using both legs in the stand-up test. The different age distributions of these three indices in both stages 1 and 2 might be conducive to their mutually independent associations with immobility. Our result that these three indices in both stages 1 and 2 independently predicted immobility shows that all three are important for predicting immobility. Furthermore, because these indices independently predicted immobility, risk severity may be classified by a positive number of indices present. In fact, in our study, accumulation of indices increased the risk of immobility exponentially, especially accumulation of indices in stage 2, which suggests the possibility of categorizing the

severity of risk for immobility, such as 0, normal; 1, mild; 2, moderate; 3, severe.

With regard to the 25-question GLFS, there might be some concern that it is too cumbersome for older people to answer 25 questions. Seichi et al. also proposed a short version of the test using only five questions with a cutoff score of  $\geq 6$ , on behalf of a 25-question GLFS score  $\geq 16$ . They selected five questions from the 25-question GLFS using the Akaike information criterion (AIC) [17–19]. They determined that a cutoff score of 6 had the lowest AIC value, representing the model with the best fit, and finally concluded that the five-question GLFS can be applied as a rapid self-check tool for locomotive syndrome [4]. We compared the results of 1535 individuals who completed both the 25-question GLFS and five-question GLFS. Supplementary Table 1 shows that these two indices (five-question GLFS and 25-question GLFS) have a strong association (sensitivity 0.859; specificity 0.985). This result shows the possibility of using the five-question GLFS instead of the 25-question GLFS as a rapid-check tool for the prediction of immobility.

There are several limitations in the present study. First, as mentioned above, our study assessed the usefulness of three indices in locomotive syndrome risk test stages 1 and 2 for predicting immobility using FTSST time and walking speed as outcome variables. Although there has been significant evidence regarding these outcome measures, such as slow FTSST time and slow walking speed, and disability [8, 10–13, 16], including our report, the direct associations of these new indices and occurrence of disability are unclear. The proposal of these new indices was published in 2013 [3], so there was not enough time to observe future occurrence of disability in our cohort. Therefore, we should continue to observe our cohort and assess the ability of such indices to predict disability directly. Second, although the ROAD study includes a large number of participants, these participants do not truly represent the general population, since the subjects in the present study were recruited from only two areas. However, we have already confirmed the representativeness of the participants of the ROAD baseline study to the Japanese population by comparing anthropometric measurements and frequencies of smoking and alcohol consumption between participants and the general Japanese population [5]. Values for the general population were obtained from the 2005 National Health and Nutrition Survey conducted by the Ministry of Health, Labour and Welfare in Japan [20]. Regarding anthropometric measurements, we found no significant differences between our participants and the total Japanese population, except that men participants aged 70–74 years in the ROAD study were significantly smaller in terms of body structure than the overall population ( $p < 0.05$ ). In addition, proportions of current

smokers and current drinkers were significantly higher in the general Japanese population than in the study population [5], suggesting that participants of the ROAD study have healthier lifestyles than the general population. This “healthy” selection bias should be taken into consideration when using reference values obtained in the ROAD study.

In conclusion, we have assessed whether the proposed clinical decision limits of the indices in locomotive syndrome risk test stages 1 and 2 could predict immobility represented by a slow FTSST time and slow walking speed using data from a population-based cohort of the third survey of the ROAD study. We found that all the indices in the locomotive syndrome risk test—the two-step test, stand-up test, and 25-question GLFS—could significantly and independently predict a decline in mobility and that the accumulation of such indices increased the risk of immobility exponentially.

**Acknowledgments** This work was supported by a Grant-in-Aid for H17-Men-eki-009 (Director, Kozo Nakamura), H20-Chouju-009 (Director, Noriko Yoshimura), H23-Chouju-002 (Director, Toru Akune), H-25-Chouju-007 (Director, Noriko Yoshimura), and H25-Nanchitou(Men)-005 (Director, Sakae Tanaka) of the Ministry of Health, Labour, and Welfare; and Scientific Research B23390172, B20390182, and Challenging Exploratory Research 24659317 to Noriko Yoshimura; B23390356, C20591774, and Challenging Exploratory Research 23659580 to Shigeyuki Muraki; Challenging Exploratory Research 24659666 and 21659349 and Young Scientists A18689031 to Hiroyuki Oka; B23390357 and C20591737 to Toru Akune; Collaborating Research with NSF 08033011-00262 (Director, Noriko Yoshimura) from the Ministry of Education, Culture, Sports, Science and Technology in Japan. This study also was supported by grants from the Japan Osteoporosis Society (Noriko Yoshimura, Shigeyuki Muraki, Hiroyuki Oka, and Toru Akune) and research aid from the Japanese Orthopaedic Association (JOA-Subsidized Science Project Research 2006-1 & 2010-2; Director, Hiroshi Kawaguchi). The authors wish to thank Dr. Takako Nojiri and Mr. Kazuhiro Hatanaka of the Gobo Public Health Centre; Dr. Naoki Hirabayashi of the Kawakami Clinic in Hidakagawa Town; Mrs. Tomoko Takijiri, Mrs. Rie Takiguchi, Mrs. Kyoko Maeda, and other members of the public office in Hidakagawa Town; Dr. Shinji Matsuda of the Shingu Public Health Centre; Mrs. Tamako Tsutsumi, Mrs. Kanami Maeda, Mrs. Megumi Takino, Mrs. Shuko Okada, Mrs. Kazuyo Setoh, Mrs. Chise Ryouno, Mrs. Miki Shimosaki, Mrs. Chika Yamaguchi, Mrs. Yuki Shimoji, and other members of the public office in Taiji Town for their assistance in locating and scheduling participants for examinations. We also thank Ms. Kyoko Yoshimura, Mrs. Toki Sakurai, Mrs. Saeko Sahara, and Mr. Noriyuki Oe for their assistance in data reduction and administration.

**Conflict of interest** No conflicts of interest have been declared by the authors.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

1. Ministry of Health, Labour and Welfare. The outline of the results of National Livelihood Survey (2013). <http://www.mhlw.go.jp/toukei/saikin/hw/k-tyosa/k-tyosa13/dl/06.pdf>. Accessed 17 June 2015
2. Nakamura K. A “super-aged” society and the “locomotive syndrome”. *J Orthop Sci.* 2008;13(1):1–2.
3. Locomotive syndrome. In: Locomotive Challenge! Council, editors. Locomotive syndrome pamphlet 2013. Japanese Orthopaedic Association: Tokyo; 2013.
4. Seichi A, Hoshino Y, Doi T, Akai M, Tobimatsu Y, Iwaya T. Development of a screening tool for risk of locomotive syndrome in the elderly: the 25-question Geriatric Locomotive Function Scale. *J Orthop Sci.* 2012;17(2):163–72.
5. Locomotive syndrome. In: Locomotive Challenge! Council, editors. Locomotive syndrome pamphlet 2015. Tokyo: Japanese Orthopaedic Association; 2015 (in press).
6. Yoshimura N, Muraki S, Oka H, Mabuchi A, En-yo Y, Yoshida M, Saika A, Suzuki T, Yoshida H, Kawaguchi H, Nakamura K, Akune T. Prevalence of knee osteoarthritis, lumbar spondylosis and osteoporosis in Japanese men and women: the research on osteoarthritis/osteoporosis against disability study. *J Bone Miner Metab.* 2009;27(5):620–8.
7. Yoshimura N, Muraki S, Oka H, Kawaguchi H, Nakamura K, Akune T. Cohort profile: research on osteoarthritis/osteoporosis against disability (road) study. *Int J Epidemiol.* 2010;39(4):988–95.
8. Muranaga S, Hirano K. Development of a convenient way to predict ability to walk, using a two-step test. *J Showa Med Assoc.* 2003;63(3):301–8 (in Japanese).
9. Muranaga S. Evaluation of the muscular strength of the lower extremities using the standing movement and clinical application. *J Showa Med Assoc.* 2001;61(3):362–7 (in Japanese).
10. Guralnik JM, Ferrucci L, Pieper CF, Leveille SG, Markides KS, Ostir GV, Studenski S, Berkman LF. Lower extremity function and subsequent disability: consistency across studies, predictive models, and value of gait speed alone compared with the short physical performance battery. *J Gerontol A Biol Sci Med Sci.* 2000;55(4):M221–31.
11. Buatois S, Miljkovic D, Manckoundia P, Gueguen R, Miget P, Vançon G, Perrin P, Benetos A. Five times sit to stand test is a predictor of recurrent falls in healthy community-living subjects aged 65 and older. *J Am Geriatr Soc.* 2008;56(8):1575–7.
12. Bohannon RW. Reference values for the five-repetition sit-to-stand test: a descriptive meta-analysis of data from elders. *Percept Mot Skills.* 2006;103(1):215–22.
13. Akune T, Muraki S, Oka H, Tanaka S, Kawaguchi H, Tokimura F, Yoshida H, Suzuki T, Nakamura K, Yoshimura N. Incidence of certified need of care in the long-term care insurance system and its risk factors in the elderly of Japanese population-based cohorts: the ROAD study. *Geriatr Gerontol Int.* 2014;14(3):695–701.
14. Mong Y, Teo TW, Ng SS. 5-repetition sit-to-stand test in subjects with chronic stroke: reliability and validity. *Arch Phys Med Rehabil.* 2010;91(3):407–13.
15. Orwoll E, Blank JB, Barrett-Connor E, Cauley J, Cummings S, Ensrud K, Lewis C, Cawthon PM, Marcus R, Marshall LM, McGowan J, Phipps K, Sherman S, Stefanick ML, Stone K. Design and baseline characteristics of the osteoporotic fractures in men (MrOS) study—a large observational study of the determinants of fracture in older men. *Contemp Clin Trials.* 2005;26(5):569–85.
16. Chen LK, Liu LK, Woo J, Assantachai P, Auyeung TW, Bahyah KS, Chou MY, Chen LY, Hsu PS, Krairit O, Lee JS, Lee WJ,

- Lee Y, Liang CK, Limpawattana P, Lin CS, Peng LN, Satake S, Suzuki T, Won CW, Wu CH, Wu SN, Zhang T, Zeng P, Akishita M, Arai H. Sarcopenia in Asia: consensus report of the Asian Working Group for Sarcopenia. *J Am Med Dir Assoc*. 2014;15(2):95–101.
17. Akaike H. Information theory and an extension of the maximum likelihood principles. In: Petrov B, Caski F, editors. *The second international symposium on information theory*. Budapest: Akademia Kiado; 1973. p. 267–81.
  18. Sakamoto Y, Ishiguro M, Kitagawa G. *Akaike information criterion statistics*. Dordrecht: Reidel; 1986.
  19. FORTRAN Program CATDAP-02. In: Sakamoto Y, ed. *Categorical data analysis by AIC*. Tokyo: KTK Scientific Publishers, Kluwer Academic Publishers; 1991: 163–206.
  20. Ministry of Health, Labour and Welfare. *The report of National Health and Nutrition Survey (2005)*. <http://www.mhlw.go.jp/bunya/kenkou/eiyou07/dl/01-04.pdf>. Accessed 17 June 2015