

Erratum to: Individual and peer factors associated with ketamine use among adolescents in Taiwan

Kun-Hua Lee¹ · Yi-Chun Yeh^{2,3} · Pin-Chen Yang^{2,3} ·
Huang-Chi Lin^{2,3} · Peng-Wei Wang^{2,3} · Tai-Ling Liu^{2,3} ·
Cheng-Fang Yen^{2,3}

Published online: 8 March 2017
© Springer-Verlag Berlin Heidelberg 2017

Erratum to: Eur Child Adolesc Psychiatry
DOI 10.1007/s00787-012-0292-7

In the original publication, the scoring method of the *Negative outcome expectancy for ketamine* is incorrectly described. The correct version is as follows:

“The response categories included: 1, very severe damage; 2, severe damage; 3, a little damage; and 4, no damage.”

The online version of the original article can be found under
doi:[10.1007/s00787-012-0292-7](https://doi.org/10.1007/s00787-012-0292-7).

✉ Cheng-Fang Yen
chfaye@cc.kmu.edu.tw

- ¹ Institute of Allied Health Sciences, College of Medicine, National Cheng Kung University, Tainan, Taiwan
- ² Department of Psychiatry, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan
- ³ Department of Psychiatry, Faculty of Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan