



Correction to: Cavin-1 is linked to lipid droplet formation in human hepatic stellate cells

Hiroaki Yokomori¹ · Wataru Ando² · Masaya Oda³

Published online: 13 May 2020

© The Japanese Society for Clinical Molecular Morphology 2020

Correction to:

Medical Molecular Morphology (2020) 53:56–59

<https://doi.org/10.1007/s00795-019-00219-4>

In the original publication, the part figures (e, f) of Fig. 2 were wrongly cited as “g, h” in the text and in Fig. 2 caption.

The last sentence in the second paragraph of page 56 should read as “By contrast, particles labeling cavin-1 were present on caveolae of plasma membranes, but only rarely on LD and ER-like membranes (Fig. 2e, f).

The first sentence in the Fig. 2 legend should read as “Transmission electron micrographs of cavin-1 (a–d) showing hepatic stellate cells (HSCs) around the portal tract (a, b)

and around central vein in normal liver tissue (a–d) and in cirrhotic liver tissue (e, f): a, c, e low magnification; b, d, f high magnification. a, b HSCs associated with low-density lipid droplets (LDs).”

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00795-019-00219-4>.

✉ Hiroaki Yokomori
yokomori@insti.kitasato-u.ac.jp

¹ Department of Internal Medicine, Kitasato University Medical Center, Saitama 364-8501, Japan

² Department of Clinical Pharmacy, School of Pharmacy, Kitasato University, Tokyo, Japan

³ Organized Center of Clinical Medicine, International University of Health and Welfare, Tokyo, Japan