CORRECTION



Correction: Resistance to FGFR1-targeted therapy leads to autophagy via TAK1/AMPK activation in gastric cancer

Rui Peng¹ · Yan Chen^{2,5} · Liangnian Wei³ · Gang Li¹ · Dongju Feng³ · Siru Liu² · Runqiu Jiang⁴ · Shaojiang Zheng² · Yun Chen^{1,3,6}

Published online: 21 March 2024 © The International Gastric Cancer Association and The Japanese Gastric Cancer Association 2024

Correction: Gastric Cancer (2020) 23:988–1002 https://doi.org/10.1007/s10120-020-01088-y.

In Fig. 4a and 5a of this article, the representative western blot images of mTOR (in Fig. 4a) and β -actin of BGC-823 cells (in Fig. 5a) were mistakenly inserted; the Fig. 4a and 5a should have appeared as shown below.

The original article can be found online at https://doi.org/10.1007/s10120-020-01088-y.

- Runqiu Jiang jiangrq@njmu.edu.cn
- Shaojiang Zheng zhenghoho@aliyun.com
- Yun Chen chenyun@njmu.edu.cn
- ¹ Department of General Surgery, Research Center for Clinical Oncology, The Affiliated Cancer Hospital of Nanjing Medical University, Jiangsu Cancer Hospital, Jiangsu Institute of Cancer Research, Nanjing 210009, Jiangsu, China
- ² Key Laboratory of Emergency and Trauma of Ministry of Education, Tumor Institute of the First Affiliated Hospital, Hainan Medical University, Haikou 571199, China

- ³ Department of Immunology, Key Laboratory of Immune Microenvironment and Disease, Nanjing Medical University, Nanjing 211166, Jiangsu, China
- ⁴ Department of Hepatobiliary Surgery, The Affiliated Drum Tower Hospital of Nanjing University Medical School, Nanjing 210093, Jiangsu, China
- ⁵ Department of Ultrasound, Hainan General Hospital (Hainan Affiliated Hospital of Hainan Medical University), Haikou 570311, China
- ⁶ Jiangsu Key Lab of Cancer Biomarkers, Prevention and Treatment, Collaborative Innovation Center for Cancer Personalized Medicine, Nanjing Medical University, Nanjing 211166, Jiangsu, China




Fig. 4a MGC-803 and BGC-823 cells, as well as resistant cells, were treated with 1 µM AZD4547 for 24 h and immunoblotted with LC3, p62, p-AMPK, total AMPK, p-mTOR, total mTOR, p-S6, total S6, and GAPDH.




Fig. 5a BGC-823 and BGC-823R and MGC-803 and MGC-803R cells were untreated or treated with 1 μ M AZD4547 for 24 h. Protein levels of p-ERK, ERK, and β -actin were measured by western blot assays.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.