

Erratum to: The role of a low glomerular density and being overweight in the etiology of proteinuria in CKD patients without known glomerular diseases

Hidekazu Okamoto · Tetsuya Kawamura ·
Hideo Okonogi · Nobuo Tsuboi · Yoichi Miyazaki ·
Takashi Yokoo

Published online: 4 April 2014
© Japanese Society of Nephrology 2014

Erratum to: Clin Exp Nephrol
DOI 10.1007/s10157-014-0940-y

Unfortunately, there was an error in the article cited above.

In the Subjects and methods section, under the heading “Items included in the clinical examination”, the third sentence should read:

The estimated glomerular filtration rate (eGFR) was calculated as follows: $194 \times \text{serum Cr level}^{-1.094} \times \text{age}^{-0.287}$ (female = $\times 0.739$) [16].

The online version of the original article can be found under doi:[10.1007/s10157-014-0940-y](https://doi.org/10.1007/s10157-014-0940-y).

H. Okamoto · T. Kawamura (✉) · H. Okonogi · N. Tsuboi ·
Y. Miyazaki · T. Yokoo
Division of Kidney and Hypertension, Department of Internal
Medicine, The Jikei University School of Medicine,
3-25-8 Nishi-shinbashi, Minato-ku, Tokyo 105-8461, Japan
e-mail: kawatetu@coral.ocn.ne.jp