CORRECTION



Correction to: Effective clearance of uremic toxins using functionalised silicon Nanoporous membranes

Ananya Ghosh $^1 \cdot$ Fidal Vallam Thodi $^2 \cdot$ Sudeshna Sengupta $^2 \cdot$ Sivasundari Kannan $^1 \cdot$ Lalitha Krishnan $^2 \cdot$ Enakshi Bhattacharya 1,2

Published online: 21 December 2021
© Springer Science+Business Media, LLC, part of Springer Nature 2021

Correction to: Biomedical Microdevices (2021) 23: 4 https://doi.org/10.1007/s10544-020-00539-8

The original version of this article unfortunately contained some mistakes. There is an error in the hemodialysis buffer composition mentioned in Sect. 2: Materials and methods, Subsection 2.1 Materials (Page No.4, Line No. 22–25). The correct composition is as follows:

Hemodialysis (HD) buffer were prepared with following components (Sodium Chloride $-5.31~{\rm g\,L^{-1}}$, Calcium Chloride $-0.22~{\rm g\,L^{-1}}$, Potassium Chloride $-0.15~{\rm g\,L^{-1}}$, Magnesium chloride $-0.10~{\rm g\,L^{-1}}$, Sodium acetate $-0.04~{\rm g\,L^{-1}}$, Citric acid $-0.15~{\rm g\,L^{-1}}$, Sodium bicarbonate $-1.63~{\rm g\,L^{-1}}$).

The caption of Fig. 6 is erroneous and the corrected caption is as follows:

Fig. 6. (A) Initial urea concentration in Cis reservoir; Concentration of urea in the Cis and Trans reservoirs after 2 h of dialysis through (B) Plasma treated SNM (C) GLYMO treated SNM (D) One side silanized acid treated SNM (E) Both sides silanized and acid treated SNM performed using The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at https://doi.org/10.1007/ $\,$ s10544-020-00539-8.



Enakshi Bhattacharya enakshi@ee.iitm.ac.in

Department of Electrical Engineering, Indian Institute of Technology-Madras, Chennai 600036, India

² Centre for NEMS and Nanophotonics, Indian Institute of Technology-Madras, Chennai 600036, India